HFT session homework 4

1) In this problem you will use the simple binomial approximation of Brownian motion to help understand Girsanov's theorem. Consider a simple binary random variable

$$X = \begin{cases} \sqrt{dt} & \text{with probability } 0.5 + \mu(dt)^n/2 \\ -\sqrt{dt} & \text{with probability } 0.5 - \mu(dt)^n/2 \end{cases}$$

- (a) For this Brownian motion with drift to have a drift term of order dt, what must n be?
- (Hint: Compute the mean of X and choose n so that the mean is of order dt.)
- (b) Given that you computed n corrected from part (a), compute the variance of X.
- (c) Can you use the results of (a) and (b) to argue that an equivalent change of measure will change the mean of a Brownian motion, but not the instantaneous variance?

$$\chi = \begin{cases} \delta t^{1/4}, & P = \frac{1}{2} + \frac{M}{2} (dt)^{1/4} \\ - \delta t^{1/4}, & P = \frac{1}{2} - \frac{M}{2} (dt)^{1/4} \end{cases}$$

(a)
$$E[X] = 4t^{1/2}(\frac{1}{2} + \frac{2}{4} dt^{n}) - dt^{1/2}(\frac{1}{2} - \frac{2}{4} dt^{n})$$

$$= \det \left(\frac{1}{2} + \frac{1}{2} \det^{1/2}\right) + \det \left(\frac{1}{2} - \frac{1}{2} \det^{1/2}\right) - \lambda^{2} dt^{2}$$

$$= \det \left(-\lambda^{2} dt^{2}\right) + \det \left(\frac{1}{2} - \frac{1}{2} dt^{2}\right) - \lambda^{2} dt^{2}$$

$$= \det \left(-\lambda^{2} dt^{2}\right)$$

Change
$$M \rightarrow$$
 measure IP changes (to IP)

 \longrightarrow probability variable X changes (to ZX)

 \longrightarrow Brownian motion B changes (to B)

 \longrightarrow But, let $E[\Delta B]$ changes P

$$E[B] \neq E[\widehat{B}]$$
: mean of BM changes

Using principles from linear pricing (or risk-neutrality), derive a PDE for the price of a European call option on a non-dividend paying stock when interest rates are random, and the short rate follows Cox-Ingersoll-Ross dynamics. That is:

where z_1 and z_2 are correlated Winner processes with correlation coefficient ρ . (i.e. $E(dz_1 dz_2) = \rho dt$) (Hint: Your answer may contain a market price of risk.)

let C(S, r,t) be the price of derivative on S. That means,

$$dc = rcdt + (?)d\tilde{\epsilon}_1 + (?)d\tilde{\epsilon}_2$$
, (?) term is not important. We will any care the drift term.

By Ito's formula,

Thus