

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/286449572>

Zoonotic Viruses and Conservation of Bats

Chapter · January 2016

DOI: 10.1007/978-3-319-25220-9_10

CITATIONS

28

READS

443

2 authors:



Karin Schneeberger
Universität Potsdam

21 PUBLICATIONS 402 CITATIONS

[SEE PROFILE](#)



Christian C Voigt
Leibniz Institute for Zoo and Wildlife Research

353 PUBLICATIONS 7,292 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Immunology of European Hibernating Bats [View project](#)



Spatiotemporal effects on bat activity above intensively managed farmland [View project](#)

Chapter 10

Zoonotic Viruses and Conservation of Bats

Karin Schneeberger and Christian C. Voigt

Abstract Many of the recently emerging highly virulent zoonotic diseases have a likely bat origin, for example Hendra, Nipah, Ebola and diseases caused by coronaviruses. Presumably because of their long history of coevolution, most of these viruses remain subclinical in bats, but have the potential to cause severe illnesses in domestic and wildlife animals and also humans. Spillovers from bats to humans either happen directly (via contact with infected bats) or indirectly (via intermediate hosts such as domestic or wildlife animals, by consuming food items contaminated by saliva, faeces or urine of bats, or via other environmental sources). Increasing numbers of breakouts of zoonotic viral diseases among humans and livestock have mainly been accounted to human encroachment into natural habitat, as well as agricultural intensification, deforestation and bushmeat consumption. Persecution of bats, including the destruction of their roosts and culling of whole colonies, has led not only to declines of protected bat species, but also to an increase in virus prevalence in some of these populations. Educational efforts are needed in order to prevent future spillovers of bat-borne viruses to humans and livestock, and to further protect bats from unnecessary and counterproductive culling.

K. Schneeberger (✉) · C.C. Voigt
Department of Evolutionary Ecology, Leibniz Institute for Zoo and Wildlife Research,
Berlin, Germany
e-mail: schneeberger@izw-berlin.de

10.1 Introduction

Over the past decades, the emergence of zoonotic viruses (those that are naturally transmitted between vertebrate animals and humans) from bats has been the subject of increasing attention from both scientists and the general public (e.g. Quammen 2013). During outbreaks of diseases in humans and livestock, bats are now often the primary focus of searches for a reservoir host (Chua et al. 2002a; Leroy et al. 2005; Li et al. 2005; Halpin et al. 2007; Towner et al. 2007; Lau et al. 2010; Wibbelt et al. 2010; Memish et al. 2013). Identification of bats as natural hosts for emerging viruses has important implications for bat conservation. We review the current state of research of four important families of emerging zoonotic viruses for which bats are natural reservoir hosts and discuss direct and indirect conservation implications.

10.2 Emerging Viral Diseases: Why Bats?

Although bats have been identified as carriers of many highly virulent human pathogens (Chen et al. 2014), evidence of pathogen-related clinical signs or disease in bats is scarce, particularly for intracellular pathogens such as viruses (Brook and Dobson 2015). Post-infection survival is supported by the frequent identification of antibodies to known viruses in apparently healthy bats and long-term survival of these bats (e.g. Hayman et al. 2010). Additionally, viruses isolated or genetically detected from bat populations are highly diverse and often ancestral to related viruses in human and other mammalian species (e.g. Towner et al. 2009; Drexler et al. 2012; Baker et al. 2013a; Tong et al. 2013; Vidgen et al. 2015). Together, these findings suggest a long history of coevolution between many bat-virus relationships identified to date. Recent progress in the field of bat immunology and genomics has identified key differences in bat immunity and physiology that evolved concomitantly with the evolution of flight, resulting not only in apparently increased immunotolerance of intracellular pathogens, but also in increased longevity and decreased tumour production (Baker et al. 2013b; Zhang et al. 2013; Brook and Dobson 2015). Immunotolerance and incomplete clearance of viral infections are also likely to favour the establishment of persistent infections (Virgin et al. 2009), as proposed for a number of bat-borne viruses (Plowright et al. 2015).

Various ecological and life-history factors play a key role in the susceptibility of individuals and populations to pathogens (Allen et al. 2009; Turmelle et al. 2010; Schneeberger et al. 2013), and notable differences exist between bats and terrestrial mammals such as rodents (Luis et al. 2013). For example, the often high-population densities and the usually gregarious roosting behaviour of bats increase the likelihood of both intra- and interspecies transmission of viruses (Luis et al. 2013; Streicker et al. 2010). Large-scale movements of bats due to

their ability for powered flight are also likely to facilitate viral transmission within and among species, including the exchange of novel viruses and virus variants across biomes or even continents (Calisher et al. 2006; Epstein et al. 2009; Peel et al. 2013). The extreme relative longevity of bats compared to other mammals of similar size (Wilkinson and South 2002) and the potential for persistent and/or subclinical viral infections could further increase transmission potential (Calisher et al. 2006). Reduction of body temperature associated with hibernation of temperate zone bats lowers both viral activity and the metabolism of hosts, leading to increased incubation periods and therefore reduced likelihood of epizootic fadeout (of rabies, for example; George et al. 2011). Bats are ancient mammals in evolutionary terms, and virus utilisation of highly conserved cellular receptors could facilitate transmission to other mammals (Calisher et al. 2006), for example, as has been suggested for henipaviruses (Negrete et al. 2005). Lastly, it was recently speculated that, similar to the febrile response of other mammals, the relatively high body temperature (about 38–41 °C) and metabolism of bats during flight may select for viruses tolerant to such conditions, meaning the normal febrile defence mechanism of other mammals is ineffective (“Flight as fever hypothesis”, O’Shea et al. 2014), making bat-borne viruses potentially more virulent and lethal for other, non-flying mammals.

10.3 Zoonotic Viruses of Bats and Their SpillOver

10.3.1 *Rhabdoviruses*

Rabies virus (RABV) is the longest and best-known member of the genus *Lyssavirus* (family Rhabdoviridae) and still one of the most significant zoonoses known from bats (recent reviews include: Banyard et al. 2011; Banyard et al. 2014 and Kuzmin 2014). The genus is rapidly expanding, with 14 of the currently recognised species (plus another known from genetic material only), and all but two (Mokola and Ikoma viruses) having been isolated from bats (Table 10.1). Lyssaviruses spill over directly from bats to domestic animals, other wildlife and humans, or indirectly to humans via these other species. All lyssaviruses are potentially neurotropic, meaning that the virus infects nerve cells and replicates in the brain, resulting in clinical signs consistent with classical rabies (Schnell et al. 2009). Although isolated from a variety of tissues and body fluids in the late stages of infection, the predominant route of transmission is via saliva (mostly via biting; Kuzmin 2014).

Lyssaviruses can be divided into two distinct “phylogroups” (Badrane et al. 2001, Table 10.1), reflecting biological and genetic differences, and they are distributed globally in bats. Classical rabies virus occurs in bats across North, Central and South America (Messenger et al. 2003; Banyard et al. 2011) and was first associated with vampire bats following an outbreak in cattle in South America in 1911 (Carini 1911). It is reported most frequently in the common

Table 10.1 Known lyssaviruses and their association with different bat species (adapted from Banyard et al. 2014)

Geographical distribution	Lyssavirus species	Phylogroup	Bat species most commonly associated with lyssavirus infection	Common name	Known human cases
The Americas	Rabies virus (RABV)	I	<i>Eptesicus fuscus</i>	Big brown bat	Yes
			<i>Tadarida brasiliensis</i>	Mexican/ Brazilian free-tail bat	
			<i>Lasionycteris noctivagans</i>	Silver-haired bat	
			<i>Perimyotis subflavus</i>	Tri-coloured bat	
			<i>Desmodus rotundus</i>	Vampire bat	
Eurasia	European bat lyssavirus type 1 (EBLV-1)	I	<i>Eptesicus serotinus</i>	Serotine bat	Yes
	European bat lyssavirus type 2 (EBLV-2)	I	<i>Myotis daubentonii</i>	Daubenton's bat	Yes
	Bokeloh bat lyssavirus (BBLV)	I	<i>Myotis nattereri</i>	Natterer's bat	No
	Aravan virus (ARAV)	I	<i>Myotis blythi</i>	Lesser mouse-eared bat	No
	Irkut virus (IRKV)	I	<i>Murina leucogaster</i>	Greater tube-nosed bat	Yes
	Khujand virus (KHUV)	I	<i>Myotis mystacinus</i>	Whiskered bat	No
	West Caucasian bat virus (WCBV)	NA ^a	<i>Miniopterus schreibersii</i>	Common bent-wing bat	No
	Lleida bat lyssavirus (LLEBV)	NA ^a	<i>Miniopterus schreibersii</i>	Common bent-wing bat	No
Africa	Duvenhage virus (DUVV)	I	<i>Miniopterus sp?</i>	Undefined	Yes
			<i>Nycteris thebaica</i>	Egyptian slit-faced bat	
	Lagos bat virus (LBV)	II	<i>Eidolon helvum</i>	Straw-coloured fruit bat	No
			<i>Rousettus aegyptiacus</i>	Egyptian fruit bat	
			<i>Epomorphorus wahlbergi</i>	Wahlberg's epauletted fruit bat	

(continued)

Table 10.1 (continued)

Geographical distribution	Lyssavirus species	Phylogroup	Bat species most commonly associated with lyssavirus infection	Common name	Known human cases
	Mokola virus (MOKV)	II	not detected		Yes
	Shimoni bat virus (SHIBV)	II	<i>Hipposideros commersoni</i>	Commerson's leaf-nosed bat	No
	Ikoma virus (IKOV)	NA ^a	not detected		No
Australasia	Australian bat lyssavirus (ABLV)	I	<i>Pteropus scapulatus</i> ^b	Little red flying fox	Yes
			<i>Saccolaimus flaviventris</i>	Yellow-bellied sheath-tailed bat	
			<i>Pteropus alecto</i>	Black flying fox	

^aLyssaviral phylogenies infer WCBV, LLEBV and IKOV that are more genetically distinct from other species, and they have not yet been assigned a phylogroup (Kuzmin 2014)

^bBarrett (2004)

vampire bat (*Desmodus rotundus*; Kuzmin et al. 2011a), which has a wide distribution across Mexico, Central America, and South America. Bites from this species appear to be responsible for the majority of human and domestic animal rabies infections of bat origin in South and Central America, with increased prey availability via expansion of livestock into new areas across the region hypothesised to be contributing to increasing incidences (Schneider et al. 2009; Ruiz and Chávez 2010). In Canada and the USA, 51 cases of human rabies transmitted by non-haematophagous bats were recognised or inferred between 1951 and 2006 (mostly silver-haired bats (*Lasionycteris noctivagans*), eastern pipistrelle bats (*Perimyotis subflavus*) and Brazilian/Mexican free-tailed bats (*Tadarida brasiliensis*)) (Constantine and Blehert 2009; Banyard et al. 2011). However, across the Americas, only 15 % of human rabies cases between 1993 and 2002 were reported as resulting from encounters with bats (Belotto et al. 2005).

Reported antibody prevalences against RABV in *D. rotundus* include 3–28 % in Peru (Streicker et al. 2012) and 12 % in Brazil (Almeida et al. 2011). Depending on the year, location and species, prevalence in other bats varies from relatively low 2 % in *T. brasiliensis* in New Mexico (Steece and Altenbach 1989) and 2.5 % in the little brown bats (*Myotis lucifugus*) in New York (Trimarchi and Debbie 1977), to 58 % in Seba's short-tailed bat (*Carollia perspicillata*) in Peru (Salmón-Mulanovich et al. 2009) and 67 % in *T. brasiliensis* in Texas (Baer and Smith 1991). As with other lyssaviruses discussed below, the potential for high antibody prevalences in bat populations and infrequent reports of mortality suggest that many individuals exposed to the virus survive, contrary to the overwhelmingly lethal nature of lyssavirus infections in other mammalian species (reviewed in Banyard et al. 2011). The mechanisms for this remain unclear.

Seven bat lyssaviruses have been isolated in Eurasia (Table 10.1). **European bat lyssavirus** type 1 and type 2 (EBLV-1 and EBLV-2; Bourhy et al. 1992) are the most widely recognised and studied. Five fatal cases of human infections with EBLV have so far been reported, three from EBLV-1 (Roine et al. 1988; Selimov et al. 1989; Botvinkin et al. 2005) and two from EBLV-2 (Lumio et al. 1986; Fooks et al. 2003; Nathwani et al. 2003). Spillover of EBLV-1 into other mammals has also been observed, but rarely, with examples including zoo bats (Rønsholt et al. 1998), sheep (Tjørnehøj et al. 2006), domestic cats (Dacheux et al. 2009) and a stone marten (Müller et al. 2004). While EBLV-1 and EBLV-2 have been detected in a range of bat species (reviewed in Schatz et al. 2013), they are most frequently associated with serotine bats (*Eptesicus serotinus*) and Daubenton's bat (*Myotis daubentonii*), respectively. The dynamics of EBLV infections in their natural hosts is poorly understood, but banding and recapture data and the frequent capture of apparently healthy bats with antibodies against EBLV suggest that many bats survive infection (Serra-Cobo et al. 2002; Amengual et al. 2007; Schatz et al. 2013). In cases where bats develop clinical symptoms of EBLV infection, the affected individuals are often unable to fly, are generally weak and show abnormal behaviour, including attempts to bite (Banyard et al. 2011). Experimental studies suggest that variable development of clinical signs may be related to inoculation route and dose (reviewed in Banyard et al. 2011).

Comparatively, little is known about the remaining Eurasian bat lyssaviruses, which have each been isolated from bats only once: West Caucasian bat virus (WCBV, Botvinkin et al. 2003), Bokeloh bat lyssavirus (BBLV, Freuling et al. 2011), Aravan virus (ARAV, Kuzmin et al. 1991), Irkut virus (IRKV, Botvinkin et al. 2003) and Khujand virus (KHUV, Kuzmin et al. 2001), or is only known from partial genetic sequence data (Lleida virus, Ceballos et al. 2013, Table 10.1). Of these, only IRKV has been detected in other mammals (a human who developed rabies after a bat bite, Leonova et al. 2009). WCBV appears to have a large geographical range. It was isolated from *Miniopterus schreibersii* in Russia, but cross-reactive antibodies have also been detected in *Miniopterus* bats in Kenya (Kuzmin et al. 2008a). The relatively wide distribution and migratory behaviour of *Miniopterus* spp. may facilitate cross-continental transmission of this virus. Alternatively, given the close relationship between WCBV and Ikoma virus (IKOV), which was recently isolated in neighbouring Tanzania, the serological findings from Kenya could in fact indicate exposure to IKOV or another related lyssavirus rather than WCBV (Marston et al. 2012; Horton et al. 2014). Similarly, serological surveys have detected antibodies against ARAV virus and KHUV virus in Indian flying foxes (*Pteropus giganteus*) from Bangladesh (Kuzmin et al. 2006), and ARAV, KHUV, IRKV or Australian bat lyssavirus in Lyle's flying foxes (*P. lylei*) and dawn bats (*Eonycteris spelaea*) from Thailand (Lumlertdacha et al. 2005). Yet, given the limited lyssavirus surveillance in bats performed to date in this region and that individuals in these studies tested positive to multiple viruses, these results likely represent cross-reactivity of serological assays to unknown lyssaviruses.

Africa also hosts significant lyssavirus diversity, with five species identified, though only three of these isolated from bats to date (Table 10.1). **Duvenhage virus** (DUVV, Meredith et al. 1971) is the only phylogroup I lyssavirus in Africa and is more closely related to RABV, ABLV and the majority of the European species than other known African lyssaviruses. Since it was first isolated from a human in 1970, two more fatal human infections of DUVV have been reported, one in South Africa in 2006 (Paweska et al. 2006) and one from the Netherlands in 2007 after obtaining the infection in Kenya (van Thiel et al. 2008). DUVV has been isolated from bats twice, once from a presumed *M. schreibersii* bat in South Africa and once from an Egyptian slit-faced bat (*Nycteris thebaica*) in Zimbabwe (Schneider et al. 1985; Foggini 1988; Paweska et al. 2006). No further information is so far available on this apparently rare African lyssavirus.

In contrast, **Lagos bat virus** (LBV) is the most widely detected lyssavirus in Africa (Banyard et al. 2011). In 1956, this virus was first isolated from a straw-coloured fruit bat (*Eidolon helvum*; Boulger and Porterfield 1958). Since then, the virus has been isolated and neutralising antibodies detected in a variety of fruit bat species, one insectivorous bat species, domestic cats, domestic dogs and a water mongoose, but not in humans (reviewed in Banyard et al. 2011). *E. helvum* and *Rousettus aegyptiacus* are likely primary reservoir hosts for LBV, with seroprevalences ranging from 6 to 80 % and 29 to 46 %, respectively, depending on the region (Hayman et al. 2008, 2012; Kuzmin et al. 2008b; Dzikwi et al. 2010; Peel et al. 2013). LBV has been isolated from healthy, rabid and dead bats (reviewed in Banyard et al. 2011), but longitudinal studies in Ghana (Hayman et al. 2012) and surveys across continental Africa (Peel et al. 2010, 2013) suggest widespread exposure, no difference in survival between seropositive and seronegative *E. helvum*, and viral persistence in very small, isolated island populations. Early infection experiments with LBV suggested that LBV and other phylogroup II viruses were less pathogenic than other lyssaviruses (Boulger and Porterfield 1958; Badrane et al. 2001). However, recent experimental infections indicated the potential for comparable mortality between LBV and RABV and indicated that significant differences might instead exist between different LBV isolates (Kuzmin et al. 2010; Markotter et al. 2009).

Of the other African lyssaviruses, only Shimoni bat virus (SHIBV) has been detected in bats (Commerson's leaf-nosed bat (*Hipposideros commersoni*) in Kenya; Kuzmin et al. 2010) and only Mokola virus (MOKV) has been detected in humans (on two occasions in Nigeria, Familusi and Moore 1972; Familusi et al. 1972). MOKV has also been isolated from cats and small wild mammals, however, the natural reservoir host is unknown (Nel 2001). Ikoma virus was isolated from a rabid African civet (*Civettictis civetta*), but it is believed that the civet was a spillover host and the true reservoir host is yet to be identified (Horton et al. 2014).

The only lyssavirus detected in Australia to date—**Australian bat lyssavirus** (ABLV)—has two known lineages, one circulating in flying foxes and one in an insectivorous bat (Fraser et al. 1996; Gould et al. 2002; Warrilow 2005). In 1996, shortly after ABLV was first isolated from a black flying fox (*P. alecto*) that was

unable to fly (Fraser et al. 1996), a 39-year-old woman died of clinical rabies after being bitten by a yellow-bellied sheath-tail bat (*Saccolaimus flaviventris*; Gould et al. 2002). Two subsequent human cases have been identified, a woman who died in 1998, 27 months after being bitten by a flying fox (Hanna et al. 2000), and a child who died in 2014 after being scratched by a flying fox (Francis et al. 2014). Experimental infection of grey-headed flying foxes (*P. poliocephalus*) with ABLV resulted in clinical signs of weakness, trembling and limb paralysis in three out of ten individuals (McColl et al. 2002). As with other bat lyssaviruses, a small proportion of ABLV-positive bats succumb to encephalitis-like symptoms (Hooper et al. 1997), yet serological tests show a high prevalence of antibodies in populations of surviving bats (McColl et al. 2000).

10.3.2 Paramyxoviruses

The most notable viruses from the *Paramyxoviridae* family in bats are those of the genus *Henipavirus*, which are the subject of many reviews (e.g. Halpin and Rota 2015; Smith and Wang 2013, Luby and Gurley 2012; Clayton et al. 2013; Middleton and Weingartl 2012; Field and Kung 2011). The first recognised henipavirus, **Hendra virus** (HeV), was first detected during an outbreak of infectious respiratory disease in horses and then humans in Hendra, Australia, in 1994 (Murray et al. 1995). Ultimately, 13 of 20 infected horses died or were euthanised, and of two humans working closely with horses who became infected, one died from acute pneumonia (Murray et al. 1995; Plowright et al. 2015). This spillover was preceded a month earlier by another involving two horses and one human over 800 km away in Mackay, but which went unrecognised until 1995 (Rogers et al. 1996; O'Sullivan et al. 1997). An initial serological survey of 46 wildlife species (excluding bats) failed to identify a reservoir host; however, serological evidence of HeV infection was later identified in all four species of flying foxes native to Australia (Young et al. 1996). Virus isolation (Halpin et al. 2000) and experimental studies (Halpin et al. 2011) have confirmed pteropodid bats as reservoir hosts for henipaviruses (with a lack of clinical signs), with evidence that black (*P. alecto*) and spectacled flying foxes (*P. conspicillatus*) are the main reservoir species for HeV (Smith et al. 2014; Goldspink et al. 2015).

Because HeV is frequently detected in the urine of wild flying foxes (Smith et al. 2014), the predominant transmission route to horses is likely via material recently contaminated with bat urine (e.g. pastures) or via direct transmission (Martin et al. 2015). Recognised spillover events from bats to horses occurred sporadically from 1994 to 2004 and annually since 2006, with five spillover events resulting in ongoing transmission to humans in close contact with horses (a total of seven human cases and four deaths; Field et al. 2010). Spillover events are spatiotemporally clustered, occurring year-round in the northern tropics, but seasonally clustered in winter with a peak in July in subtropical regions (Plowright et al. 2015).

The relative importance of various hypothesised drivers of HeV dynamics in bats and subsequent spillover to horses is still unclear (Plowright et al. 2015).

Nipah virus (NiV), the second henipavirus to be recognised, was first isolated in 1999 from pigs and encephalitic pig workers in Malaysia (Center of Disease Control and Prevention 1999). NiV spillover has not been observed since this time in Malaysia; however, annual seasonal outbreaks with high case fatality (average 73 %) have occurred in people in Bangladesh since 2001 (Hsu et al. 2004; Luby et al. 2009; Luby and Gurley 2012), with occasional spillover also occurring in neighbouring India (Chadha et al. 2006; Harit et al. 2006). Due to the close relatedness of HeV and NiV, fruit bats were targeted, and serological evidence quickly identified them to be the natural reservoir of NiV (Enserink 2000; Yob et al. 2001). This was subsequently supported by isolation of NiV from the urine of *P. hypomelanus* (Chua et al. 2002a), *P. vampyrus* (Rahman et al. 2010) and *P. lylei* (Reynes et al. 2005), and seroconversion in the absence of clinical signs following experimental infections in *P. vampyrus* (Halpin et al. 2011). Antibodies against NiV and NiV-related viruses have now been detected in a variety of bat species (including non-pteropid bats) across a wide geographical area (summarised in Breed et al. 2013). NiV transmission to humans appears to occur via a wider variety of routes compared with HeV. Infection of domestic animal intermediate hosts (via consumption of saliva- or urine- contaminated partially eaten fruits or raw date palm sap) has been implicated as a source of human infections in both Malaysia and Bangladesh (Chua et al. 2002b; Chowdhury et al. 2014). In Malaysia, human infections resulted from direct contacts with infected pigs (Chua et al. 1999; Paton et al. 1999; Parashar et al. 2000), whereas in Bangladesh, transmission to humans regularly occurs via consumption of contaminated date palm sap (Luby et al. 2006; Rahman et al. 2012) or directly from human to human (e.g. via nursing sick individuals or preparation for burial; Hughes et al. 2009). The risk of direct human infection with NiV from bats is considered to be lower than horizontal transmission once the virus enters the human population (Gurley et al. 2007; Luby et al. 2009; Chong et al. 2003).

A third henipavirus, **Cedar Virus** (Marsh et al. 2012), has been isolated from urine collected under a mixed *P. alecto*/*P. scapulatus* roost in Australia. In contrast to HeV and NiV, however, it appears to be of low pathogenicity and failed to induce clinical signs in experimentally infected laboratory animal species (Marsh et al. 2012). Serological evidence from South-East Asia and Australasia (Breed et al. 2013) and the wide diversity of paramyxovirus sequences detected in Australia (Vidgen et al. 2015) suggest more henipaviruses are yet to be found. Additionally, although henipaviruses were long thought to be restricted to Asia and Australia, antibodies cross-reactive to HeV and NiV were detected in Madagascar in 2007, suggesting a potentially wider geographical distribution of henipa-related paramyxoviruses (Iehl   et al. 2007). This was supported by serological findings and molecular detection of henipa- or henipa-like viruses in mainland Africa and its offshore islands (Hayman et al. 2008, 2012; Peel et al. 2010, 2013; Drexler et al. 2012). Indeed, a recent serological study indicates that these

viruses are also occasionally transmitted to humans in Africa (Pernet et al. 2014), though no African henipavirus has been isolated to date.

Viruses from the paramyxovirus genus *Rubulavirus* (a genus which includes the human mumps virus) have also been frequently detected in bats (Barr et al. 2015). **Menangle virus** was isolated from pigs following the birth of unusually high numbers of stillborn and deformed piglets in Australia (Philbey et al. 1998). Two piggery personnel had neutralising antibodies against Menangle virus after having recovered from an unexplained febrile illness (Philbey et al. 1998). Flying fox colonies roosting in close proximity to the piggeries were a suspected source of infection for pigs, with subsequent transmission to humans (Philbey et al. 1998). This was supported by serological evidence from *P. poliocephalus*, *P. alecto* and *P. conspicillatus*, and recent virus isolation from *P. alecto* (Barr et al. 2012). Other isolated bat rubulaviruses with unknown or limited understanding of their zoonotic potential include Tioman virus from Malaysia (Chua et al. 2001), Tuhokovirus 1, 2 and 3 from China (Lau et al. 2010), Achimota virus 1 and 2 from Ghana (Baker et al. 2013c) and Hervey, Grove, Teviot and Yeppoon paramyxoviruses from Australia (Barr et al. 2015). Neutralising antibodies to Tioman virus and Achimota viruses have been detected in humans, suggesting previous exposure and infection with the virus (Yaiw et al. 2007; Baker et al. 2013c). Pigs experimentally infected with Tioman virus produced neutralising antibodies and excreted virus in saliva, but were either asymptomatic or developed only a fever (Yaiw et al. 2008). Undetected infection in pigs could therefore facilitate transmission to humans.

Finally, viral fragments related to rubulaviruses and the proposed genus *Jeilongvirus* have also been detected outside the range of fruit bats, in European insectivorous bat species (Kurth et al. 2012). However, nothing is yet known about the relevance of these viruses as potentially zoonotic threats to humans.

10.3.3 Coronaviruses

Bat coronaviruses were first identified from species of the genus *Miniopterus* (Poon et al. 2005), however, with unknown zoonotic potential. The most prominent coronavirus, the one causing **severe acute respiratory syndrome (SARS)**, was followed by a pandemic spread in humans after the first outbreak in China in 2002 (Rota et al. 2003). Soon after the outbreak, the virus was detected in masked palm civet (*Paguma larvata*) and raccoon dogs (*Nyctereutes procyonoides*) in a market in Guangdong Province, where SARS was first reported (Guan et al. 2003). A survey of common wildlife species in the area identified bats to be the natural reservoir of SARS coronavirus, with viruses from bats showing greater genetic diversity than the ones isolated from other species, including humans (Li et al. 2005). Bats can regularly be found in markets in China, which makes direct transmission of the virus from bats to humans likely (Li et al. 2005). The followed pandemic spread with 8096 confirmed cases of which 774 were fatal can be

accounted to rapid interindividual transmission of the virus once it entered the human population (World Health Organization 2003).

Outside Asia, **SARS-like coronaviruses** have been detected in the lesser horseshoe bat (*Rhinolophus hipposideros*) from Europe (Rihtarič et al. 2010), in *Chaerephon* sp. from Kenya (Tong et al. 2009) and in *Hipposideros commersoni* from Nigeria (Quan et al. 2010). Antibodies against SARS coronavirus are present in various African bat species (Müller et al. 2007). As with many newly detected viruses, their potential threat as a zoonotic disease is yet unclear.

Since the outbreak of SARS in Asia has been traced to bats as natural hosts of the virus, the same was suspected to be the case for **Middle East respiratory syndrome (MERS)**, an infection that has been occasionally spreading among humans of the Arabian peninsula since 2012 (Zaki et al. 2012). Most human infections have been traced down to close contacts with dromedary camels (*Camelus dromedarius*), which carry a virus with a similar genome organisation as human MERS (Hemida et al. 2014). There is at least one report of direct transmission of the virus from camels to humans via contact with infected animals (Memish et al. 2014). However, a small fragment of a coronavirus PCRed from an Egyptian tomb bat (*Taphozous perforatus*) showed 100 % nucleotide identity to virus from the human index case-patient of MERS, suggesting that this species may be one of the putative natural reservoirs of the virus (Memish et al. 2013). Bat-derived MERS virus has been shown to be able to use human receptors and thus could potentially infect human cells (Yang et al. 2014). However, given the generally low prevalence of MERS virus in bat populations, a direct spillover from bats to humans is unlikely, and transmission probably happens mainly via camels as intermediate hosts (Memish et al. 2013). In fact, no other bat has yet been found to carry MERS virus since the one reported by Memish and colleagues in 2013.

The intensified search for viruses in bats worldwide has led to the detection of coronaviruses other than SARS and MERS, whose potential to be or become zoonotic has yet to be investigated (Woo et al. 2006; Tang et al. 2006; Dominguez et al. 2007; Carrington et al. 2008; Brandão et al. 2008; Misra et al. 2009; Pfefferle et al. 2009; Donaldson et al. 2010; Watanabe et al. 2010; Drexler et al. 2010; Falcón et al. 2011; Annan et al. 2013; Ge et al. 2013; Anthony et al. 2013; Ithete et al. 2013). No clinical symptoms associated with infections with SARS-like and other coronaviruses have yet been described for bats.

10.3.4 Filoviruses

Ebola virus is the most prominent filovirus, causing severe haemorrhagic fever in humans with high mortality and fast spreading among African populations. The recent outbreak in 2013 in west Africa has resulted in the most severe epidemic of Ebola so far, with more than 11,000 lethal cases (as by September 2015; according to World Health Organization; <http://apps.who.int/ebola/ebola-situation-reports>).

All Ebola outbreaks recorded until 2004 in Gabon and the Republic of the Congo have been linked to handling of gorilla, chimpanzee or duiker carcasses, species that can carry the Ebola virus (Leroy et al. 2004; Pigott et al. 2014). It has thus become apparent that spillover from animals to humans occurs through hunting, butchering and consumption of bushmeat (Gonzalez et al. 2005; Li and Chen 2014; Chap. 12), followed by fast human-to-human transmission (World Health Organization 2014). An outbreak of Ebola in Congo in 2007 that resulted in 260 infected humans of whom 186 died has been traced to a potential direct transmission from a dead fruit bat that the first human victim bought from hunters to eat (Leroy et al. 2009). Antibodies against Ebola virus have since been detected in a total of 14 bat species, with seroprevalences of up to 44 % depending on species and location (Olival and Hayman 2014). Experimental infection of several bat species with Ebola led to high replication of the virus, but to no apparent signs of illness, suggesting that Ebola infections are subclinical in these species (Swanepoel et al. 1996). One *Eidolon helvum* has survived for at least 13 months after being tested seropositive for Ebola virus and Lagos bat virus, indicating long-term survival of an individual bat following exposure to these viruses (Hayman et al. 2010). The recent outbreak of Ebola in Guinea and neighbouring countries in 2013—countries that are at significant distance to the previous outbreaks in central Africa—has caused speculations about a possible transmission of the virus by migrating fruit bats (Bausch and Schwarz 2014; Vogel 2014). However, as the strain of the west African Ebola virus is a genetic outlier within the known Ebola viruses, it has been argued that the west African variant may have emerged from local wildlife populations rather than from migrating individuals (Gatherer 2014). Furthermore, although speculated (Saéz et al. 2015), it is yet not clear whether the spillover of Ebola virus in west Africa originated from bats.

Marburg virus is the only filovirus that has so far been directly isolated from bats (Towner et al. 2009; Amman et al. 2012; Pourrut et al. 2005). The first outbreak of the virus was caused by a spillover from laboratory monkeys to humans in Marburg, Germany, in 1967 (Jacob and Solcher 1968). In 2007, mine workers in a cave in Uganda were diagnosed with Marburg haemorrhagic fever that potentially resulted from a spillover of the virus from a colony of *Rousettus aegyptiacus*, where 5.1 % of tested individuals carried the virus (Towner et al. 2009). The high divergence of the genome sequence of Marburg in this population suggests a long-term association of the virus with the host, leading to the assumption that bats are the natural reservoir (Towner et al. 2009). However, given that no other bat species has yet tested positive for the virus (Towner et al. 2007), and seroprevalence being generally low in *R. aegyptiacus* (Pourrut et al. 2009), spillovers from bats to humans may be rare events.

The **Reston Ebolavirus** has first been detected in 1989 in crab-eating macaques (*Macaca fascicularis*) imported from the Philippines to be used for animal testing in laboratories in Reston, USA (Jahrling et al. 1990). During a second outbreak in 1990, animal handlers developed antibodies but did not get sick (Center for Disease Control and Prevention 1990). In 2008, Reston Ebolavirus was isolated from pigs in the Philippines (Marsh et al. 2011), and soon after, some

sampled *R. amplexicaudatus* had antibodies against the virus, while 16 other bat species tested negative against Reston Ebolavirus (Taniguchi et al. 2011). Screening for antibodies of the Ebola virus and Reston Ebolavirus in bats in Bangladesh has found seropositive *R. leschenaultii*, suggesting that these filoviruses or related strains are distributed at a much larger geographic range than previously assumed (Olival et al. 2013).

10.4 Main Conservation Issues Related to Bat Viruses

10.4.1 Direct Effect: Viruses Killing Bats

From all the viruses described above, only a few seem to affect bats. Although experimental infection with RABV leads to mortalities between 40 and 90 % depending on the bat species (Sétien et al. 1998; Jackson et al. 2008; Turmelle et al. 2010), there are no observed mass mortalities in natural populations (Pawan 1959). The only virus that may be largely lethal for bats is the **Lloviu virus**, which is closely related to Ebola and Marburg virus, but not yet of zoonotic relevance. It was detected during investigations of a massive die-off of *Miniopterus schreibersii* in a cave in Spain (Negredo et al. 2011). However, a causal connection between the detected virus and death of the bats has not yet been confirmed, and other bat species roosting in the same caves appeared to remain unaffected (Roué and Nemoz 2004).

The lack of reports of viruses that are detrimental for bat health should not imply that viruses in general are not of importance for the conservation of bat populations. Similar to white-nose syndrome causing mass mortalities in North American bats (Frick et al. 2010), newly emerging viruses may put local populations at threat. This may be especially the case if pathogens cross geographical borders and infect naïve bat populations. *Pseudogymnoascus destructans*—the causative fungus responsible for white-nose syndrome—likely originated from Europe, where it seemingly causes no bat fatalities, in contrast to North America (Puechmaille et al. 2010; Frick et al. 2010; Frick et al. 2015, Chap. 9).

10.4.2 Indirect Effects: Biased Public Perception

Generally, the public perception of bats as aesthetically less appealing mammals as well as folklores that often associate bats with negative stigma makes bat-related conservation efforts time-consuming and demanding (Fenton 1997; Allen 2004; Knight 2008). The recent outbreaks of viral zoonotic diseases with the identification of bats as putative natural hosts have further complicated bat conservation efforts (Li et al. 2005; Knight 2008). Following numerous and often lurid reports of fatal zoonotic diseases by the media, public perception of bats is mostly

skewed by fear and lack of information (Kingston 2016, Chap. 18). Therefore, it is important to highlight the context of bat-associated infections in order to provide more evidence-based information about the emergence and transmission of bat-related zoonotic diseases, which may lead to a more balanced reputation of bats. Depending on educational, cultural, legal and medial background of the targeted audience, specific aspects need to be taken into account.

In Europe and North America, rabies is, so far, the only viral disease that is associated with bats. The fact that lyssaviruses are occasionally found in temperate zone bats sometimes finds its way to the media, not always in favour of bats. Biased newspaper articles or press campaigns may result in the public misconception that bats are aggressive animals or that their mere presence can lead to human infections with these viruses. Although there are anecdotal reports of unprovoked attacks of bats on humans and dogs (Baer and Smith 1991), bats, as is the case of most mammals, usually only bite when handled or provoked. Furthermore, once bitten or scratched by a bat, immediate post-exposure vaccination can prevent a person from contracting rabies (see Sect. 10.5.2). In the case of the 37-year-old woman who died from a bat lyssavirus infection in Kenya, staff members of the health facility which the woman visited after being scratched by a bat were unaware of the possibility of rabies transmission (van Thiel et al. 2009). Likewise, two persons in Europe who worked regularly with bats and died from rabies after being bitten and scratched by bats received neither pre- nor post-exposure treatment (Roine et al. 1988; Nathwani et al. 2003). These two cases triggered a Europe-wide serological screening effort involving more than 11,000 bats, with seroprevalences varying depending on the species and location (Racey et al. 2012). EBLV-1 was most commonly detected in the serotine bat (*Eptesicus serotinus*), while EBLV-2 was very uncommon in all bat species. As a result, the public has been persuaded not to handle bats or to do so only with gloves and, in the case of bat workers, to receive pre- and/or post-exposure immunisation. Two fatal cases in which persons contracted rabies in Australia (Samaratunga et al. 1998; Hanna et al. 2000) triggered a similar campaign on this continent (Speare et al. 1997, but see Francis et al. 2014). Efficient education of medical professionals worldwide seems to be pivotal for implementing the correct treatment after scratches or bites from bats. In addition, vaccination should be mandatory for those who are frequently exposed to bats (Rupprecht and Gibbons 2004). Studies on animal models have shown that rabies vaccine also provides protection against other, although not all, lyssaviruses' variants (Brookes et al. 2005; Hanlon et al. 2005). However, there is no known case of a person developing bat-associated rabies despite having been vaccinated, neither pre- nor post-exposure. Thus, getting infected by some sort of bat-related virus is unlikely in Europe and North America and decreases virtually to zero if people who experienced bat bites and scratches are treated appropriately.

There is no case known for paramyxoviruses having spilled over to humans by direct contact with bats. An extensive serological survey among people frequently handling bats in Australia revealed no antibodies against Hendra virus (Arklay et al. 1996). The virus apparently needs horses as amplifier hosts, from where the virus can further be transmitted to persons in close contact with infected

individuals. Nevertheless, the outbreak of Hendra increased the unpopularity of flying foxes in Australia, making conservation of the four native species challenging (Thiriet 2011). Unlike Hendra, Nipah virus has likely been acquired by humans via consumption of contaminated date palm sap (Luby et al. 2006; Rahman et al. 2012), followed by person-to-person transmission (Gurley et al. 2007). Although diseases associated with Hendra virus and Nipah virus have high mortality rates, the risk of infection for humans seems to be low (Chong et al. 2003), and countermeasures may be taken in order to prevent future spillover events (see Sect. 10.5.2). MERS, just as Hendra virus, apparently needs livestock as an amplifier host. In contrast to dromedaries (Hemida et al. 2014), seroprevalence of MERS seems to be low in bats (Memish et al. 2013), making direct transmission from bats to humans unlikely. As long as details on MERS infections in dromedaries and how to mitigate them are missing, it is hard to give recommendations to people who might be at risk.

In contrast to MERS, the spillover of SARS into the human populations most likely happened via the wildlife market, either directly from a bat, or from other wildlife species. Likewise, the hunting, butchering and consumption of chimpanzees, gorillas and bats seem to have been sources of Ebola spillovers from wildlife to humans. The education of local communities needs to carefully balance information about the potential risk of acquiring infectious diseases by consuming bushmeat, without implying that bats need to be eradicated in order to prevent spillovers. The recent outbreak of Ebola resulting in several thousand human victims, and with bats frequently being reported as the likely source of origin, has undoubtedly led to severe loss of reputation of bats on this continent, which makes the conservation of threatened populations and species even more challenging, not only in Africa, but also worldwide.

10.4.3 Indirect Effect—Culling

The direct persecution of bats often seems to be the most effective way to deal with bat-borne diseases to members of the public. Killing of bats has long been acceptable, even if they are protected (Chap. 14). Even though culling may be officially banned and thus not supported by authorities or governmental programs, large-scale killing of bats or the destruction of roost trees may still be commonly practiced in areas where zoonotic diseases are spreading.

In Australia, for example, flying foxes are frequently harassed and killed, both legally (under permits issued by state wildlife management agencies) and illegally. This happened most prominently during periods when Hendra virus emerged in Australian flying fox populations (Roberts et al. 2012). Half of the flying fox species native to Australia have declined about 30 % in population size during the last decade, and killing of bats usually does not lead to legal measures (Booth 2005). Furthermore, large-scale culling leads to a change of movement behaviour of bats, with new, susceptible individuals being recruited from nearby colonies

(Field 2009). Instead of reducing the viral prevalence, this may therefore lead to the exact opposite (see below).

In the attempt to reduce rabies incidences, vampire bats are regularly culled in many parts of Latin America (Streicker et al. 2012). In Brazil, for example, governmental programs are in action that involve targeted campaigns against vampire bats. During these measures, vampire bats are captured and poisoned or coated with anticoagulant and released, so that allogrooming kills their conspecifics (Medellin 2003). Furthermore, bat roosts are destroyed using fire and explosives (Mayen 2003), which also leads to dramatic declines of non-target bats (Furey and Racey 2015, Chap. 15). Besides the questionable methods involved, instead of reducing viral abundance in the population, culling of wildlife can lead to an increase in viral spreading. New hosts are recruited and the dispersal probability of infected individuals increases, which results in transmission of the disease to naïve hosts (Donnelly et al. 2005; Choisy and Rohani 2006; Streicker et al. 2012). This was the case for vampire bats in Peru, where culling failed to reduce seroprevalence of rabies in bat populations, but rather had the opposite effect (Streicker et al. 2012). Therefore, persecution of bats as potential carriers of zoonotic diseases has been denounced as useless and even counterproductive by both conservationists and experts on disease transmission (Hutson and Mickleburgh 2001; Knight 2008).

10.4.4 Indirect Effect—Killing of Bats for Virus Surveys

In the scope of recently emerging zoonotic diseases, the search for new bat-borne viruses has become a well-funded field in the scientific community. While research is important to advance our understanding about the emergence of diseases and to possibly prevent further spillover events, the methods involved in these surveys are sometimes questionable from the perspective of bat conservation (Racey 2015). Some of the investigated bat species are listed as near threatened or vulnerable by the International Union for the Conservation of Nature (IUCN), with decreasing population sizes even in many species of least concern. While most surveillance studies that involve species of conservation concern use non-lethal methods such as antibody screening in blood (Hayman et al. 2008; Young et al. 1996; Lumlertdacha et al. 2005; Wacharapluesadee et al. 2005; Reynes et al. 2005), others have involved the killing of a considerable number of bats of various conservation status (e.g. in Yob et al. 2001; Kuzmin et al. 2008b, 2010, 2011b; Dzikwi et al. 2010 and Sasaki et al. 2012). In order to limit such detrimental surveys, the Food and Agriculture Organization of the United Nations (2011) has published a guideline for investigating the role of bats in emerging zoonotic diseases, including non-invasive protocols, which not only reduce the impact on bat populations, but also minimise the transmission risk of viral diseases. Such protocols have now been widely adopted, as for example by Ecohealth Alliance and other international research groups and networks.

10.5 Counter Measures in Favour of Bat Conservation

10.5.1 *Preventing the Emergence of New Viral Diseases*

In general, preventing the emergence of infectious diseases in wildlife populations is extremely challenging and usually underfunded, with only few practical suggestions being discussed (Daszak et al. 2000). For example, it is important that translocations of animals across geographical borders need to follow strict guidelines in order to prevent the introduction of exotic pathogens in novel areas (e.g. Woodroffe 1999). Furthermore, an integration of knowledge about disease dynamics, as well as ecological and immunological aspects of the host, may contribute to a better understanding of emerging infectious diseases in wildlife species such as bats (Daszak et al. 2000).

10.5.2 *Educational Efforts*

As many bat-borne viral diseases have high lethality rates for humans, preventing spillover events are of central importance. In particular, spillover by direct contact to bats, such as via bites or bat consumption, may bear severe risks to humans that could be minimised by educational programs (Kingston 2016, Chap. 18). Reducing the risk of outbreaks of zoonotic viruses may also lead to more positive attitudes towards bats, which may further be increased by highlighting their ecological importance as pollinators, seed dispersers and pest control for agriculture (Ghanem and Voigt 2012). Moreover, conservation measures that promote the preservation of bat habitats serve a dual role as they can decrease the contact zone between bats and humans, thus reducing the risk of spillover.

As aforementioned vaccination against rabies and other lyssaviruses should be mandatory for persons working with bats and recommended for other people at risk. A significant problem is that both pre- and post-exposure treatments are expensive and thus may not be readily available in developing countries, such as in Central and South America. Here, building houses in a bat-proof manner in order to avoid vampire bites during sleep and decreasing the risk of direct contact with other bats has so far been the best solution (Greenhall 1964; Voigt et al. 2016, Chap. 14).

A different issue is the transmission of Nipah viruses via consuming raw date palm sap contaminated by urine, faeces or saliva of bats (Luby et al. 2006; Rahman et al. 2012). Here, cooking the sap at temperatures above the level that viruses tolerate is an effective measure to prevent spillover (Hughes et al. 2009). Additionally, preventing bats from accessing date palms and thus contaminating the sap has been proved to be both efficient and relatively cheap (Nahar et al. 2010, 2013). The traditional “bamboo skirt” method for example uses inexpensive, recyclable bamboo to cover the part of the date palm where the sap is collected, preventing bats and



Fig. 10.1 Intact trees with colonies of *Eidolon helvum* (left) in Yaoundé, Cameroon, as compared to former roosting trees that have been cut (right) after bats were suspected to be the source of the recent Ebola outbreak in western Africa (photograph credits: Simon Ghanem)

other vertebrates from getting access. Furthermore, in contrast to bird nets, this measure is non-lethal to the bats and therefore of high conservation value to local populations. However, such protective measures are reported to be rarely used in Bangladesh (Nahar et al. 2010, 2013). This could potentially be changed by encouraging local farmers to use this method, emphasising its inexpensiveness and efficiency while highlighting the reduced risk of acquiring Nipah virus disease.

One of the key issues both for conservation and public health is the direct transmission of SARS and Ebola via wildlife markets. In South-East Asia, flying foxes are hunted regularly for the purpose of food (Mickleburgh et al. 2002; Mildenstein et al. 2016, Chap. 12), sometimes even authorised by the local Wildlife Department such as in Malaysia (Breed et al. 2006). Likewise, fruit bats are consumed regularly throughout Africa (Mickleburgh et al. 2009; Mildenstein et al. 2016, Chap. 12). Since bats are suggested as potential reservoir for the recent outbreak of Ebola, Guinea banned bats for sale from markets (Gatherer 2014). Educational efforts to reduce the threat both to public health by zoonotic diseases and to the conservation of local bat populations are challenging, as they are usually impeded by the lack of understanding of entrenched cultural behaviours and social components (Pooley et al. 2015; Kingston 2016, Chap. 18). In Ghana, for example, where the consumption of bats is part of the local culture and traditions, a survey revealed that knowledge about the ecological and economical value of bats would not make people refrain from killing and eating bats (Kamins et al.

2014). Usually, the direct economic benefit from selling hunted bats is more valuable to an individual person than the indirect, not always obvious economic value of bats, for example, for agriculture. However, about half of the hunters stated they would stop hunting bats if they could make them sick (Kamins et al. 2014). This highlights the potential effectiveness of public education, but careful consideration is needed to avoid demonising bats in the process (Pooley et al. 2015). The recent Ebola epidemic in western Africa for example has led to an increase in the persecution of bats, with roosts being destroyed and colonies being killed by communities (Fig. 10.1). Although preventing bats from being consumed may have higher priorities due to public health reasons, the culling of whole colonies as a likely result may be much more of a threat for the conservation of bats than the bushmeat trade (Pooley et al. 2015).

10.5.3 Environmental Conservation

Combining knowledge about the ecology of the host species as well as the disease dynamics of the virus may be crucial for establishing efficient disease prevention programs (e.g. Plowright et al. 2015). Here, it needs to be noted that the emergence of zoonotic diseases from bats also seems to be a consequence of anthropogenic alteration of natural environments (e.g. Daszak et al. 2001). For example, in Central and South America, the conversion of forested habitats into pastures shifted the dominant food source for vampire bats from native vertebrates to livestock. This has increased rabies transmission from vampire bats to livestock and domestic animals in many parts of Latin America (Schneider et al. 2009). Where bat habitats have been converted largely into agricultural farmland, the remaining bat populations are forced to concentrate in patches that provide them with resources they need. Flying foxes, for example, are highly sensitive to landscape modifications, as they require large forested areas for foraging. Where natural habitats are scarce, flying foxes may use fruiting or flowering trees in agricultural, suburban and urban areas, which increases the contact zone and spillover risk between bats and livestock or humans (Daszak et al. 2006; Plowright et al. 2015). Indeed, contact between bats and naïve hosts as a consequence of human landscape modification and encroachment likely sparked the transmission of Hendra viruses to horses (Epstein et al. 2006) and Nipah virus to pigs (Chua et al. 1999; Field et al. 2001).

10.5.4 Conservation of Bat Populations and Population Dynamics

Removing individuals or colonies from regional populations, either by unsustainable hunting or culling, can cause an increase in relative local resource availability, creating regional gradients along which bats from other populations may move,

which may lead to an increase of virus movement (Field 2009). In Australia, for example, roosts that became empty after culling, disturbing or relocating colonies of flying foxes are usually reoccupied by immigrating individuals (Roberts et al. 2012).

Anthropogenic transformation of bat habitats in Australia has also been shown to lead to decreased migration in *Pteropus* bats, which can itself lead to a decline in population immunity (Plowright et al. 2011). This could give rise to more viral shedding after local viral reintroduction, a mechanism that may be facilitated by urban habituation of fruit bat and the resulting increased contact with human and domestic animal populations (Epstein et al. 2006; Plowright et al. 2011). In Australia, all recently emerged bat-associated viruses—Hendra, Menangle and Australian bat lyssavirus—are hypothesised to be associated with habitat loss due to deforestation and agricultural intensification (Jones et al. 2013). Therefore, protection of remaining natural habitats of bats along with farm management aiming at decreasing the contact zone between bats and livestock as well as education plans increasing awareness of environmental issues and safety may play a crucial role in the avoidance of future spillovers of bat-borne diseases to livestock and human populations, and promote further protection of local bat populations.

10.6 Conclusion

Bats harbour viruses that may become zoonotic. Circumstances facilitating spillover include direct contact with bats (bites, scratches, consumption of bats), contact with material contaminated by bat saliva, faeces or urine and amplification via intermediate hosts such as domestic animals or other wildlife species. Conservational actions are not only important to prevent spillovers, but also because emerging zoonotic viruses often lead to persecution of bats. In order to reduce the transmission risk of viruses from bats to human and livestock and to protect bat species at threat, educational efforts are needed. However, entrenched cultural and social components often act as barriers to efficient changes on how people think about and respond to bats. Whenever possible, educational efforts should be done in an informative, non-lurid way, presenting the facts rather than provoking additional fears to the already bad reputation of bats. Wherever possible, solutions should be found to enable the existence of bats in anthropogenic landscape, including the development of more affordable and readily available vaccinations (e.g. against rabies), and the reduction of potential contact between bats and humans and livestock. This however also includes that the natural habitats of bats need to be better protected to provide bat populations with sufficient space and to prevent range expansion into urban and suburban areas, where contact with humans and livestock may increase the risk of spillover events. Bat-borne viruses should be considered during bat conservation efforts, and it should be equally noticed that appropriate conservation measures may even reduce the risk of viral spillover from bat populations into human populations.

Acknowledgments We thank Alison Peel, Paul Racey and an anonymous reviewer for constructive comments that helped to improve the manuscript.

Open Access This chapter is distributed under the terms of the Creative Commons Attribution Noncommercial License, which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Allen GM (2004) Bats: biology, behavior, and folklore. Dover Publications
- Allen LC, Turmelle AS, Mendonça MT, Navara KJ, Kunz TH, McCracken GF (2009) Roosting ecology and variation in adaptive and innate immune system function in the Brazilian free-tailed bat (*Tadarida brasiliensis*). *J Comp Physiol B* 179(3):315–323
- Almeida MFD, Martorelli LFA, Sodré MM et al (2011) Rabies diagnosis and serology in bats from the State of São Paulo, Brazil. *Rev Soc Bras Med Trop* 44(2):140–145
- Amengual B, Bourhy H, López-Roig M, Serra-Cobo J (2007) Temporal dynamics of European bat Lyssavirus type 1 and survival of *Myotis myotis* bats in natural colonies. *PLoS ONE* 2(6):e566
- Amman BR, Carroll SA, Reed ZD et al (2012) Seasonal pulses of Marburg virus circulation in juvenile *Rousettus aegyptiacus* bats coincide with periods of increased risk of human infection. *PLoS Pathog* 8:e1002877
- Annan A, Baldwin HJ, Corman VM et al (2013) Human betacoronavirus 2c EMC/2012-related viruses in bats, Ghana and Europe. *Emerg Infect Dis* 19(3):456
- Anthony S, Ojeda-Flores R, Rico-Chávez O et al (2013) Coronaviruses in bats from Mexico. *J Gen Virol* 94(Pt 5):1028–1038
- Arklay A, Selvey L, Taylor R, Gerrard J (1996) Screening of bat carers for antibodies to equine morbillivirus. *Commun Dis Intell* 20:477
- Badrane H, Bahloul C, Perrin P, Tordo N (2001) Evidence of two Lyssavirus phylogroups with distinct pathogenicity and immunogenicity. *J Virol* 75(7):3268–3276
- Baer GM, Smith J (1991) Rabies in nonhematophagous bats. *Nat Hist Rabies* 2:105–120
- Baker KS, Leggett RM, Bexfield NH et al (2013a) Metagenomic study of the viruses of African straw-coloured fruit bats: detection of a chiropteran poxvirus and isolation of a novel adenovirus. *Virology* 441(2):95–106
- Baker ML, Schountz T, Wang LF (2013b) Antiviral immune responses of bats: a review. *Zoonoses Public Health* 60(1):104–116
- Baker KS, Todd S, Marsh GA et al (2013c) Novel, potentially zoonotic paramyxoviruses from the African straw-colored fruit bat *Eidolon helvum*. *J Virol* 87(3):1348–1358
- Banyard AC, Hayman D, Johnson N, McElhinney L, Fooks AR (2011) Bats and lyssaviruses. *Adv Virus Res* 79:239–289
- Banyard AC, Evans JS, Luo TR, Fooks AR (2014) Lyssaviruses and bats: emergence and zoonotic threat. *Viruses* 6(8):2974–2990
- Barr JA, Smith C, Marsh GA, Field H, Wang LF (2012) Evidence of bat origin for Menangle virus, a zoonotic paramyxovirus first isolated from diseased. *J Gen Virol* 93:2590–2594
- Barr J, Smith C, Smith I et al (2015) Isolation of multiple novel paramyxoviruses from pteropid bat urine. *J Gen Virol* 96:24–29
- Barrett JL (2004) Australian bat lyssavirus. Ph.D. thesis, University of Queensland, Brisbane
- Bausch DG, Schwarz L (2014) Outbreak of Ebola virus disease in Guinea: where ecology meets economy. *PLoS Neglected Trop Dis* 8(7):e3056
- Belotto A, Leanes L, Schneider M, Tamayo H, Correa E (2005) Overview of rabies in the Americas. *Virus Res* 111(1):5–12

- Booth C (2005) Time to stop the killing. The Australasian Bat Society Newsletter 24
- Botvinkin AD, Poleschuk EM, Kuzmin IV et al (2003) Novel lyssaviruses isolated from bats in Russia. *Emerg Infect Dis* 9(12):1623
- Botvinkin A, Selnikova O, Antonova L, Moiseeva A, Nesterenko EY, Gromashevsky L (2005) Human rabies case caused from a bat bite in Ukraine. *Rabies Bull Eur* 29(3):5–7
- Boulger L, Porterfield J (1958) Isolation of a virus from Nigerian fruit bats. *Trans R Soc Trop Med Hyg* 52(5):421–424
- Bourhy H, Kissi B, Lafon M, Sacramento D, Tordo N (1992) Antigenic and molecular characterization of bat rabies virus in Europe. *J Clin Microbiol* 30(9):2419–2426
- Brandão PE, Scheffer K, Villarreal LY et al (2008) A coronavirus detected in the vampire bat *Desmodus rotundus*. *Braz J Infect Dis* 12(6):466–468
- Breed AC, Field HE, Epstein JH, Daszak P (2006) Emerging henipaviruses and flying foxes—conservation and management perspectives. *Biol Conserv* 131(2):211–220
- Breed AC, Meers J, Sendow I et al (2013) The distribution of Henipaviruses in Southeast Asia and Australasia: Is Wallace's line a barrier to Nipah Virus? *PLoS ONE* 8(4):e61316
- Brook CE, Dobson AP (2015) Bats as 'special' reservoirs for emerging zoonotic pathogens. *Trends Microbiol* 23:172–180
- Brookes S, Healy D, Fooks A (2005) Ability of rabies vaccine strains to elicit cross-neutralising antibodies. *Dev Biol (Basel)* 125:185–193
- Calisher CH, Childs JE, Field HE, Holmes KV, Schountz T (2006) Bats: important reservoir hosts of emerging viruses. *Clin Microbiol Rev* 19(3):531–545
- Carini A (1911) About one large epizootic of rabies. *Ann Inst Pasteur* 25:843–846
- Carrington CV, Foster JE, Zhu HC et al (2008) Detection and phylogenetic analysis of group 1 coronaviruses in South American bats. *Emerg Infect Dis* 14 (12):1890
- Ceballos NA, Morón SV, Berciano JM et al (2013) Novel lyssavirus in bat, Spain. *Emerg Infect Dis* 19(5):793
- Center for Disease Control and Prevention (1990) Epidemiologic notes and reports updates: filovirus infection in animal handlers. *Morb Mortal Wkly Rep* 39:221
- Center for Disease Control and Prevention (1999) Outbreak of Hendra-like virus—Malaysia and Singapore, 1998–1999. *Morb Mortal Wkly Rep* 48:265
- Chadha MS, Comer JA, Lowe L et al (2006) Nipah virus-associated encephalitis outbreak, Siliguri, India. *Emerg Infect Dis* 12(2):235–240
- Chen L, Liu B, Yang J, Jin Q (2014) DBatVir: the database of bat-associated viruses. Database:bau021. doi:[10.1093/database/bau021](https://doi.org/10.1093/database/bau021)
- Choisy M, Rohani P (2006) Harvesting can increase severity of wildlife disease epidemics. *Proc R Soc B: Biol Sci* 273(1597):2025–2034
- Chong HT, Tan C, Goh K, Lam S, Chua K (2003) The risk of human Nipah virus infection directly from bats (*Pteropus hypomelanus*) is low. *Neurol J Southeast Asia* 8:31–34
- Chowdhury S, Khan SU, Crameri G et al (2014) Serological evidence of Henipavirus exposure in cattle, goats and pigs in Bangladesh. *PLoS Neglected Trop Dis* 8:e3302
- Chua KB, Goh KJ, Wong KT et al (1999) Fatal encephalitis due to Nipah virus among pig-farmers in Malaysia. *Lancet* 354(9186):1257–1259
- Chua K, Wang L, Lam SK et al (2001) Tioman virus, a novel paramyxovirus isolated from fruit bats in Malaysia. *Virology* 283 (2):215–229
- Chua KB, Lek Koh C, Hooi PS et al (2002a) Isolation of Nipah virus from Malaysian Island flying-foxes. *Microbes Infect* 4(2):145–151
- Chua KB, Chua BH, Wang CW (2002b) Anthropogenic deforestation, El Nino and the emergence of Nipah virus in Malaysia. *Malays J Pathol* 24(1):15–21
- Clayton BA, Wang LF, Marsh GA (2013) Henipaviruses: an updated review focusing on the pteropid reservoir and features of transmission. *Zoonoses Public Health* 60(1):69–83
- Constantine DG, Blehert DS (2009) Bat rabies and other lyssavirus infections. *US Geological Survey*
- Dacheux L, Larrous F, Mailles A et al (2009) European bat lyssavirus transmission among cats, Europe. *Emerg Infect Dis* 15(2):280

- Daszak P, Cunningham AA, Hyatt AD (2000) Emerging infectious diseases of wildlife—threats to biodiversity and human health. *Science* 287(5452):443–449
- Daszak P, Cunningham AA, Hyatt AD (2001) Anthropogenic environmental change and the emergence of infectious diseases in wildlife. *Acta Trop* 78:103–116
- Daszak P, Plowright R, Epstein J et al (2006) The emergence of Nipah and Hendra virus: pathogen dynamics across a wildlife-livestock-human continuum. *Dis Ecol: Community Struct Pathog Dyn*: 186–201
- Dominguez SR, O’Shea TJ, Oko LM, Holmes KV (2007) Detection of group 1 coronaviruses in bats in North America. *Emerg Infect Dis* 13(9):1295
- Donaldson EF, Haskew AN, Gates JE, Huynh J, Moore CJ, Frieman MB (2010) Metagenomic analysis of the viromes of three North American bat species: viral diversity among different bat species that share a common habitat. *J Virol* 84(24):13004–13018
- Donnelly CA, Woodroffe R, Cox D et al (2005) Positive and negative effects of widespread badger culling on tuberculosis in cattle. *Nature* 439(7078):843–846
- Drexler JF, Gloza-Rausch F, Glende J et al (2010) Genomic characterization of severe acute respiratory syndrome-related coronavirus in European bats and classification of coronaviruses based on partial RNA-dependent RNA polymerase gene sequences. *J Virol* 84(21):11336–11349
- Drexler JF, Corman VM, Müller MA et al (2012) Bats host major mammalian paramyxoviruses. *Nat Commun* 3:796
- Dzikwi AA, Kuzmin II, Umoh JU, Kwaga JK, Ahmad AA, Rupprecht CE (2010) Evidence of Lagos bat virus circulation among Nigerian fruit bats. *J Wildl Dis* 46(1):267–271
- Enserink M (2000) Malaysian researchers trace Nipah virus outbreak to bats. *Science* 289(5479):518–519
- Epstein JH, Field HE, Luby S, Pulliam JR, Daszak P (2006) Nipah virus: impact, origins, and causes of emergence. *Curr Infect Dis Rep* 8(1):59–65
- Epstein JH, Olival KJ, Pulliam JRC et al (2009) *Pteropus vampyrus*, a hunted migratory species with a multinational home-range and a need for regional management. *J Appl Ecol* 46:991–1002
- Falcón A, Vázquez-Morón S, Casas I et al (2011) Detection of alpha and betacoronaviruses in multiple Iberian bat species. *Arch Virol* 156(10):1883–1890
- Familusi JB, Moore DL (1972) Isolation of a rabies related virus from the cerebrospinal fluid of a child with ‘aseptic meningitis’. *Afr J Med Sci* 3(1):93–96
- Familusi JB, Osunkoya BO, Moore DL, Kemp GE, Fabiyi A (1972) A fatal human infection with Mokola virus. *Am J Trop Med Hyg* 21(6):959–963
- Fenton MB (1997) Science and the conservation of bats. *J Mammal*: 1–14
- Field H, Young P, Yob JM, Mills J, Hall L, Mackenzie J (2001) The natural history of Hendra and Nipah viruses. *Microbes Infect* 3(4):307–314
- Field HE (2009) Bats and emerging zoonoses: Henipaviruses and SARS. *Zoonoses Public Health* 56:278–284
- Field H, Schaaf K, Kung N et al (2010) Hendra virus outbreak with novel clinical features, Australia. *Emerg Infect Dis* 16(2):338
- Field H, Kung N (2011) Henipaviruses—unanswered questions of lethal zoonoses. *Curr Opin Virol* 1(6):658–661
- Foggini C (1988) Rabies and rabies-related viruses in Zimbabwe: historical, virological and ecological aspects. University of Zimbabwe
- Food and Agricultural Organisation of the United Nations (2011) Investigating the role of bats in emerging zoonoses: balancing ecology, conservation and public health interests. In: Newman SH, Field HE, de Jong CE, Epstein JH (eds) FAO animal production and health manual no 12, Rome
- Fooks AR, McElhinney LM, Pounder DJ et al (2003) Case report: isolation of a European bat lyssavirus type 2a from a fatal human case of rabies encephalitis. *J Med Virol* 71(2):281–289

- Francis JR, Nourse C, Vaska VL et al (2014) Australian bat lyssavirus in a child: the first reported case. *Pediatrics* 133(4):e1063–e1067
- Fraser GC, Hooper PT, Lunt RA et al (1996) Encephalitis caused by a Lyssavirus in fruit bats in Australia. *Emerg Infect Dis* 2(4):327
- Freuling CM, Beer M, Conraths FJ et al (2011) Novel lyssavirus in Natterer's bat, Germany. *Emerg Infect Dis* 17(8):1519–1522
- Frick WF, Pollock JF, Hicks AC et al (2010) An emerging disease causes regional population collapse of a common North American bat species. *Science* 329:679–682
- Frick WF, Puechmaille SJ, Willis CKR (2016) White-nose syndrome in bats. In: Voigt CC, Kingston T (eds) *Bats in the Anthropocene: conservation of bats in a changing world*. Springer International AG, Cham, pp. 245–259
- Furey N, Racey P (2016) Conservation ecology of cave bats. In: Voigt, CC, Kingston, T (eds) *Bats in the Anthropocene: conservation of bats in a changing world*. Springer International AG, Cham, pp. 463–492
- Gatherer D (2014) The 2014 Ebola virus disease outbreak in West Africa. *J Gen Virol*: vir 0.067199-067190
- Ge X-Y, Li J-L, Yang X-L et al (2013) Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. *Nature* 503:535–538
- George DB, Webb CT, Farnsworth ML et al (2011) Host and viral ecology determine bat rabies seasonality and maintenance. *Proc Natl Acad Sci* 108(25):10208–10213
- Ghanem SJ, Voigt CC (2012) Increasing awareness of ecosystem services provided by bats. *Adv Study Behav* 44:279–302
- Gonzalez J-P, Herbreteau V, Morvan J, Leroy EM (2005) Ebola virus circulation in Africa: a balance between clinical expression and epidemiological silence. *Bull de la Société de Pathol Exotique* 98(3):210–217
- Gould AR, Kattenbelt JA, Gumley SG, Lunt RA (2002) Characterisation of an Australian bat lyssavirus variant isolated from an insectivorous bat. *Virus Res* 89(1):1–28
- Goldspink LK, Edson D, Vidgen M, Bingham J, Field H, Smith C (2015) Natural Hendra virus infection in flying-foxes, tissue tropism and risk factors. *PLoS ONE* 10(6):e0128835
- Greenhall AM (1964) Bats: their public health importance and control with special reference to Trinidad. In: *Proceedings of the 2nd vertebrate pest control conference*, p. 18
- Guan Y, Zheng B, He Y et al (2003) Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. *Science* 302(5643):276–278
- Gurley ES, Montgomery JM, Hossain MJ et al (2007) Person-to-person transmission of Nipah virus in a Bangladeshi community. *Emerg Infect Dis* 13(7):1031
- Halpin K, Young P, Field H, Mackenzie J (2000) Isolation of Hendra virus from pteropid bats: a natural reservoir of Hendra virus. *J Gen Virol* 81(8):1927–1932
- Halpin K, Hyatt AD, Plowright RK et al (2007) Emerging viruses: coming in on a wrinkled wing and a prayer. *Clin Infect Dis* 44:711–717
- Halpin K, Hyatt AD, Fogarty R et al (2011) Pteropid bats are confirmed as the reservoir hosts of henipaviruses: a comprehensive experimental study of virus transmission. *Am J Trop Med Hyg* 85(5):946–951
- Halpin K, Rota P (2015) A review of Hendra virus and Nipah virus infections in man and other animals. In: *Zoonoses-infections affecting humans and animals*. Springer Netherlands, pp. 997–1012
- Hanlon CA, Kuzmin IV, Blanton JD, Weldon WC, Manangan JS, Rupprecht CE (2005) Efficacy of rabies biologics against new lyssaviruses from Eurasia. *Virus Res* 111(1):44–54
- Hanna JN, Carney IK, Smith GA et al (2000) Australian bat lyssavirus infection: a second human case, with a long incubation period. *Med J Aust* 172(12):597–599
- Harit AK, Ichhpujani RL, Gupta S et al (2006) Nipah/Hendra virus outbreak in Siliguri, West Bengal, India in 2001. *Indian J Med Res* 123(4):553–560
- Hayman DT, Fooks AR, Horton D et al (2008) Antibodies against Lagos bat virus in megachiroptera from West Africa. *Emerg Infect Dis* 14(6):926
- Hayman DT, Emmerich P, Yu M et al (2010) Long-term survival of an urban fruit bat seropositive for Ebola and Lagos bat viruses. *PLoS ONE* 5(8):e11978

- Hayman DT, Fooks AR, Rowcliffe JM et al (2012) Endemic Lagos bat virus infection in *Eidolon helvum*. *Epidemiol Infect* 140(12):2163–2171
- Hemida MG, Chu D, Poon L, Perera R, Alhammadi M, Ng H (2014) MERS coronavirus in dromedary camel herd, Saudi Arabia. *Emerg Infect Dis* 20(7):1231–1234
- Hooper P, Lunt R, Gould A et al (1997) A new lyssavirus—the first endemic-related virus recognized in Australia. *Bull de l'Institut Pasteur* 95(4):209–218
- Horton DL, Banyard AC, Marston DA et al (2014) Antigenic and genetic characterization of a divergent African virus, Ikoma lyssavirus. *J Gen Virol* 95:1025–1032
- Hsu VP, Hossain MJ, Parashar UD et al (2004) Nipah virus encephalitis reemergence, Bangladesh. *Emerg Infect Dis* 10(12):2082–2087
- Hughes JM, Wilson ME, Luby SP, Gurley ES, Hossain MJ (2009) Transmission of human infection with Nipah virus. *Clin Infect Dis* 49(11):1743–1748
- Hutson AM, Mickleburgh SP (2001) Microchiropteran bats: global status survey and conservation action plan, vol 56. IUCN
- Iehlé C, Razafitrimo G, Razainirina J (2007) Henipavirus and Tioman virus antibodies in pteropodid bats, Madagascar. *Emerg Infect Dis* 13:159
- Ithete NL, Stoffberg S, Corman VM et al (2013) Close relative of human Middle East respiratory syndrome coronavirus in bat, South Africa. *Emerg Infect Dis* 19(10):1697
- Jackson FR, Turmelle AS, Farino DM, Franka R, McCracken GF, Rupprecht CE (2008) Experimental rabies virus infection of big brown bats (*Eptesicus fuscus*). *J Wildl Dis* 44(3):612–621
- Jacob H, Solcher H (1968) An infectious disease transmitted by *Cercopithecus aethiops* (“marburg disease”) with glial nodule encephalitis. *Acta Neuropathol* 11(1):29
- Jahrling P, Geisbert T, Johnson E, Peters C, Dalgard D, Hall W (1990) Preliminary report: isolation of Ebola virus from monkeys imported to USA. *Lancet* 335(8688):502–505
- Jones BA, Grace D, Kock R et al (2013) Zoonosis emergence linked to agricultural intensification and environmental change. *Proc Natl Acad Sci* 110(21):8399–8404
- Kamins AO, Rowcliffe JM, Ntiama-Baidu Y, Cunningham AA, Wood JL, Restif O (2014) Characteristics and risk perceptions of Ghanaians potentially exposed to bat-borne zoonoses through bushmeat. *EcoHealth*:1–17
- Kingston T (2016) Cute, Creepy, or Crispy—how values, attitudes and norms shape human behavior toward bats. In: Voigt, CC, Kingston, T (eds) *Bats in the anthropocene: conservation of bats in a changing world*. Springer International AG, Cham, pp. 571–588
- Knight AJ (2008) “Bats, snakes and spiders, Oh my!” How aesthetic and negativistic attitudes, and other concepts predict support for species protection. *J Environ Psychol* 28(1):94–103
- Kurth A, Kohl C, Brinkmann A et al (2012) Novel paramyxoviruses in free-ranging European bats. *PLoS ONE* 7(6):e38688
- Kuzmin IV, Botvinkin AD, Rybin SN, Baialiev AB (1991) A lyssavirus with an unusual antigenic structure isolated from a bat in southern Kyrgyzstan. *Vopr Virusol* 37(5–6):256–259
- Kuzmin IV, Botvinkin AD, Khabilov TK (2001) The lyssavirus was isolated from a whiskered bat in northern Tajikistan. *Plecotus* et al 4:75–81
- Kuzmin IV, Niezgoda M, Carroll DS et al (2006) Lyssavirus surveillance in bats, Bangladesh. *Emerg Infect Dis* 12(3):486
- Kuzmin IV, Niezgoda M, Franka R et al (2008a) Possible emergence of West Caucasian bat virus in Africa. *Emerg Infect Dis* 14(12):1887
- Kuzmin IV, Niezgoda M, Franka R et al (2008b) Lagos bat virus in Kenya. *J Clin Microbiol* 46(4):1451–1461
- Kuzmin IV, Mayer AE, Niezgoda M et al (2010) Shimoni bat virus, a new representative of the Lyssavirus genus. *Virus Res* 149(2):197–210
- Kuzmin IV, Bozick B, Guagliardo SA et al (2011a) Bats, emerging infectious diseases, and the rabies paradigm revisited. *Emerg Health Threats* 4
- Kuzmin IV, Turmelle AS, Agwanda B et al (2011b) Commerson's leaf-nosed bat (*Hipposideros commersoni*) is the likely reservoir of Shimoni bat virus. *Vector-Borne Zoonotic Dis* 11(11):1465–1470
- Kuzmin IV (2014) Basic facts about Lyssaviruses. *Curr Lab Tech Rabies Diagn Res Prev* 1:3

- Lau SK, Woo PC, Wong BH et al (2010) Identification and complete genome analysis of three novel paramyxoviruses, Tuhoko virus 1, 2 and 3, in fruit bats from China. *Virology* 404(1):106–116
- Leonova GN, Belikov SI, Kondratov IG et al (2009) A fatal case of bat lyssavirus infection in Primorye territory of the Russian far East. *Rabies Bull Europe* 33(4):5–8
- Leroy EM, Rouquet P, Formenty P et al (2004) Multiple Ebola virus transmission events and rapid decline of central African wildlife. *Science* 303(5656):387–390
- Leroy EM, Kumulungui B, Pourrut X et al (2005) Fruit bats as reservoirs of Ebola virus. *Nature* 438(7068):575–576
- Leroy EM, Epelboin A, Mondonge V et al (2009) Human Ebola outbreak resulting from direct exposure to fruit bats in Luebo, democratic Republic of Congo, 2007. *Vector-Borne Zoonotic Dis* 9(6):723–728
- Li W, Shi Z, Yu M et al (2005) Bats are natural reservoirs of SARS-like coronaviruses. *Science* 310(5748):676–679
- Li Y, Chen S (2014) Evolutionary history of Ebola virus. *Epidemiol Infect* 142(06):1138–1145
- Luby SP, Rahman M, Hossain MJ et al (2006) Foodborne transmission of Nipah virus, Bangladesh. *Emerg Infect Dis* 12(12):1888
- Luby SP, Hossain MJ, Gurley ES et al (2009) Recurrent zoonotic transmission of Nipah virus into humans, Bangladesh, 2001–2007. *Emerg Infect Dis* 15(8):1229
- Luby SP, Gurley ES (2012) Epidemiology of henipavirus disease in humans. In: *Henipavirus*. Springer Berlin Heidelberg, pp. 25–40
- Luis AD, Hayman DT, O'Shea TJ et al (2013) A comparison of bats and rodents as reservoirs of zoonotic viruses: are bats special? *Proc R Soc B: Biol Sci* 280(1756):20122753
- Lumio J, Hillbom M, Roine R et al (1986) Human rabies of bat origin in Europe. *Lancet* 327(8477):378
- Lumlertdacha B, Boongird K, Wanghongsa S et al (2005) Survey for bat lyssaviruses, Thailand. *Emerg Infect Dis* 11(2):232
- Markotter W, Kuzmin IV, Rupprecht CE, Nel LH (2009) Lagos bat virus virulence in mice inoculated by the peripheral route. *Epidemiol Infect* 137(8):1155
- Marsh GA, Haining J, Robinson R et al (2011) Ebola Reston virus infection of: clinical significance and transmission potential. *J Infect Dis* 204(Suppl 3):S804–S809
- Marsh GA, De Jong C, Barr JA et al (2012) Cedar virus: a novel henipavirus isolated from Australian bats. *PLoS Pathog* 8(8):e1002836
- Marston DA, Horton DL, Ngeleja C et al (2012) Ikoma lyssavirus, highly divergent novel lyssavirus in an African civet. *Emerg Infect Dis* 18(4):664–667
- Martin G, Plowright R, Chen C, Kault D, Selleck P, Skerratt L (2015) Hendra virus survival does not explain spillover patterns and implicates relatively direct transmission routes from flying foxes to horses. *J Gen Virol* 96(6):1229–1237
- Mayen F (2003) Haematophagous bats in Brazil, their role in rabies transmission, impact on public health, livestock industry and alternatives to an indiscriminate reduction of bat population. *J Vet Med Ser B* 50(10):469–472
- McColl KA, Tordo N, Aguilar SA (2000) Bat lyssavirus infections. *Revue scientifique et technique (Int Office Epizootics)* 19(1):177–196
- McColl K, Chamberlain T, Lunt R, Newberry K, Middleton D, Westbury H (2002) Pathogenesis studies with Australian bat lyssavirus in grey-headed flying foxes (*Pteropus poliocephalus*). *Aust Vet J* 80(10):636–641
- Medellin RA (2003) Diversity and conservation of bats in Mexico: research priorities, strategies and Actions. *Wildl Soc Bull* 31:87–97
- Memish ZA, Mishra N, Olival KJ et al (2013) Middle East respiratory syndrome coronavirus in bats, Saudi Arabia. *Emerg Infect Dis* 19(11):1819
- Memish ZA, Cotten M, Meyer B et al (2014) Human infection with MERS coronavirus after exposure to infected camels, Saudi Arabia, 2013. *Emerg Infect Dis* 20(6):1012
- Meredith C, Prossouw A, Koch H (1971) An unusual case of human rabies thought to be of chiropteran origin. *South African Med J=Suid-Afrikaanse tydskrif vir geneeskde* 45(28):767

- Messenger SL, Rupprecht CE, Smith JS (2003) Bats, emerging virus infections, and the rabies paradigm. In: Kunz TH, Fenton MB (eds) Bat ecology. The University of Chicago Press, Chicago, pp 622–679
- Mickleburgh SP, Hutson AM, Racey PA (2002) A review of the global conservation status of bats. *Oryx* 36(01):18–34
- Mickleburgh S, Waylen K, Racey P (2009) Bats as bushmeat: a global review. *Oryx* 43(02):217–234
- Middleton DJ, Weingartl HM (2012) Henipaviruses in their natural animal hosts. In: Henipavirus. Springer Berlin Heidelberg, pp 105–121
- Mildenstein T, Tanshi I, Racey PA (2016) Exploitation of bats for bushmeat and medicine. In: Voigt, CC, Kingston, T (eds) Bats in the Anthropocene: conservation of bats in a changing world. Springer International AG, Cham, pp 325–376
- Misra V, Dumonceaux T, Dubois J et al (2009) Detection of polyoma and corona viruses in bats of Canada. *J Gen Virol* 90(8):2015–2022
- Müller T, Cox J, Peter W et al (2004) Spill-over of European bat Lyssavirus type 1 into a Stone Marten (*Martes foina*) in Germany. *J Vet Med Ser B* 51(2):49–54
- Müller MA, Paweska JT, Leman PA et al (2007) Coronavirus antibodies in African bat species. *Emerg Infect Dis* 13(9):1367
- Murray K, Selleck P, Hooper P et al (1995) A morbillivirus that caused fatal disease in horses and humans. *Science* 268(5207):94–97
- Nahar N, Sultana R, Gurley ES, Hossain MJ, Luby SP (2010) Date palm sap collection: exploring opportunities to prevent Nipah transmission. *EcoHealth* 7(2):196–203
- Nahar N, Mondal UK, Sultana R et al (2013) Piloting the use of indigenous methods to prevent Nipah virus infection by interrupting bats' access to date palm sap in Bangladesh. *Health Promot Int* 28(3):378–386
- Nathwani D, McIntyre P, White K et al (2003) Fatal human rabies caused by European bat Lyssavirus type 2a infection in Scotland. *Clin Infect Dis* 37(4):598–601
- Negredo A, Palacios G, Vázquez-Morón S et al (2011) Discovery of an ebolavirus-like filovirus in europe. *PLoS Pathog* 7:e1002304
- Negrete OA, Levroney EL, Aguilar HC et al (2005) EphrinB2 is the entry receptor for Nipah virus, an emergent deadly paramyxovirus. *Nature* 436(7049):401–405
- Nel LH (2001) Mokola virus: a brief review of the status quo. In Sixth SEARG meeting, Lilongwe/Malawi:80–86
- O'Shea TJ, Cryan PM, Cunningham AA et al (2014) Bat flight and zoonotic viruses. *Emerg Infect Dis* 20(5):741
- O'Sullivan J, Allworth A, Paterson D et al (1997) Fatal encephalitis due to novel paramyxovirus transmitted from horses. *Lancet* 349:93–95
- Olival KJ, Islam A, Yu M et al (2013) Ebola virus antibodies in fruit bats, Bangladesh. *Emerg Infect Dis* 19:270–273
- Olival KJ, Hayman DTS (2014) Filoviruses in bats: current knowledge and future directions. *Viruses* 6:1759–1788
- Parashar UD, Sunn LM, Ong F et al (2000) Case-control study of risk factors for human infection with a new zoonotic paramyxovirus, Nipah virus, during a 1998–1999 outbreak of severe encephalitis in Malaysia. *J Infect Dis* 181:1755–1759
- Paton NI, Leo YS, Zaki SR et al (1999) Outbreak of Nipah-virus infection among abattoir workers in Singapore. *Lancet* 354:1253–1256
- Pawan J (1959) Rabies in the vampire bat of Trinidad with special reference to the clinical course and the latency of infection. *Carib Med J* 21:137–156
- Paweska JT, Blumberg LH, Liebenberg C et al (2006) Fatal human infection with rabies-related Duvenhage virus, South Africa. *Emerg Infect Dis* 12:1965–1967
- Peel AJ, Feldman K, Hayman DTS et al (2010) Widespread seroprevalence to Lagos bat virus and henipavirus in straw coloured fruit bats (*Eidolon helvum*) across East Africa. In: Proceedings of the 2nd International Berlin bat meeting: bat biology and infectious diseases, pp 15–16

- Peel AJ, Sargan DR, Baker KS et al (2013) Continent-wide panmixia of an African fruit bat facilitates transmission of potentially zoonotic viruses. *Nat Commun* 4:2770
- Pernet O, Schneider BS, Beaty SM et al (2014) Evidence for henipavirus spillover into human populations in Africa. *Nat Commun* 5:5342
- Pfefferle S, Oppong S, Drexler JF et al (2009) Distant relatives of severe acute respiratory syndrome coronavirus and close relatives of human coronavirus 229E in bats, Ghana. *Emerg Infect Dis* 15:1377–1384
- Philbey AW, Kirkland PD, Ross AD et al (1998) An apparently new virus (family Paramyxoviridae) infectious for pigs, humans, and fruit bats. *Emerg Infect Dis* 4:269–271
- Pigott DM, Golding N, Mylne A et al (2014) Mapping the zoonotic niche of Ebola virus disease in Africa. *Elife* 3:e04395
- Plowright RK, Foley P, Field HE et al (2011) Urban habituation, ecological connectivity and epidemic dampening: the emergence of Hendra virus from flying foxes (*Pteropus* spp.). *Proc R Soc B* 278:3703–3712
- Plowright RK, Eby P, Hudson PJ et al (2015) Ecological dynamics of emerging bat virus spillover. *Proc R Soc B* 282(1798):20142124
- Pooley S, Fa JE, Nasi R (2015) No conservation silver lining to Ebola. *Conserv Biol* 29(3):965–967
- Poon L, Chu D, Chan K et al (2005) Identification of a novel coronavirus in bats. *J Virol* 79:2001–2009
- Pourrut X, Kumulungui B, Wittmann T et al (2005) The natural history of Ebola virus in Africa. *Microbes Infect/Inst Pasteur* 7:1005–1014
- Pourrut X, Souris M, Towner JS et al (2009) Large serological survey showing cocirculation of Ebola and Marburg viruses in Gabonese bat populations, and a high seroprevalence of both viruses in *Rousettus aegyptiacus*. *BMC Infect Dis* 9:159
- Puechmaille SJ, Verdeyroux P, Fuller H, Gouilh MA, Bekaert M, Teeling EC (2010) White-nose syndrome fungus (*Geomyces destructans*) in bat, France. *Emerg Infect Dis* 16(2):290
- Quammen D (2013) Spillover: animal infections and the next human pandemic. WW Norton & Company
- Quan P-L, Firth C, Street C et al (2010) Identification of a severe acute respiratory syndrome coronavirus-like virus in a leaf-nosed bat in Nigeria. *MBio* 1:e00208–e00210
- Racey PA, Hutson AM, Lina PHC (2012) Bat rabies, public health and european bat conservation. *Zoonoses Public Health (Special Issue—Bats)*. 60:58–68
- Racey PA (2015) The uniqueness of bats. In: Lin-Fa W, Cowled C (eds) *Bats and Viruses: A New Frontier of Emerging Infectious Diseases*. Wiley Blackwell. p. 1–22
- Rahman SA, Hassan SS, Olival KJ et al (2010) Characterization of Nipah virus from naturally infected *Pteropus vampyrus* bats, Malaysia. *Emerg Infect Dis* 16(12):1990
- Rahman MA, Hossain MJ, Sultana S et al (2012) Date palm sap linked to Nipah virus outbreak in Bangladesh, 2008. *Vector-Borne Zoonotic Dis* 12:65–72
- Reynes J-M, Counor D, Ong S et al (2005) Nipah virus in Lyle's flying foxes, Cambodia. *Emerg Infect Dis* 11:1042–1047
- Rihtarič D, Hostnik P, Steyer A, Grom J, Toplak I (2010) Identification of SARS-like coronaviruses in horseshoe bats (*Rhinolophus hipposideros*) in Slovenia. *Arch Virol* 155:507–514
- Roberts BJ, Catterall CP, Eby P, Kanowski J (2012) Long-distance and frequent movements of the flying-fox *Pteropus poliocephalus*: implications for management. *PLoS ONE* 7:e42532
- Rogers RJ, Douglas IC, Baldock FC et al (1996) Investigation of a second focus of equine morbillivirus infection in coastal Queensland. *Aust Vet J* 74(3):243–244
- Roine R, Hillbom M, Valle M et al (1988) Fatal encephalitis caused by a bat-borne rabies-related virus. Clinical findings. *Brain: a journal of neurology* 111:1505–1516
- Rønsholt L, Sørensen K, Bruschke C et al (1998) Clinically silent rabies infection in (zoo) bats. *Vet Rec* 142:519–520
- Rota PA, Oberste MS, Monroe SS et al (2003) Characterization of a novel coronavirus associated with severe acute respiratory syndrome. *Science* 300:1394–1399
- Roué SY, Nemoz M (2004) Unusual mortality in Schreiber's long fingered bat, at several nurseries. *Wild Dis Assoc Newsl* 14:4

- Ruiz M, Chávez CB (2010) Rabies in Latin America. *Neurol Res* 32:272–277
- Rupprecht CE, Gibbons RV (2004) Prophylaxis against rabies. *New Engl J Med* 351:2626–2635
- Saéz AM, Weiss S, Nowak K, Lapeyre V, Zimmermann F et al (2015) Investigating the zoonotic origin of the West African Ebola epidemic. *EMBO Mol Med* 7(1):17–23
- Salmón-Mulanovich G, Vásquez A, Albújar C, Guevara C (2009) Human rabies and rabies in vampire and nonvampire bat species, Southeastern Peru, 2007. *Emerg Inf Dis* 15:1308–1310
- Samaratunga H, Searle J, Hudson N (1998) Non-rabies Lyssavirus human encephalitis from fruit bats: Australian bat Lyssavirus (pteropid Lyssavirus) infection. *Neuropathol Appl Neurobiol* 24:331–335
- Sasaki M, Setiyono A, Handharyani E et al (2012) Molecular detection of a novel paramyxovirus in fruit bats from Indonesia. *Virol J* 9:240
- Schatz J, Fooks AR, McElhinney L et al (2013) Bat rabies surveillance in Europe. *Zoonoses Public Health* 60(1):22–34
- Schneeberger K, Cziráj GÁ, Voigt CC (2013) Measures of the constitutive immune system are linked to diet and roosting habits of neotropical bats. *PLoS ONE* 8:e54023
- Schneider L, Barnard B, Schneider H et al (1985) Application of monoclonal antibodies for epidemiological investigations and oral vaccination studies. In: *Rabies in the tropics*. Springer, Verlag
- Schneider MC, Romijn PC, Uieda W et al (2009) Rabies transmitted by vampire bats to humans: an emerging zoonotic disease in Latin America? *Rev panam de salud pública* 25:260–269
- Schnell MJ, McGettigan JP, Wirblich C, Papaneri A (2009) The cell biology of rabies virus: using stealth to reach the brain. *Nat Rev Microbiol* 8:51–61
- Selimov MA, Tatarov AG, Botvinkin AD et al (1989) Rabies-related Yuli virus; identification with a panel of monoclonal antibodies. *Acta Virol* 33(6):542–546
- Serra-Cobo J, Amengual B, Abellán C, Bourhy H (2002) European bat lyssavirus infection in Spanish bat populations. *Emerg Inf Dis* 8:413–420
- Sétien A, Brochier B, Tordo N et al (1998) Experimental rabies infection and oral vaccination in vampire bats (*Desmodus rotundus*). *Vaccine* 16:1122–1126
- Smith I, Wang LF (2013) Bats and their virome: an important source of emerging viruses capable of infecting humans. *Curr Opin Virol* 3(1):84–91
- Smith C, Skelly C, Kung N, Roberts B, Field H (2014) Flying-fox species density—a spatial risk factor for Hendra virus infection in horses in Eastern Australia. *PLoS ONE* 9(6):e99965
- Speare R, Skerratt L, Foster R et al (1997) Australian bat lyssavirus infection in three fruit bats from north Queensland. *Commun Dis Intell* 21:117–119
- Steece R, Altenbach JS (1989) Prevalence of rabies specific antibodies in the Mexican free-tailed bat (*Tadarida brasiliensis mexicana*) at Lava Cave, New Mexico. *J Wildl Dis* 25:490–496
- Streicker DG, Turmelle AS, Vonhof MJ et al (2010) Host phylogeny constrains cross-species emergence and establishment of rabies virus in bats. *Science* 329(5992):676–679
- Streicker DG, Recuenco S, Valderrama W et al (2012) Ecological and anthropogenic drivers of rabies exposure in vampire bats: implications for transmission and control. *Proc R Soc B* 279:3384–3392
- Swanepoel R, Leman PA, Burt FJ et al (1996) Experimental inoculation of plants and animals with Ebola virus. *Emerg Inf Dis* 2:321–325
- Tang X, Zhang J, Zhang S et al (2006) Prevalence and genetic diversity of coronaviruses in bats from China. *J Virol* 80:7481–7490
- Taniguchi S, Watanabe S, Masangkay JS et al (2011) Reston Ebolavirus antibodies in bats, the Philippines. *Emerg Infect Dis* 17:1559–1560
- Thiriet D (2011) Conservation shouldn't be a popularity contest. *Conversation*:1–4
- Tjørnehøj K, Fooks AR, Agerholm JS, Rønsholt L (2006) Natural and experimental infection of sheep with European bat lyssavirus type-1 of Danish bat origin. *J Comp Pathol* 134(2):190–201
- Tong S, Conrardy C, Ruone S et al (2009) Detection of novel SARS-like and other coronaviruses in bats from Kenya. *Emerg Inf Dis* 15:482–485

- Tong S, Zhu X, Li Y et al (2013) New world bats harbor diverse influenza A viruses. *PLoS Pathog* 9(10):e1003657
- Towner JS, Pourrut X, Albariño CG et al (2007) Marburg virus infection detected in a common African bat. *PLoS ONE* 2:e764
- Towner JS, Amman BR, Sealy TK et al (2009) Isolation of genetically diverse Marburg viruses from Egyptian fruit bats. *PLoS Pathog* 5:e1000536
- Trimarchi C, Debbie J (1977) Naturally occurring rabies virus and neutralizing antibody in two species of insectivorous bats of New York State. *J Wildl Dis* 13:366–369
- Turmelle A, Jackson F, Green D, McCracken G, Rupprecht C (2010) Host immunity to repeated rabies virus infection in big brown bats. *J Gen Virol* 91:2360–2366
- van Thiel P, Van den Hoek J, Eftimov F et al (2008) Fatal case of human rabies (Duvénhage virus) from a bat in Kenya: The Netherlands, December 2007. *Euro Surveill* 13:118
- van Thiel P-PA, de Bie RM, Eftimov F et al (2009) Fatal human rabies due to Duvénhage virus from a bat in Kenya: failure of treatment with coma-induction, ketamine, and antiviral drugs. *PLoS Negl Trop Dis* 3:e428
- Vidgen ME, de Jong C, Rose K et al (2015) Novel paramyxoviruses in Australian flying-fox populations support host-virus coevolution. *J Gen Virol* 96(7):1619–1625
- Virgin HW, Wherry EJ, Ahmed R (2009) Redefining chronic viral infection. *Cell* 138(1):30–50
- Vogel G (2014) Are bats spreading ebola across sub-saharan Africa? *Science* 344:140
- Voigt, CC, Phelps KL, Aguirre L, Schoeman MC, Vanitharani J, Zubaid A (2016) Bats and buildings: the conservation of synanthropic bats. In: Voigt, CC, Kingston, T (eds) *Bats in the Anthropocene: conservation of bats in a changing world*. Springer International AG, Cham, pp. 427–453
- Wacharapluesadee S, Lumlerdacha B, Boongird K et al (2005) Bat Nipah virus, Thailand. *Emerg Inf Dis* 11:1949–1951
- Warrilow D (2005) Australian bat lyssavirus: a recently discovered new rhabdovirus. In: Fu ZF (ed) *The world of Rhabdoviruses*. Springer, Berlin
- Watanabe S, Masangkay JS, Nagata NK et al (2010) Bat coronaviruses and experimental infection of bats, the Philippines. *Emerg Inf Dis* 16:1217–1223
- Wilkinson GS, South JM (2002) Life history, ecology and longevity in bats. *Aging Cell* 1(2):124–131
- Woodroffe R (1999) Managing disease threats to wild mammals. *Anim Conserv* 2(03):185–193
- World Health Organisation (2003) Summary of probable SARS cases with onset of illness from 1 November 2002 to 31 July 2003 (http://www.who.int/csr/sars/country/table2004_04_21)
- World Health Organization (2014) Ebola virus disease, Fact Sheet. (<http://www.who.int/mediacentre/factsheets/fs103/en>)
- Wibbelt G, Moore MS, Schountz T, Voigt CC (2010) Emerging diseases in Chiroptera: why bats? *Biol Lett* 6:438–440
- Woo PC, Lau SK, Li KS et al (2006) Molecular diversity of coronaviruses in bats. *Virology* 351:180–187
- Yaiw KC, Crameri G, Wang L et al (2007) Serological evidence of possible human infection with Tioman virus, a newly described paramyxovirus of bat origin. *J Infect Dis* 196:884–886
- Yaiw KC, Bingham J, Crameri G et al (2008) Tioman virus, a paramyxovirus of bat origin, causes mild disease in pigs and has a predilection for lymphoid tissues. *J Virol* 82:565–568
- Yang Y, Du L, Liu C et al (2014) Receptor usage and cell entry of bat coronavirus HKU4 provide insight into bat-to-human transmission of MERS coronavirus. *Proc Nat Acad Sci USA* 111:12516–12521
- Yob JM, Field H, Rashdi AM et al (2001) Nipah virus infection in bats (order Chiroptera) in peninsular Malaysia. *Emerg Inf Dis* 7:439–441
- Young PL, Halpin K, Selleck PW et al (1996) Serologic evidence for the presence in Pteropus bats of a paramyxovirus related to equine morbillivirus. *Emerg Inf Dis* 2:239–240
- Zaki AM, Van Boheemen S, Bestebroer TM et al (2012) Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *New Engl J Med* 367:1814–1820
- Zhang G, Cowled C, Shi Z et al (2013) Comparative analysis of bat genomes provides insight into the evolution of flight and immunity. *Science* 339(6118):456–460