

Complex Analysis Problem Set 3

D. Zack Garza

Sunday 29th March, 2020

Contents

1	Problems From Tie	2
1.1	1	2
	1.1.1 Solution	2
1.2	2	3
	1.2.1 Solution	3
1.3	3	4
	1.3.1 Solution	4
1.4	4	5
	1.4.1 Solution	5
1.5	5	5
1.6	6	5
1.7	7	5
1.8	8	5
1.9	9	6
	1.9.1 Solution	6
1.10	10	6
1.11	11	8
1.12	12	8
	1.12.1 Solution	8
1.13	13	10
1.14	14	10
	1.14.1 Solution	10
2	Stein and Shakarchi	10
2.1	S&S 3.8.1	10
2.2	S&S 3.8.2	10
2.3	S&S 3.8.4	11
2.4	S&S 3.8.5	11
2.5	S&S 3.8.6	11
2.6	S&S 3.8.7	11
2.7	S&S 3.8.8	11
2.8	S&S 3.8.9	12
2.9	S&S 3.8.10	12
2.10	S&S 3.8.14	12

2.11	S&S 3.8.15	12
2.11.1	a	12
2.11.2	b	12
2.11.3	c	12
2.11.4	d	13
2.12	S&S 3.8.17	13
2.12.1	a	13
2.12.2	b	13
2.13	S&S 3.8.19	13
2.13.1	a	13
2.13.2	b	13

1 Problems From Tie

1.1 1

Prove that if f has two Laurent series expansions,

$$f(z) = \sum c_n(z-a)^n \quad \text{and} \quad f(z) = \sum c'_n(z-a)^n$$

then $c_n = c'_n$.

1.1.1 Solution

By taking the difference of two such expansions, it suffices to show that if f is identically zero and $f(z) = \sum_{n=-\infty}^{\infty} c_n(z-a)^n$ about some point a , then $c_n = 0$ for all n .

Under this assumption, let $D_\varepsilon(a)$ be a disc about a and γ be any contour contained in its interior. Then for each n , we can apply the formula

$$\begin{aligned}
c_n &= \frac{1}{2\pi i} \int_{\gamma} \frac{f(\xi)}{(\xi-a)^{n+1}} d\xi \\
&= \frac{1}{2\pi i} \int_{\gamma} \frac{0}{(\xi-a)^{n+1}} d\xi \quad \text{by assumption} \\
&= 0,
\end{aligned}$$

which shows that $c_n = 0$ for all n . ■

1.2 2

Find Laurent series expansions of

$$\frac{1}{1-z^2} + \frac{1}{3-z}$$

How many such expansions are there? In what domains are each valid?

1.2.1 Solution

Note that f has poles at $z = -1, 1, 3$, all with multiplicity 1, and so there are 3 regions to consider:

1. $|z| < 1$
2. $1 < |z| < 3$
3. $3 < |z|$.



Region 1: Take the following expansion:

$$\begin{aligned} f(z) &= \frac{1}{1-z^2} + \frac{1}{3-z} \\ &= \sum_{n \geq 0} z^{2n} + \frac{1}{3} \left(\frac{1}{1 - \frac{z}{3}} \right) \\ &= \sum_{n \geq 0} z^{2n} + \frac{1}{3} \sum_{n \geq 0} \left(\frac{1}{3} \right)^n z^n \\ &= \sum_{n \geq 0} z^{2n} + \sum_{n \geq 0} \left(\frac{1}{3} \right)^{n+1} z^n \end{aligned}$$

Noting $|z^2| < 1$ implies then $|z| < 1$, and that the first term converges for $|z^2| < 1$ and the second for $\left| \frac{z}{3} \right| < 1 \iff |z| < 3$, this expansion converges to f on the region $|z| < 1$.

Region 2: Take the following expansion:

$$\begin{aligned}
 f(z) &= \frac{1}{1-z^2} + \frac{1}{3-z} \\
 &= -\frac{1}{z^2} \left(\frac{1}{1-\frac{1}{z^2}} \right) - \frac{1}{3} \left(\frac{1}{1-\frac{z}{3}} \right) \\
 &= -\frac{1}{z^2} \sum_{n \geq 0} z^{-2n} + \sum_{n \geq 0} \left(\frac{1}{3} \right)^{n+1} z^n \\
 &= -\sum_{n \geq 2} \frac{1}{z^{2n}} + \sum_{n \geq 0} \left(\frac{1}{3} \right)^{n+1} z^n
 \end{aligned}$$

By construction, the first term converges for $\left| \frac{1}{z^2} \right| < 1 \iff |z| > 1$ and the second for $|z| < 3$.

Region 3: Take the following expansion:

$$\begin{aligned}
 f(z) &= \frac{1}{1-z^2} + \frac{1}{3-z} \\
 &= -\frac{1}{z^2} \left(\frac{1}{1-\frac{1}{z^2}} \right) - \frac{1}{z} \left(\frac{1}{1-\frac{z}{3}} \right) \\
 &= -\frac{1}{z^2} \sum_{n \geq 0} \frac{1}{z^{2n}} - \frac{1}{z} \sum_{n \geq 0} 3^n \frac{1}{z^n} \\
 &= -\sum_{n \geq 2} \frac{1}{z^{2n}} - \sum_{n \geq 1} \left(\frac{1}{3} \right)^{n-1} \frac{1}{z^n}.
 \end{aligned}$$

Note: in principle, terms could be collected here.

By construction, this converges on $\{|z|^2 > 1\} \cap \{|z| > 3\} = \{|z| > 3\}$.

■

1.3 3

Let P, Q be polynomials with no common zeros. Assume a is a root of Q . Find the principal part of P/Q at $z = a$ in terms of P and Q if a is (1) a simple root, and (2) a double root.

1.3.1 Solution

todo

1.4 4

Let f be non-constant, analytic in $|z| > 0$, where $f(z_n) = 0$ for infinitely many points z_n with $\lim_{n \rightarrow \infty} z_n = 0$.

Show that $z = 0$ is an essential singularity for f .

Example: $f(z) = \sin(1/z)$.

1.4.1 Solution

It suffices to show that $z_0 = 0$ is neither a pole nor a removable singularity, i.e.

1. $\lim_{z \rightarrow z_0} f(z) \neq \infty$
2. $|f(z)|$ is not bounded on any neighborhood $D_\varepsilon(z_0)$.

The first property follows because if f is analytic,

1.5 5

Show that if f is entire and $\lim_{z \rightarrow \infty} f(z) = \infty$, then f is a polynomial.

1.6 6

Problem : a. Show that

$$\int_0^{2\pi} \log |1 - e^{i\theta}| d\theta = 0$$

b. Show that this identity is equivalent to SS 3.8.9.

1.7 7

Let $0 < a < 4$ and evaluate

$$\int_0^\infty \frac{x^{\alpha-1}}{1+x^3} dx$$

1.8 8

Prove the fundamental theorem of Algebra using

- a. Rouché's Theorem.
- b. The maximum modulus principle.

1.9 9

Let f be analytic in a region D and γ a rectifiable curve in D with interior in D .
Prove that if $f(z)$ is real for all $z \in \gamma$, then f is constant.

1.9.1 Solution

Since f is analytic in D (wlog assuming $0 \in D$ by translation), take its series expansion $f(z) = c_0 + c_1z + \cdots$ for $z \in D$.

Without loss of generality, suppose that γ is not entirely contained in \mathbb{R} , so for $z \in \gamma$ we can write $z = x + iy$ where $y \neq 0$.

Then

$$\begin{aligned} f(z) &= f(x + iy) \\ &= c_0 + c_1(x + iy) + \cdots \\ &= c_0 + c_1x + ic_1y + \cdots \end{aligned} \quad \subset \mathbb{R} \quad \text{by assumption,}$$

and so we must have $c_1y = 0 \implies c_1 = 0$. The same argument applies to further terms in the expansion, so we in fact have $c_i = 0$ for every $i \geq 1$.

But this says $f(z) = c_0$ for an arbitrary z , i.e. f is constant.

■

1.10 10

For $a > 0$, evaluate

$$\int_0^{\pi/2} \frac{d\theta}{a + \sin^2 \theta}$$

We have

$$\begin{aligned}
\int_0^{\pi/2} \frac{1}{1 + \sin^2(\theta)} d\theta &= \int_{\gamma_1} \frac{1}{a + \left(\frac{z-z^{-1}}{2i}\right)^2} \frac{-i dz}{z} \quad \text{where } \gamma_1 \text{ is } \frac{1}{4} \text{ of the unit circle } S^1 \\
&= -i \int_{\gamma_1} \frac{1}{z} \left(\frac{1}{a + \left(-\frac{1}{4}\right)(z^2 - 2 + z^{-2})} \right) dz \\
&= 4i \int_{\gamma_1} \frac{1}{z} \left(\frac{1}{z^2 - (2 + 4a) + z^{-2}} \right) dz \\
&= 4i \int_{\gamma_1} \frac{z}{z^4 - (2 + 4a)z^2 + 1} dz \\
&= i \oint_{S^1} \frac{z}{z^4 - (2 + 4a)z^2 + 1} dz \\
&= \frac{i}{2} \oint_{2 \cdot S^1} \frac{1}{u^2 - (2 + 4a)u + 1} du \quad \text{using } u = z^2, \frac{1}{2} du = z dz \\
&:= i \oint_{2 \cdot S^1} \frac{1}{f_a(u)} du \\
&= 2i \cdot 2\pi i \cdot \sum \text{Res}_{u=r_i} \frac{1}{f_a(u)},
\end{aligned}$$

where $2 \cdot S^1$ denotes the contour wrapping around the unit circle twice and r_i denote the poles contained in the region bounded by S^1 . We can now compute the last integral by the residue theorem.

Factor the denominator as

$$f_a(u) = u^2 - (2 + 4a)u + 1 = (u - r_1)(u - r_2),$$

where the r_i are given by $(1 + 2a) \pm 4\sqrt{a^2 + a}$ using the quadratic formula. We can then write a partial fraction decomposition

$$\begin{aligned}
\frac{1}{f_a(u)} &:= \frac{1}{u^2 - (2 + 4a)u + 1} \\
&= \frac{1}{(u - r_1)(u - r_2)} \\
&= \frac{A}{u - r_1} + \frac{B}{u - r_2} \\
&= \frac{\text{Res}_{u=r_1} 1/f(u)}{u - r_1} + \frac{\text{Res}_{u=r_2} 1/f(u)}{u - r_2} \\
&= \frac{1/f'(r_1)}{u - r_1} + \frac{1/f'(r_2)}{u - r_2} \\
&= \frac{1}{8\sqrt{a^2 + a}(u - r_1)} - \frac{1}{8\sqrt{a^2 + a}(u - r_2)}.
\end{aligned}$$

Since $|r_2| = \left| (1 + 2a) + 4\sqrt{a^2 + a} \right| > 1$, we find that the only relevant pole inside of S^1 is r_1 .

1.11 11


Find the number of roots of $p(z) = 4z^4 - 6z + 3$ in $|z| < 1$ and $1 < |z| < 2$ respectively.

1.12 12

Prove that $z^4 + 2z^3 - 2z + 10$ has exactly one root in each open quadrant.

1.12.1 Solution

Let $f(z) = z^4 + 2z^3 - 2z + 10$, and consider the following contour:



By the argument principle, we have

$$\Delta_{\Gamma} \arg f(z) = 2\pi(Z - P),$$

where Z is the number of zeros of f in the region Ω enclosed by Γ and P is the number of poles in Ω .

Since polynomials are holomorphic on \mathbb{C} , by the argument principle it suffices to show that

- f does not have any roots on the real or imaginary axes
- f does not vanish on Γ , and
- $\Delta_\Gamma \arg f(z) = 1$, where Δ_Γ denotes the total change in the argument of f over Γ .

It will follow by symmetry that f has exactly one root in each quadrant.

Claim 1.1.

- f has no roots on the coordinate axes.
- $\Delta_{\gamma_1} \arg f(z) = 0$
- $\Delta_{\gamma_2} \arg f(z) = 2\pi$
- $\Delta_{\gamma_3} \arg f(z) = 0$

Given the claim, we would have

$$\Delta_\Gamma \arg f(z) = 2\pi = 2\pi(Z - 0) \implies Z = 1,$$

which is what we wanted to show.

Proof of Claim:

γ_2 : For $R \gg 0$, we have $f(z) \sim z^4$. Along γ_2 , the argument of z ranges from 0 to $\frac{\pi}{2}$, and thus the argument of z^4 ranges from 0 to $4 \cdot \frac{\pi}{2} = 2\pi$.

γ_1 : By cases, for $z \in \mathbb{R}$,

- If $|z| > 1$, then $z^3 > z$ and so

$$\begin{aligned} f(z) &= (z^4 + 10) + (2z^3 - 2z) \\ &> (z^4 + 10) + (2z - 2z) \\ &= z^4 + 10 \\ &> 0, \end{aligned}$$

so f is strictly positive and does not change argument on $(\pm 1, \pm\infty)$ or $i \cdot (\pm 1, \pm\infty)$.

- If $|z| \leq 1$,

$$\begin{aligned} \left| -z^4 - 2z^3 + 2z \right| &\leq |z|^4 + 2|z|^3 + 2|z| \\ &\leq 1 + 2 + 2 \\ &= 5 \\ &< 10 \\ \implies f(z) &= 10 - (-z^4 - 2z^3 + 2z) > 0, \end{aligned}$$

so f is strictly positive and does not change argument $(0, \pm 1)$ or $i \cdot (0, \pm 1)$.

■

1.13 13

Prove that for $a > 0$, $z \tan z - a$ has only real roots.

1.14 14

Let f be nonzero, analytic on a bounded region Ω and continuous on its closure $\bar{\Omega}$. Show that if $|f(z)| \equiv M$ is constant for $z \in \partial\Omega$, then $f(z) \equiv Me^{i\theta}$ for some real constant θ .

1.14.1 Solution

By the maximum modulus principle applied to f in $\bar{\Omega}$, we know that $\max |f| = M$. Similarly, the maximum modulus principle applied to $\frac{1}{f}$ in $\bar{\Omega}^c$ since f is nonzero in Ω , and we can conclude that $\min |f| = M$ as well. Thus $|f| = M$ is constant on $\bar{\Omega}$.

So consider the function $g(z) = |f(z)|$; from the above observation, we find that $g(\bar{\Omega}) = \{M\}$. Letting S_M be the circle of radius M , this implies that $f(\Omega) \subseteq S_M$. In particular, $S_M \subset \mathbb{C}$ is a closed set.

However, by the open mapping theorem, $f(\Omega) \subset \mathbb{C}$ must be an open set. A basis for the topology on \mathbb{C} is given by open discs, so in particular, the open sets of \mathbb{C} have real dimension either zero or two. Since S_M has real dimension 1, $f(\Omega)$ must have dimension zero and is thus a collection of points. Since f is continuous, the image can only be one point, i.e. $f(\Omega) = \{\text{pt}\} \in S_M$. So f is constant. ■

2 Stein and Shakarchi**2.1 S&S 3.8.1**

Using Euler's formula

$$\sin(\pi z) = \frac{1}{2i}(e^{i\pi z} - e^{-i\pi z})$$

show that the complex zeros of $\sin(\pi z)$ are exactly the integers, each of order one.

Calculate the residue of $\frac{1}{\sin(\pi z)}$ at $z = n \in \mathbb{Z}$.

2.2 S&S 3.8.2

Evaluate the integral

$$\int_{\mathbb{R}} \frac{dx}{1+x^4}$$

What are the poles of the integrand?

2.3 S&S 3.8.4

Show that

$$\int_{\mathbb{R}} \frac{x \sin x}{x^2 + a^2} = \frac{\pi e^{-a}}{a} \quad a > 0$$

2.4 S&S 3.8.5

Show that for $\xi \in \mathbb{R}$,

$$\int_{\mathbb{R}} \frac{e^{2\pi i x \xi}}{(1 + x^2)^2} = \frac{\pi}{2} (1 + 2\pi |\xi|) e^{-2\pi |\xi|}$$

2.5 S&S 3.8.6

Show that

$$\int_{\mathbb{R}} \frac{dx}{(1 + x^2)^{n+1}} = \frac{1 \cdot 3 \cdots (2n - 1)\pi}{2 \cdot 4 \cdots (2n)}$$

2.6 S&S 3.8.7

Show that for $a > 1$,

$$\int_0^{2\pi} \frac{d\theta}{(a + \cos \theta)^2} = \frac{2\pi a}{(a^2 - 1)^{3/2}}$$

2.7 S&S 3.8.8

Show that if $a, b \in \mathbb{R}$ with $a > |b|$ then

$$\int_0^{2\pi} \frac{d\theta}{a + b \cos \theta} = \frac{2\pi a}{\sqrt{a^2 - b^2}}$$

2.8 S&S 3.8.9

Show that

$$\int_0^1 \log(\sin \pi x) \, dx = -\log 2$$

2.9 S&S 3.8.10

Show that if $a > 0$

$$\int_0^\infty \frac{\log x}{x^2 + a^2} \, dx = \frac{\pi \log a}{2a}$$

2.10 S&S 3.8.14

Prove that if f is entire and injective, then $f(z) = az + b$ with $a, b \in \mathbb{C}$ with $a \neq 0$.

Hint: apply the Casorati-Weierstrass theorem to $f(1/z)$.

2.11 S&S 3.8.15

Use the Cauchy inequalities or the maximum modulus principle to solve the following problems:

2.11.1 a

If f is entire and for all $R > 0$, there are constants $A, B > 0$ such that $\sup_{|z|=R} |f(z)| \leq AR^k + B$, then f is a polynomial of degree less than k .

2.11.2 b

Show that if f is holomorphic on the unit disk, is bounded, and converges to zero uniformly in the sector $\theta \leq \arg z \leq \phi$ as $|z| \rightarrow 1$, then $f \equiv 0$.

2.11.3 c

Let w_1, \dots, w_n be points on $S^1 \subset \mathbb{C}$. Show that there exists a point $z \in S^1$ such that

$$\prod_{i=1}^n |z - w_i| \geq 1.$$

Conclude that there exists a point $w \in S^1$ such that

$$\prod_{i=1}^n |w - w_i| = 1.$$

2.11.4 d

Show that if f is entire and $\Re(f)$ is bounded, then f is constant.

2.12 S&S 3.8.17

Let f be non-constant, and holomorphic in an open set containing the open unit disc.

2.12.1 a

Show that $|z| = 1 \implies |f(z)| = 1$, then the image of f contains the unit disc.

Hint: Show that $f(z) = w_0$ has a root for every $w_0 \in \mathbb{D}$, for which it suffices to show that $f(z) = 0$ has a root, then use the maximum modulus principle.

2.12.2 b

Show that if $|z| \geq 1 \implies |f(z)| = 1$ **and** there exists a point $z_0 \in \mathbb{D}$ such that $|f(z_0)| < 1$, then the image of f contains the unit disc.

2.13 S&S 3.8.19

Prove the maximum modulus principle for harmonic functions; i.e.,

2.13.1 a

If u is a non-constant real-valued harmonic function on Ω , then u can not attain its extrema on Ω .

2.13.2 b

Suppose Ω has compact closure $\overline{\Omega}$, then if u is harmonic on Ω and continuous on $\overline{\Omega}$, then

$$\sup_{z \in \Omega} |u(z)| \leq \sup_{z \in \overline{\Omega} - \Omega} |u(z)|$$

Hint: to prove (a), assume that u attains a local maximum at z_0 , and let f be holomorphic near z_0 with $u = \Re(f)$, then show that f is not open. Part (b) is a direct consequence.