

# Title

D. Zack Garza

August 21, 2019

## Contents

0.1 Exercises . . . . .	1
-------------------------	---

### 0.1 Exercises

#### Problem 1.

Let  $C$  denote the Cantor set.

1. Show that  $C$  contains point that is not an endpoint of one of the removed intervals.
2. Show that  $C$  is nowhere dense, meager, and has measure zero.
3. Show that  $C$  is uncountable.

#### Solution 1.

1. First we will characterize the endpoints of the removed intervals. Let  $C_n$  be the  $n$ th stage of the deletion process that is used to define the Cantor set; then what remains is a union of intervals:

$$C_n = [0, \frac{1}{3^n}] \cup [\frac{2}{3^n}, \frac{3}{3^n}] \cup \cdots \cup [\frac{3^n - 1}{3^n}, 1],$$

and so the endpoints are precisely the numbers of the form  $\frac{k}{3^n}$  where  $0 \leq k \leq 3^n$ . Moreover, any endpoint appearing in  $C_n$  is never removed in any later step, and so all endpoints remaining in  $C$  are of this form where we allow  $0 \leq n < \infty$ .

Thus, our goal is to produce a number  $x \in [0, 1]$  such that  $x \neq \frac{k}{3^n}$  for any  $k$  or  $n$ , but also satisfies  $x \in C$ . So we will need a general characterization of all of the points in  $C$ .

---

Lemma: If  $x \in C$ , then one can find a ternary expansion for which all of the digits are either 0 or 2, i.e.

$$x = \sum_{k=1}^{\infty} a_k 3^{-k} \quad \text{where } a_k \in \{0, 2\}.$$

Proof: By induction on the index  $k$  in  $a_k$ , first consider note that if  $x \in C$  then  $x \in C_1 = [0, 1] \setminus [\frac{1}{3}, \frac{2}{3}] = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$ . So if  $x \in C_1$ , then  $x \notin (\frac{1}{3}, \frac{2}{3})$ . But note that  $a_1$  is computed in the following way:

$$a_1 = \begin{cases} 0 & 0 \leq x < \frac{1}{3}, \\ 1 & \frac{1}{3} \leq x < \frac{2}{3}, \\ 2 & \frac{2}{3} \leq x < 1. \end{cases}$$

Since the interval  $(\frac{1}{3}, \frac{2}{3})$  is deleted in  $C_1$ , we find that  $a_1 = 1 \iff x = \frac{1}{3}$ . In this case, however, we claim that we can find a ternary expansion of  $x$  that does not contain a 1. We first write

$$x = \frac{1}{3} = \sum_{k=1}^{\infty} a_k 3^{-k} \quad \text{where } a_1 = 1, a_{k>1} = 0,$$

and then define

$$x' = \sum_{k=1}^{\infty} b_k 3^{-k} \quad \text{where } b_1 = 0, b_{k>1} = 2.$$

The claim now is that  $x = x'$ , which follows from the fact that this is a geometric sum that can be written in closed form:

$$\begin{aligned} x' &= \sum_{k=2}^{\infty} (2) 3^{-k} \\ &= \left( \sum_{k=0}^{\infty} (2) 3^{-k} \right) - 2 - 2(3^{-1}) \\ &= 2 \left( \sum_{k=0}^{\infty} 3^{-k} \right) - 2 - 2(3^{-1}) \\ &= 2 \left( \frac{1}{1 - \frac{1}{3}} \right) - 2 - 2(3^{-1}) \\ &= 2 \left( \frac{3}{2} \right) - 2 - 2(3^{-1}) \\ &= 1 - \frac{2}{3} \\ &= \frac{1}{3} = x. \end{aligned}$$

In short, we have  $\frac{1}{3} = (0.1)_3 = (0.222\cdots)_3$  as ternary expansions, and a similar proof shows that such an expansion without 1s can be found for any endpoint.

For the inductive step, consider  $a_n$ : the claim is that if  $a_n = 1$ , then  $x \notin C_{n+1}$  – that is, it is contained in one of the intervals deleted at the  $n + 1$ st stage. Writing the deleted interval at this stage as  $(a, b)$ , we find that  $a_n = 1$  if and only if  $x \in [a, b)$ . Since  $x \in C$ , the only way  $a_n$  can be 1 is if  $x$  was in fact the endpoint  $a$  (since no previous digit was a 1, by hypothesis). However, as shown above, every such endpoint has a ternary expansion containing no 1s.  $\square$

---

Therefore, if we can produce an  $x$  that satisfies  $x \neq \frac{k}{3^n}$  for any  $k, n$  **and**  $x$  has no 1s in its ternary expansion, we will have an  $x \in C$  that is not an endpoint.

So take

$$x = (0.\overline{02})_3 = (0.020202\cdots)_3.$$

This evidently has no 1s in its ternary expansion, and if we sum the corresponding geometric series, we find  $x = \frac{1}{4}$ . This is not of the form  $\frac{k}{3^n}$  for any  $k, n$ , and thus fulfills both conditions.

2. We first show that  $C$  is nowhere dense by showing that the interior of its closure is empty, i.e.  $(\overline{C})^\circ = \emptyset$ .

To do so, we note that  $C$  is itself closed and so  $C = \overline{C}$ . To see why this is, consider  $C^c$ ; we'll show that it is open. By construction,  $C_1^c$  is the open interval  $(\frac{1}{3}, \frac{2}{3})$  that is deleted, and similarly  $C_n^c$  is the finite union of the open intervals that are deleted at the  $n$ th stage. But then

$$C^c = \left(\bigcap C_n\right)^c = \bigcup C_n^c$$

is an infinite union of open sets, which is also open. So  $C$  is closed.