

# Title

*D. Zack Garza*

# Contents

1	Lecture 15: The $L$ -Polynomial	3
---	---------------------------------	---

# 1 | Lecture 15: The $L$ -Polynomial

Recall that we had  $Z(t) + F(t) + G(t)$ :

$$(q-1)F(t) = \sum_{0 \leq \deg C \leq 2g-2} q^{\ell(C)} t^{\deg(C)}$$

$$(q-1)G(t) = h \left( \frac{q^g t^{2g-1}}{1-qt} - \frac{1}{1-t} \right).$$

Note that  $F(t)$  is a polynomial of degree at most  $2g-2$ , and clearing denominators in  $G(t)$  yields a polynomial of degree at most  $2g$

**Definition 1.0.1** (The  $L$ -polynomial)

The  $L$ -polynomial is defined as

$$L(t) := (1-t)(1-qt)Z(t) = (1-t)(1-qt) \sum_{n=0}^{\infty} A_n t^n \in \mathbb{Z}[t].$$

It turns out that the degree bound of  $2g$  is sharp, and the coefficients closer to the middle are most interesting:

**Theorem 1.0.2 (?)**.

Let  $K/\mathbb{F}_q$  be a function field of genus  $g \geq 1$ , then

- $\deg L = 2g$ .
- $L(1) = h$
- $L(t) = q^g t^{2g} L\left(\frac{1}{qt}\right)$ .
- Writing  $L(t) = \sum_{j=1}^{2g} a_j t^j$ ,
  - $a_0 = 1$  and  $a_{2g} = q^g$ .
  - For all  $0 \leq j \leq g$ , we have  $a_{2g-j} = q^{g-j} a_j$ .
  - $a_1 = |\Sigma(K/\mathbb{F}_q)| - (q+1)$ , which notably does not depend on  $g$ .
  - Write  $L(t) = \prod_{j=1}^{2g} (1 - \alpha_j t) \in \mathbb{C}[t]$  <sup>a</sup>
- The  $\alpha_j \in \bar{\mathbb{Z}}$  <sup>b</sup> (which were *a priori* in  $\mathbb{C}$ ) and can be ordered such that for all  $1 \leq j \leq g$ , we have  $a_j a_{g+j} = q$ . <sup>c</sup>

f. If  $L_r(t) = (1-t)(1-q^r t)Z_r(t)$  then  $L_r(t) = \prod_{j=1}^{2g} (1 - \alpha_j^r t)$ , where  $K_r$  is the constant extension  $K\mathbb{F}_{q^r}/\mathbb{F}_{q^r}$

<sup>a</sup>The polynomial isn't monic, but rather has a constant coefficient, so this expansion is somewhat more natural than (say)  $\prod (t - \alpha)$ .

<sup>b</sup> $\bar{\mathbb{Z}}$  denotes the algebraic integers.

<sup>c</sup>This is the first hint at the Riemann hypothesis: if for example they all had the same complex modulus, this would force  $|a_j| = \sqrt{q}$ . Thus proving that they all have the same absolute value is 99% of the content!

Note that the  $\alpha_i$  are reciprocal roots.

*Proof (of a).*

We saw from  $Z(t) = F(t) + G(t)$  that  $\deg L \leq 2g$ . Equality will follow from the proof of (d) part 1, since this would imply that  $a_{2g} = q^g \neq 0$ . ■

*Proof (of b).*

Our formula  $Z(t) = F(t) + G(t)$  and Schmidt's theorem (showing  $\delta = 1$ ) gives

$$L(t) = (1-t)(1-qt)F(t) + \frac{h}{q-1} \left( q^g t^{2g-2} (1-t) - (1-qt) \right),$$

where we've expanded  $G$  but not  $F$  because it involves various  $\ell(D)$  which are difficult to compute. It is some polynomial though, and we can evaluate  $L$  at 1 to get  $L(1) = h$ . Thus the class number is the sum of the coefficients! ■

*Proof (of c).*

This follows easily from the functional equation for  $Z(t)$ , which we already established using the Riemann-Roch theorem:

$$Z(t) = q^{g-1} t^{2g-2} Z\left(\frac{1}{qt}\right).$$

We can compute

$$\begin{aligned} q^g t^{2g} L\left(\frac{1}{qt}\right) &= q^g t^{2g} \left(1 - \frac{1}{qt}\right) \left(1 - \frac{1}{t}\right) Z\left(\frac{1}{qt}\right) \\ &= q^{g-1} t^{2g-2} (1-t)(1-qt) Z\left(\frac{1}{qt}\right) \\ &= (1-t)(1-qt) Z(t) \\ &:= L(t), \end{aligned}$$

where we've distributed one  $q$  and two  $t$ s in the first steps. ■

*Proof (of  $d$ ).*

Using the functional equation from (c), we can write

$$L(t) = q^g t^{2g} L\left(\frac{1}{qt}\right) = \left(\frac{a_{2g}}{q^g}\right) + \left(\frac{a_{2g-1}}{q^{g-1}}\right)t + \cdots + (a_0 q^g) t^{2g},$$

where we're correcting by enough in  $t$  but not enough in  $q$  and seeing what we get. Equating coefficients, for  $0 \leq j \leq g$  we have

$$a_{2g-j} = q^{g-j} a_j. \quad (1)$$

Using the fact that  $A_0$  is the number of effective degree zero divisors, which is only zero, we have  $A_0 = 1$  and we can multiply formal power series to obtain

$$\begin{aligned} L(t) &= a_0 + a_1 t + \cdots + a_{2g} t^{2g} = (1-t)(1-qt) \sum_{n=0}^{\infty} A_n t^n \\ &= \left(1 - (q+1)t + qt^2\right) (1 + A_1 t + A_2 t^2 + \cdots) \\ &= 1 + (A_1 - (q+1))t + \cdots \end{aligned}$$

From this, we can read off

- $L(0) = a_0 = 1$
- $a_1 = A_1 - (q+1) = \Sigma_1(K/k) - (q+1)$
- $a_{2g} = a_{2g-0} = q^{g-0} a_0 = a^g$  by taking  $j = 0$  in eq. 1, and thus  $\deg L = 2g$ .

■

*Proof (of  $e$  (the most interesting!)).*

Consider the **reciprocal polynomial**

$$L^\perp(t) := t^{2g} L\left(\frac{1}{t}\right) = t^{2g} + a_1 t^{2g-1} + \cdots + q^g.$$

The original polynomial had  $\mathbb{Z}$  coefficients and constant term 1, so this polynomial is monic and has a nonzero constant term. Thus its roots are patently nonzero algebraic integers in  $\overline{\mathbb{Z}}^\bullet$ .

If  $L^\perp(t) = \prod_{j=1}^{2g} (t - \alpha_j)$ , then

$$L(t) = t^{2g} L^\perp\left(\frac{1}{t}\right) = \prod_{j=1}^{2g} (1 - \alpha_j t)$$

and if the roots of  $L(t)$  are  $r_j$ , then the roots of  $L^\perp(t)$  are the reciprocal roots  $1/r_j$  and vice-versa. This shows the first assertion that  $r_j \in \overline{\mathbb{Z}}$  as well.

■