

Title

D. Zack Garza

Sunday 13th September, 2020

Contents

1	Sunday, September 13	1
1.1	General Notes	1
1.2	1.a	1
1.3	3.a	2

1 | Sunday, September 13

1.1 General Notes

- If flipping logic and not using a direct proof (contradiction, contrapositive, etc), then signpost/announce it near the beginning of the proof.
- Say what you're assuming at the start of the proof.
- Put any important equations (i.e. major steps of the proof) on their own lines or in `displaymath` environments.
- Use some whitespace to separate parts of the proof and increase readability.
- Remember that limits of sequences need not exist, but \liminf s/ \limsup s always do (just may be $\pm\infty$).
- Try to avoid abbreviating the names of major theorems (example: "AP" can stand for many results, not just the Archimedean property!)
- It's not generally true that $a \leq M \implies |a| \leq M$, e.g. take $a = -1$. This only holds $a \geq 0$.

1.2 1.a

Proof ($A \implies B$).

- Suppose $\{a_n\}$ is not bounded above.
- Then any $k \in \mathbb{N}$ is not an upper bound for $\{a_n\}$.

- So choose a subsequence $a_{n_k} > k$, then by order-limit laws,

$$a_{n_k} > k \implies \liminf_{k \rightarrow \infty} a_{n_k} > \liminf_{k \rightarrow \infty} k = \infty.$$

■

Proof ($\neg A \implies \neg B$).

- Suppose $\{a_n\}$ is bounded by M , so $a_n < M < \infty$ for all $n \in \mathbb{N}$.
- Then if $\{a_{n_k}\}$ is a subsequence, we have $a_{n_k} \in \{a_n\}$, so $a_{n_k} < M$ for all $k \in \mathbb{N}$.
- But then

$$a_{n_k} < M \implies \limsup_{k \rightarrow \infty} a_{n_k} \leq M,$$

- Now note that if $\lim_{k \rightarrow \infty} a_{n_k}$ exists,

$$\lim_{k \rightarrow \infty} a_{n_k} < \limsup_{k \rightarrow \infty} a_{n_k} \leq M < \infty,$$

so every subsequence is bounded and thus can not converge to ∞ .

■

1.3 3.a

Proof (Using definition (i)).

- Suppose $x_n \leq M$ for all n , we will show that every subsequential limit is also bounded by M .
- Let

$$S := \left\{ x \in \mathbb{R} \mid x \text{ is a subsequential limit of } \{x_n\} \right\}$$

be the set of subsequential limits.

- Note that $\inf S := \liminf_{n \rightarrow \infty} x_n$ by definition (i).
- Let $\{x_{n_k}\} \in S$ be an arbitrary convergent subsequence (since we are only concerned about subsequences with well-defined limits).
- Then for every k we have $x_{n_k} \in \{x_n\}$, so

$$|x_{n_k}| \leq M.$$

- By order limit laws,

$$|x_{n_k}| \leq M \implies \lim_{k \rightarrow \infty} |x_{n_k}| \leq M,$$

- Since the map $x \mapsto |x|$ is continuous, using the sequential definition of continuity we can pass the limit through the absolute value to obtain

$$\left| \lim_{k \rightarrow \infty} x_{n_k} \right| \leq M.$$

- Since the subsequence was arbitrary, we find that M is an upper bound for S and so $\sup S \leq M$.
- But

$$\inf S \leq \sup S \leq M \implies \inf S \leq M.$$

■

Proof (Using definition (ii)).

- Suppose $|x_n| \leq M$ for every n , we will directly show that $\left| \liminf_{n \rightarrow \infty} x_n \right| \leq M$.
- Let $\{x_{n_k}\}$ be an arbitrary subsequence, then since $x_{n_k} \in \{x_n\}$ for all k , $|x_{n_k}| \leq M$ for all k .
- By order-limit laws, for every fixed n we have

$$|x_{n_k}| \leq M \iff -M \leq x_{n_k} \leq M \implies -M \leq \inf_{k > n} x_{n_k} \leq M.$$

- Again applying order-limit laws,

$$-M \leq \inf_{k > n} x_{n_k} \leq M \implies -M \leq \lim_{n \rightarrow \infty} \inf_{k > n} x_{n_k} \leq M \iff \left| \lim_{n \rightarrow \infty} \inf_{k \geq n} x_{n_k} \right| \leq M.$$

- But by definition (i), this precisely says that $\left| \liminf_{n \rightarrow \infty} x_n \right| \leq M$.

■