

Title

D. Zack Garza

August 19, 2019

Contents

0.1 Exercises	1
0.2 Qual Problems	2

0.1 Exercises

Problem 1 (Hungerford 1.6.3).

If $\sigma = (i_1 i_2 \cdots i_r) \in S_n$ and $\tau \in S_n$, then show that $\tau\sigma\tau^{-1} = (\tau(i_1)\tau(i_2) \cdots \tau(i_r))$.

Solution 1. Let

$$\sigma = (s_1 s_2 \cdots s_r) \in S_n$$

in cycle notation, and $\tau \in S_n$ be arbitrary. Define $t_j = \tau(s_j)$; we would then like to show that

$$(t_1, t_2, \cdots t_r) := (\tau(s_1)\tau(s_2) \cdots \tau(s_r)) = \tau\sigma\tau^{-1}$$

To this end, it suffices to show that t_i maps to $t_{i+1 \bmod r}$, under $\tau\sigma\tau^{-1}$, which is to say

$$\tau\sigma\tau^{-1}(t_i) = \begin{cases} t_{i+1} & i+1 \leq r, \\ t_1 & i = r \end{cases}.$$

Bearing this in mind, we will immediately suppress notation and take all indices $\bmod r$ for the rest of this problem.

The following then follows simply by definitions:

$$\begin{aligned} \tau\sigma\tau^{-1}(t_i) &= \tau\sigma(s_i) \\ &= \tau(s_{i+1}) \\ &= t_{i+1}. \end{aligned}$$

□

Problem 2 (Hungerford 1.6.4).

Show that $S_n \cong \langle (12), (123 \cdots n) \rangle$ and also that $S_n \cong \langle (12), (23 \cdots n) \rangle$

Problem 3 (Hungerford 2.2.1).

Let G be a finite abelian group that is not cyclic. Show that G contains a subgroup isomorphic to $\mathbb{Z}_p \oplus \mathbb{Z}_p$ for some prime p .

Problem 4 (Hungerford 2.2.12.b).

Determine all abelian groups of order n for $n \leq 20$.

Problem 5 (Hungerford 2.4.1).

Let G be a group and $A \trianglelefteq G$ be a normal abelian subgroup. Show that G/A acts on A by conjugation and construct a homomorphism $\varphi : G/A \rightarrow \text{Aut}(A)$.

Problem 6 (Hungerford 2.4.9).

Let $Z(G)$ be the center of G . Show that if $G/Z(G)$ is cyclic, then G is abelian.

Note that Hungerford uses the notation $C(G)$ for the center.

Problem 7 (Hungerford 2.5.6).

Let G be a finite group and $H \trianglelefteq G$ a normal subgroup of order p^k . Show that H is contained in every Sylow p -subgroup of G .

Problem 8 (Hungerford 2.5.9).

Let $|G| = p^n q$ for some primes $p > q$. Show that G contains a unique normal subgroup of index q .

0.2 Qual Problems**Problem 9.**

Let G be a finite group and p a prime number. Let X_p be the set of Sylow- p subgroups of G and n_p be the cardinality of X_p . Let $\text{Sym}(X)$ be the permutation group on the set X .

1. Construct a homomorphism $\rho : G \rightarrow \text{Sym}(X_p)$ with image a transitive subgroup (i.e. with a single orbit).
2. Deduce that G is simple and the order of G divides $n_p!$.
3. Show that for any $1 \leq a \leq 4$ and any prime power p^k , no group of order ap^k is simple.

Problem 10.

Let G be a finite group and $H < G$ a subgroup. Let n_H be the number of subgroups of G that are conjugate to H . Show that n_H divides the order of G .

Problem 11.

Let $G = S_5$, the symmetric group on 5 elements. Identify all conjugacy classes of elements in G , provide a representative from each class, and prove that this list is complete.