

Homework 7

D. Zack Garza

November 5, 2019

Contents

1	Problem 1	1
1.1	Part 1	1
1.2	Part 2	2
2	Problem 2	3
2.1	Part 1	3
2.2	Part 2	3
3	Problem 3	3
3.1	Part 1	3
4	Problem 4	5

1 Problem 1

1.1 Part 1

In order for IS to be a submodule of A , we need to show the following implication:

$$x \in IS, a \in A \implies xa, ax \in IS.$$

Suppose $x \in IS$. Then by definition, $x = \sum_{i=1}^n r_i a_i$ for some $r_i \in R, a_i \in A$.

But then

$$\begin{aligned} xa &= \left(\sum_{i=1}^n r_i a_i \right) a \\ &= \sum_{i=1}^n r_i a_i a \\ &:= \sum_{i=1}^n r_i a'_i, \end{aligned}$$

where $a'_i := a_i a$ for each i , which is still an element of A since A itself is a module and thus closed under multiplication.

But this expresses xa as an element of IS . Similarly, we have

$$\begin{aligned} ax &= a \left(\sum_{i=1}^n r_i a_i \right) \\ &= \sum_{i=1}^n a r_i a_i a \\ &:= \sum_{i=1}^n r_i a a_i, \\ &:= \sum_{i=1}^n r_i a'_i, \end{aligned}$$

and so $ax \in IS$ as well.

1.2 Part 2

Letting $R/I \curvearrowright A/IA$ be the action given by $r + I \curvearrowright +IA := ra + IA$, we need to show the following:

- $r.(x + y) = r.x + r.y$,
- $(r + r').x = r.x + r'.x$,
- $(rs).x = r.(s.x)$, and
- $1.x = x$.

Letting \oplus denote the addition defined on cosets, we have

$$\begin{aligned} r \curvearrowright (x + IA \oplus y + IA) &:= r \curvearrowright x + y + IA \\ &:= r(x + y) + IA \\ &= rx + ry + IA \\ &:= rx + IA \oplus ry + IA \\ &:= (r \curvearrowright x + IA) \oplus (r \curvearrowright y + IA). \end{aligned}$$

$$\begin{aligned} (r + s) \curvearrowright x + IA &:= (r + s)x + IA \\ &:= rx + sx + IA \\ &:= rx + IA \oplus sx + IA \\ &:= (rs \curvearrowright IA) \oplus (sx \curvearrowright IA). \end{aligned}$$

$$\begin{aligned}
(rs) \curvearrowright x + IA &:= rsx + IA \\
&= r(sx) + IA \\
&:= r \curvearrowright (sx + IA) \\
&= r \curvearrowright (s \curvearrowright x + IA).
\end{aligned}$$

$$1 \curvearrowright x + IA := 1x + IA = x + IA.$$

2 Problem 2

2.1 Part 1

We want to show that every simple R -module M is cyclic, i.e. if the only ideals of M are (0) and M itself, that $M = \langle m \rangle$ for some element $m \in M$.

Towards a contradiction, let M be a simple R -module and suppose M is not cyclic, so $M \neq \langle m \rangle$ for any $m \in M$. But then let $a \in M$ be an arbitrary nontrivial element; then (a) is a non-empty ideal (since it contains a), so $(a) \neq 0$. Since M is simple, we must have $(a) = M$, a contradiction.

2.2 Part 2

Let $\phi : A \rightarrow A$ be a module endomorphism on a simple module A . Then $\text{im } \phi := \phi(A)$ is a submodule of A . Since A is simple, we have either $\text{im } \phi = 0$, in which case ϕ is the zero map, or $\text{im } \phi = A$, so ϕ is surjective. In this case, we can also consider $\ker \phi$, which is a submodule of A . Since A is simple, we can again only have $\ker \phi = A$, which can not happen if ϕ is not the zero map, or $\ker \phi = 0$, in which case ϕ is both a surjective and an injective map and thus an isomorphism of modules.

3 Problem 3

3.1 Part 1

We want to show that if A, B are R -modules then $X = (\text{hom}_{R\text{-mod}}(A, B), +)$ is an abelian group. Let $f, g, h \in X$, we then need to show the following:

- a. Closure: $f + g \in X$
- b. Associativity: $f + (g + h) = (f + g) + h$
- c. Identity: $\text{id} \in X$
- d. Inverses: $f^{-1} \in X$
- e. Commutativity: $f + g = g + f$

Closure: This follows from the definition, because $(f + g) \curvearrowright x := f(x) + g(x)$ pointwise, which is well-defined homomorphism $A \rightarrow B$.

Associativity: We have

$$\begin{aligned}
f + (g + h) \curvearrowright x &:= f(x) + (g + h)(x) \\
&:= f(x) + (g(x) + h(x)) \\
&= (f(x) + g(x)) + h(x) \\
&= (f + g) + h \curvearrowright x.
\end{aligned}$$

Identity: We can define $\mathbf{0} : A \rightarrow B$ by $\mathbf{0}(x) = 0 \in B$. Then

$$(f + \mathbf{0}) \curvearrowright x = f(x) + 0 = f(x) = 0 + f(x) = (\mathbf{0} + f) \curvearrowright x.$$

Inverses: Given $f \in X$, we can define $-f : A \rightarrow B$ as $-f(x) = -x$. Then

$$\begin{aligned}
(f + -f) \curvearrowright x &= f(x) + -f(x) = f(x) - f(x) = x - x = 0 = \mathbf{0} \curvearrowright x \\
(-f + f) \curvearrowright x &= -f(x) + f(x) = -f(x) + f(x) = -x + x = 0 = \mathbf{0} \curvearrowright x.
\end{aligned}$$

Commutativity: Since B is a module, by definition $(B, +)$ is an abelian group. Thus

$$(f + g) \curvearrowright x = f(x) + g(x) = g(x) + f(x) = (g + f) \curvearrowright x.$$

3.2 Part 2

By part 1, $(\text{hom}_{R\text{-mod}}(A, A), +)$ is an abelian group, We just need to check that $(\text{hom}_R(A, A), \circ)$ is a monoid, i.e.:

- Associativity: $f \circ (g \circ h) = (f \circ g) \circ h$
- Identity: $\text{id} \circ f = f$
- Closure: $f \circ g \in \text{hom}_{R\text{-mod}}(A, A)$

Associativity: We have

$$\begin{aligned}
f \circ (g \circ h) \curvearrowright x &:= (f \circ (g \circ h))(x) \\
&= f((g \circ h)(x)) \\
&= f(g(h(x))) \\
&= (f \circ g)(h(x)) \\
&= ((f \circ g) \circ h)(x) \\
&:= (f \circ g) \circ h \curvearrowright x.
\end{aligned}$$

Identity: Take $\text{id}_A : A \rightarrow A$ given by $\text{id}_A(x) = x$, then

$$f \circ \text{id}_A \curvearrowright x = f(\text{id}_A(x)) = f(x) = \text{id}_A(f(x)) = \text{id}_A \circ f \curvearrowright x.$$

Closure: If $f : A \rightarrow A$ and $g : A \rightarrow A$ are homomorphisms, then $f \circ g : A \rightarrow A$ as a set map, and is an R -module homomorphism because

$$\begin{aligned} f \circ g \curvearrowright (r + s)(x + y) &= f(g((r + s)(x + y))) \\ &= f((r + s)(g(x) + g(y))) \\ &= (r + s)(f(g(x)) + f(g(y))) \\ &= (f \curvearrowright (r + s)(x + y)) \circ (g \curvearrowright (r + s)(x + y)). \end{aligned}$$

3.3 Part 3

For arbitrary $x, y \in A$, we need to check the following:

- a. $f \curvearrowright (x + y) = f \curvearrowright x + f \curvearrowright y$
- b. $(f + g) \curvearrowright x = f \curvearrowright x + g \curvearrowright x$
- c. $f \circ g \curvearrowright x = f \curvearrowright (g \curvearrowright x)$
- d. $\text{id}_A \curvearrowright x = x$

For (a):

$$\begin{aligned} f \curvearrowright (x + y) &:= f(x + y) \\ &= f(x) + f(y) \quad \text{since } f \text{ is a homomorphism} \\ &= f \curvearrowright x + f \curvearrowright y \end{aligned}$$

For (b):

$$\begin{aligned} (f + g) \curvearrowright x &= (f + g)(x) \\ &= f(x) + g(x) \\ &= f \curvearrowright x + g \curvearrowright x. \end{aligned}$$

For (c):

$$\begin{aligned} f \circ g \curvearrowright x &= (f \circ g)(x) \\ &= f(g(x)) \\ &= f \curvearrowright g(x) \\ &= f \curvearrowright (g \curvearrowright x). \end{aligned}$$

For (d):

$$\text{id}_A \curvearrowright x = \text{id}_A(x) = x.$$

4 Problem 4

We have the following situation:

$$\begin{array}{c} A_1 \\ \downarrow \\ Y \end{array}$$