

Notes on Lee's Manifolds

D. Zack Garza

Sunday 5th July, 2020

Contents

1	Preface: Point Set Review	1
1.1	Quotients	1
1.2	Subspaces	2
1.3	Products	2
1.4	Misc	3
2	Chapter 1: Point-Set Properties of Topological Manifolds	3
2.1	Notes	3
3	Chapter 1: Smooth Manifolds	10
4	Chapter 1 Problems	13
4.1	Recommended Problems	13
5	Chapter 2	20
6	Chapter 3	21

1 Preface: Point Set Review

1.1 Quotients

Definition 1.0.1 (Saturated).

A subset $A \subseteq X$ is *saturated* with respect to $p : X \rightarrow Y$ if whenever $p^{-1}(\{y\}) \cap A \neq \emptyset$, then $p^{-1}(\{y\}) \subseteq A$.

Equivalently, $A = p^{-1}(B)$ for some $B \subseteq Y$, i.e. it is a complete inverse image of some subset of Y , i.e. A is a union of fibers $p^{-1}(b)$.

Definition 1.0.2 (Quotient Map).

A continuous surjective map $p : X \rightarrow Y$ is a *quotient map* if $U \subseteq Y$ is open **iff** $p^{-1}(U) \subset X$ is open.

Note that \Rightarrow comes from the definition of continuity of p , but \Leftarrow is a stronger condition.

Equivalently, p maps saturated subsets of X to open subsets of Y .

Definition 1.0.3 (Universal Property of Quotients).

For $\pi : X \rightarrow Y$ a quotient map, if $g : X \rightarrow Z$ is a map that is constant on each $p^{-1}(\{y\})$, then there is a unique map f making the following diagram commute:

$$\begin{array}{ccc} X & & \\ \downarrow \pi & \searrow g & \\ Y & \xrightarrow{\quad f \quad} & Z \end{array}$$

Fact: an injective quotient map is a homeomorphism.

Fact: a product of quotient maps need not be a quotient map.

1.2 Subspaces**Definition 1.0.4** (The Subspace Topology).

$U \subset A$ is open iff $U = V \cap A$ for some open $V \subseteq X$.

Proposition 1.1 (*Universal Property of Subspaces*).

If X and $\iota_S : S \hookrightarrow Y$ is a subspace, then every continuous map $f : X \rightarrow S$ lifts to a continuous map $\tilde{f} : X \rightarrow Y$ where $\tilde{f} := \iota_S \circ f$:

$$\begin{array}{ccc} & Y & \\ \exists! \tilde{f} \nearrow & \uparrow \iota_S & \\ X & \xrightarrow{\quad f \quad} & S \end{array}$$

Note that we can view $\iota_S := \text{id}_Y|_S$. The subspace topology is the unique topology for which this property holds.

Some properties of subspace:

- The inclusion ι_S is a topological embedding.
- Restricting a continuous map to a subspace is still continuous.
- A basis for the subspace topology for $A \subset X$ can be obtained by intersecting basis elements of X with A .
- If X is Hausdorff/first/second-countable, then so is A .

1.3 Products**Definition 1.1.1** (The Product Topology).

The coarsest topology such that every projection map $p_\alpha : \prod_{\beta} X_\beta \rightarrow X_\alpha$ is continuous, i.e. for

every $U_\alpha \subseteq X_\alpha$ open, $p_\alpha^{-1}(U_\alpha) \in \prod_{\beta} X_\beta$ is open. For finite index sets, we can take the box

topology: the collection of sets of the form $\prod_{i=1}^N U_i$ with each U_i open in X_i forms a basis for the

product topology on $\prod_{i=1}^N X_i$.

Why these differ: in \mathbb{R}^∞ , the set $S = \prod(-1, 1)$ is open in the box topology but not the product topology, since $\{0\}^\infty$ is not contained in any basic open neighborhood contained in S .

Some properties of products:

- Projections π_i are continuous by definition.
- A basis for the product topology can be obtained by taking the product of bases.
- A map $f : X \rightarrow \prod Y_i$ into a product is continuous iff each component function $F_i := \pi_i \circ f : X \rightarrow Y_i$ is continuous.
 - I.e. if we have continuous maps $f_i : X \rightarrow Y_i$ then the composite map $F = [f_1, f_2, \dots]$ is continuous.
- Separate continuity does not imply joint continuity: A map $f : \prod X_i \rightarrow Y$ out of a product need not be continuous even if (defining $\iota_j : X_j \hookrightarrow \prod X_i$) the map $f \circ \iota_j : X_j \rightarrow Y$ is continuous for all arbitrary inclusions ι_j .
- Any map of the form $f_{\mathbf{a}_j} : X_j \rightarrow \prod_{i=1}^n X_i$ where $x \mapsto (a_1, \dots, a_{j-1}, x, a_{j+1}, \dots, a_n)$ is a topological embedding.
- If X_i are Hausdorff/first/second-countable, then so is $\prod_{i=1}^n X_i$.

1.4 Misc

Definition 1.1.2 (Precompact).

A subset $A \subseteq X$ is *precompact* iff its closure $\text{cl}_X(A)$ is compact in X .

Definition 1.1.3 (Locally Compact).

A space X is *locally compact* iff every $x \in X$ has a neighborhood which is contained in some compact subset of X .

2 Chapter 1: Point-Set Properties of Topological Manifolds

Pages 1- 29.

2.1 Notes

Definition 2.0.1 (Topological Manifold).

A topological space M that satisfies

1. M is Hausdorff, i.e. points can be separated by open sets
2. M is second-countable, i.e. has a countable basis
3. M is locally Euclidean, i.e. every point has a neighborhood homeomorphic to an open subset \hat{U} of \mathbb{R}^n for some fixed n .

The last property says $p \in M \implies \exists U$ with $p \in U \subseteq M$, $\hat{U} \subseteq \mathbb{R}^n$, and a homeomorphism $\varphi : U \rightarrow \hat{U}$.

Note that second countability is primarily needed for existence of partitions of unity.

Exercise Show that in the last condition, \hat{U} can equivalently be required to be an open ball or \mathbb{R}^n itself.

Theorem 2.1 (Topological Invariance of Dimension).

Two nonempty topological manifolds of different dimensions can not be homeomorphic.

Exercise Show that in a Hausdorff space, finite subsets are closed and limits of convergent sequences are unique.

Exercise Show that subspaces and finite products of Hausdorff (resp. second countable) spaces are again Hausdorff (resp. second countable).

Thus any open subset of a topological manifold with the subspace topology is again a topological manifold.

Exercise Give an example of a connected, locally Euclidean Hausdorff space that is not second countable.

Definition 2.1.1 (Charts).

A chart on M is a pair (U, φ) where $U \subseteq M$ is open and $\varphi : U \rightarrow \hat{U}$ is a homeomorphism from U to $\hat{U} = \varphi(U) \subseteq \mathbb{R}^n$. If $p \in M$ and $\varphi(p) = 0 \in \hat{U}$, then the chart is said to be *centered* at p . Note that any chart about p can be modified to a chart (φ_1, \hat{U}_1) that is centered at p by defining $\varphi_1(x) = \varphi(x) - \varphi(p)$.




Fig. 1.2 A coordinate chart

U is the *coordinate domain* and φ is the *coordinate map*.

Note that we can write φ in components as $\varphi(p) = [x^1(p), \dots, x^n(p)]$ where each x^i is a map $x^i : U \rightarrow \mathbb{R}$. The component functions x^i are the *local coordinates* on U .

Shorthand notation: $[x^i] := [x^1, \dots, x^n]$.

Example 2.1 (Graphs of Continuous Functions).

Define

$$\Gamma(f) = \left\{ (x, y) \in \mathbb{R}^n \times \mathbb{R}^k \mid x \in U, y = f(x) \in \hat{U} \right\}.$$

This is a topological manifold since we can take $\varphi : \Gamma(f) \longrightarrow U$ by restricting $\pi_1 : \mathbb{R}^n \times \mathbb{R}^k \longrightarrow \mathbb{R}^n$ to the subspace $\Gamma(f)$. Projections are continuous, restrictions of continuous functions are continuous.

This is a homeomorphism because the map $g : x \mapsto (x, f(x))$ is continuous and $g \circ \pi_1 = \text{id}_{\mathbb{R}^n}$ is continuous with $\pi_1 \circ g = \text{id}_{\Gamma(f)}$. Note that $U \cong \Gamma(f)$, and thus $(U, \varphi) = (\Gamma(f), \varphi)$ is a single *global* coordinate chart, called the *graph coordinates* of f .

Thus graphs of continuous functions $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ are locally Euclidean?

Note that this works in greater generality:: “The same observation applies to any subset of \mathbb{R}^{n+k} by setting *any* k of the coordinates equal to some continuous function of the other n .”

Coordinates as numbers vs functions?

Example 2.2 (Spheres).

S^n is a subspace of \mathbb{R}^{n+1} and is thus Hausdorff and second-countable by exercise 2.1.




Fig. 1.3 Charts for S^n

To see that it's locally Euclidean, take

$$U_i^+ := \left\{ [x^1, \dots, x^n] \in \mathbb{R}^{n+1} \mid x^i > 0 \right\} \quad \text{for } 1 \leq i \leq n+1$$

$$U_i^- := \left\{ [x^1, \dots, x^n] \in \mathbb{R}^{n+1} \mid x^i < 0 \right\} \quad \text{for } 1 \leq i \leq n+1.$$

Define

$$f : \mathbb{R}^n \longrightarrow \mathbb{R}^{\geq 0}$$

$$\mathbf{x} \mapsto \sqrt{1 - \|\mathbf{x}\|^2}.$$

Note that we immediately need to restrict the domain to $\mathbb{D}^n \subset \mathbb{R}^n$, where $\|x\|^2 \leq 1 \implies 1 - \|x\|^2 \geq 0$, to have a well-defined real function $f : \mathbb{D}^n \longrightarrow \mathbb{R}^{\geq 0}$.

Then (claim)

$$\begin{aligned} U_i^+ \cap S^n & \text{ is the graph of } x^i = f(x^1, \dots, \widehat{x^i}, \dots, x^{n+1}) \\ U_i^- \cap S^n & \text{ is the graph of } x^i = -f(x^1, \dots, \widehat{x^i}, \dots, x^{n+1}). \end{aligned}$$

This is because

$$\begin{aligned} \Gamma(x^i) &:= \{(\mathbf{x}, f(\mathbf{x})) \subseteq \mathbb{R}^n \times \mathbb{R}\} \\ &= \left\{ [x_1, \dots, \widehat{x^i}, \dots, x^{n+1}], f([x_1, \dots, \widehat{x^i}, \dots, x^{n+1}]) \subseteq \mathbb{R}^n \times \mathbb{R} \right\} \\ &= \left\{ [x_1, \dots, \widehat{x^i}, \dots, x^{n+1}], \left(1 - \sum_{\substack{j=1 \\ j \neq i}}^{n+1} (x^j)^2 \right)^{\frac{1}{2}} \subseteq \mathbb{R}^n \times \mathbb{R} \right\} \end{aligned}$$

and any vector in this set has norm satisfying

$$\|(\mathbf{x}, y)\|^2 = \sum_{\substack{j=1 \\ j \neq i}}^{n+1} (x^j)^2 + \left(1 - \sum_{\substack{j=1 \\ j \neq i}}^{n+1} (x^j)^2 \right) = 1$$

and is thus in S^n .

To see that any such point also has positive i coordinate and is thus in U_i^+ , we can rearrange (?) coordinates to put the value of f in the i th coordinate to obtain

$$\Gamma(x_i) = \left\{ [x^1, \dots, f(x^1, \dots, \widehat{x^i}, \dots, x^n), \dots, x^n] \right\}$$

Seems like f is always the *last* coordinate in the graph

and note that the square root only takes on positive values.

Thus each $U_i^\pm \cap S^n$ is the graph of a continuous function and thus locally Euclidean, and we can define chart maps

$$\begin{aligned} \varphi_i^\pm : U_i^\pm \cap S^n &\longrightarrow \mathbb{D}^n \\ [x^1, \dots, x^n] &\mapsto [x^1, \dots, \widehat{x^i}, \dots, x^{n+1}] \end{aligned}$$

yield $2(n+1)$ charts that are graph coordinates for S^n .

Example 2.3 (Projective Space).

Define \mathbb{RP}^n as the space of 1-dimensional subspaces of \mathbb{R}^{n+1} with the quotient topology determined by the map

$$\begin{aligned} \pi : \mathbb{R}^{n+1} \setminus \{0\} &\longrightarrow \mathbb{RP}^n \\ \mathbf{x} &\mapsto \text{span}_{\mathbb{R}} \{\mathbf{x}\}. \end{aligned}$$

How is this map a quotient map?

Notation: for $\mathbf{x} \in \mathbb{R}^{n+1} \setminus \{0\}$ write $[\mathbf{x}] := \pi(\mathbf{x})$, the line spanned by \mathbf{x} .

Define charts:

$$\tilde{U}_i := \left\{ \mathbf{x} \in \mathbb{R}^{n+1} \setminus \{0\} \mid x^i \neq 0 \right\}, \quad U_i = \pi(\tilde{U}_i) \subseteq \mathbb{RP}^n$$

and chart maps

$$\begin{aligned} \tilde{\varphi}_i : \tilde{U}_i &\longrightarrow \mathbb{R}^n \\ [x^1, \dots, x^{n+1}] &\mapsto \left[\frac{x^1}{x^i}, \dots, \widehat{x^i}, \dots, \frac{x^{n+1}}{x^i} \right]. \end{aligned}$$

Then (claim) this descends to a continuous map $\varphi_i : U_i \longrightarrow \mathbb{R}^n$ by the universal property of the quotient:

$$\begin{array}{ccc} \tilde{U}_i & & \\ \pi_U \downarrow & \searrow \tilde{\varphi}_i & \\ U_i & \xrightarrow{\varphi_i} & \mathbb{R}^n \end{array}$$

- The restriction $\pi_U : \tilde{U}_i \longrightarrow U_i$ of π is still a quotient map because $\tilde{U}_i = \pi_U^{-1}(U_i)$ where $U_i \subseteq \mathbb{RP}^n$ is open in the quotient topology and thus \tilde{U}_i is saturated.

Thus π_U sends saturated sets to open sets and is thus a quotient map.

- $\tilde{\varphi}_i$ is constant on preimages under π_U : fix $y \in U_i$, then $\pi_U^{-1}(\{y\}) = \{\lambda \mathbf{y} \mid \lambda \in \mathbb{R} \setminus \{0\}\}$, i.e. the point $y \in \mathbb{RP}^n$ pulls back to every nonzero point on the line spanned by $\mathbf{y} \in \mathbb{R}^n$.

But

$$\begin{aligned} \tilde{\varphi}_i(\lambda \mathbf{y}) &= \varphi_i([\lambda y^1, \dots, \lambda y^i, \dots, \lambda y^n]) \\ &= \left[\frac{\lambda y^1}{\lambda y^i}, \dots, \widehat{\lambda y^i}, \dots, \frac{\lambda y^{n+1}}{\lambda y^i} \right] \\ &= \left[\frac{y^1}{y^i}, \dots, \widehat{y^i}, \dots, \frac{y^{n+1}}{y^i} \right] \\ &= \tilde{\varphi}_i(\mathbf{y}). \end{aligned}$$

So this yields a continuous map

$$\varphi_i : U_i \longrightarrow \mathbb{R}^n.$$

We can now verify that φ is a homeomorphism since it has a continuous inverse given by

$$\begin{aligned} \varphi_i^{-1} : \mathbb{R}^n &\longrightarrow U_i \subseteq \mathbb{RP}^n \\ \mathbf{u} := [u^1, \dots, u^n] &\mapsto [u^1, \dots, u^{i-1}, \mathbf{1}, u^{i+1}, \dots, u^n]. \end{aligned}$$

It remains to check:

Exercise

1. The $n + 1$ sets U_1, \dots, U_{n+1} cover \mathbb{RP}^n .
2. \mathbb{RP}^n is Hausdorff
3. \mathbb{RP}^n is second-countable.

Exercise (1.6) Show that \mathbb{RP}^n is Hausdorff and second countable.

Exercise (1.7) Show that \mathbb{RP}^n is compact. (Hint: show that π restricted to S^n is surjective.)

Definition 2.1.2 (Topological Embedding).

A continuous map $f : X \rightarrow Y$ is a *topological embedding* iff it is injective and $\tilde{f} : X \rightarrow f(X)$ is a homeomorphism.

Example 2.4 (Product Manifolds).

Let $M := M_1 \times \dots \times M_k$ be a product of manifolds of dimensions n_1, \dots, n_k respectively. A product of Hausdorff/second-countable spaces is still Hausdorff/second-countable, so just need to check that it's locally Euclidean.

- Let $\mathbf{p} \in \prod_{i=1}^N M_i$, so $p_i \in M_i$
- Choose a chart (U_i, φ_i) with $p_i \in U_i$ and assemble a product map:

$$\Phi := \prod \varphi_i : \prod U_i \rightarrow \prod \mathbb{R}^{n_i} \cong \mathbb{R}^{\sum n_i} := \mathbb{R}^N.$$

- Claim: Φ is a homeomorphism onto its image in \mathbb{R}^N .
 - Each φ_i is a homeomorphism onto $\varphi_i(U_i)$ (by the definition of a chart on M_i)
 - It suffices to show that Φ^{-1} exists and is continuous, where

$$\Phi^{-1}(V) := \left(\prod \varphi_i \right)^{-1} \left(\prod V_i \right).$$

- Φ is a product of continuous functions and thus continuous.
- $\Phi^{-1} := \left(\prod \varphi_i \right)^{-1} = \prod \varphi_i^{-1}$, which are all assumed continuous since φ_i were homeomorphisms.

Example 2.5 (Tori).

$T^n := \prod_{i=1}^n S^1$ is a topological n -manifold.

Definition 2.1.3 (Precompact).

A subset $A \subseteq X$ is *precompact* iff its closure $\text{cl}_X(A)$ is compact in X .

Proposition 2.2.

Every topological manifold has a countable basis of precompact coordinate balls.

Proposition 2.3.

Let M be a topological manifold.

- M is locally path-connected.
- M is connected $\iff M$ is path-connected
- The connected components and path components of M coincide.
- $\pi_0(M)$ is countable and each component is open and a connected topological manifold.

Proposition 2.4.

Every topological manifold M is locally compact.

Proof .

M has a basis of precompact open sets. ■

Theorem 2.5 (Manifolds are Paracompact).

Given any open cover $\mathcal{U} \rightrightarrows M$ of a topological manifold and any basis \mathcal{B} for the topology on M , there exists a countable locally finite open refinement of \mathcal{U} consisting of elements of \mathcal{B} .

Proposition 2.6.

$\pi_1(M)$ is countable.

3 Chapter 1: Smooth Manifolds

Definition 3.0.1 (Smooth Functions).

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $[f_1(\mathbf{x}^n), f_2(\mathbf{x}^n), \dots, f_m(\mathbf{x}^n)]$ (or any subsets thereof) is said to be C^∞ or **smooth** iff each f_i has continuous partial derivatives of all orders.


Definition 3.0.2 (Diffeomorphism).

A smooth bijective map with a smooth inverse is a *diffeomorphism*.

Remark A diffeomorphism is necessarily a homeomorphism, but not conversely.

Definition 3.0.3 (Transition Maps).

If $(U, \varphi), (V, \psi)$ are two charts on M such that $U \cap V \neq \emptyset$, the composite map $\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V)$ is a function $\mathbb{R}^n \rightarrow \mathbb{R}^n$ and is called the *transition map* from φ to ψ .



Two charts are *smoothly compatible* iff $U \cap V = \emptyset$ or $\psi \circ \varphi^{-1}$ is a diffeomorphism.

Definition 3.0.4.

A collection of charts $\mathcal{A} := \{(U_\alpha, \varphi_\alpha)\}$ is an *atlas* for M iff $\{U_\alpha\} \rightrightarrows M$, and is a *smooth atlas* iff all of the charts it contains are pairwise smoothly compatible.

Remark To show an atlas is smooth, it suffices to show that an arbitrary $\psi \circ \varphi^{-1}$ is smooth.

This is because this immediately implies that its inverse is smooth, and these are diffeomorphisms. Alternatively, one can show that $\psi \circ \varphi^{-1}$ is smooth, injective, and has nonsingular Jacobian at each point.

Remark Attempting to define a function $f : M \rightarrow \mathbb{R}$ to be smooth iff $f \circ \varphi^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}$ is smooth for each φ may not work because many atlases give the “same” smooth structure in the sense that they all determine the same collection of smooth functions on M .

For example, take the following two atlases on \mathbb{R}^n :

$$\begin{aligned} \mathcal{A}_1 &= \{(\mathbb{R}^n, \text{Id}_{\mathbb{R}^n})\} \\ \mathcal{A}_2 &= \left\{ \left(\mathbb{D}_1(\mathbf{x}), \text{id}_{\mathbb{D}_1(\mathbf{x})} \right) \mid \mathbf{x} \in \mathbb{R}^n \right\} . \end{aligned}$$

Claim: a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is smooth wrt either atlas iff it is smooth in the usual sense.

What does “determine the same collection of smooth functions” mean?

Definition 3.0.5 (Maximal or Complete Atlas).

A smooth atlas on M is *maximal* iff it is not properly contained in any larger smooth atlas.

Remark Not every topological manifold admits a smooth structure. See Kervaire’s 10-dimensional manifold from 1960.

Definition 3.0.6 (Smooth Structures and Smooth Manifolds).

If M is a topological manifold, a maximal smooth atlas \mathcal{A} is a *smooth structure* on M . The triple (M, τ, \mathcal{A}) where \mathcal{A} is a smooth structure is a *smooth manifold*.

Remark To show that two smooth structures are *distinct*, it suffices to show that they are not smoothly compatible, i.e. one of the transition functions $\psi \circ \varphi^{-1}$ is not smooth. This is because any maximal atlas \mathcal{A}_1 must contain ψ and likewise \mathcal{A}_2 contains φ^{-1} , but no maximal atlas can contain φ and ψ because all charts in a maximal atlas are smoothly compatible by definition.

Proposition 3.1.

Let M be a topological manifold.

1. Every smooth atlas \mathcal{A} for M is contained in a unique maximal smooth atlas, called the *smooth structure determined by \mathcal{A}* .
2. Two smooth atlases for M determine the same smooth structure \iff their union is a smooth atlas.

Remark That we can place many requirements on the functions $\psi \circ \varphi^{-1}$ and get various other structures: C^k , real-analytic, complex-analytic, etc. C^0 structures recover topological manifolds.

Definition 3.1.1 (Smooth Charts, Maps, Domains).

If (M, τ, \mathcal{A}) is a smooth manifold, any chart $(U, \varphi) \in \mathcal{A}$ is a *smooth chart*, where U is a *smooth coordinate domain* and φ is a *smooth coordinate map*. A *smooth coordinate ball* is a smooth coordinate domain U such that $\varphi(U) = \mathbb{D}^n$.

Definition 3.1.2 (Regular Coordinate Ball).

A set $B \subseteq M$ is a *regular coordinate ball* if there is a smooth coordinate ball B' such that $\text{cl}_M(B) \subseteq B'$, and a smooth coordinate map $\varphi : B' \rightarrow \mathbb{R}^n$ such that for some positive numbers $r < r'$,

- $\varphi(B) = \mathbb{D}_r(\mathbf{0})$,
- $\varphi(B') = \mathbb{D}_{r'}(\mathbf{0})$, and
- $\varphi(\text{cl}_M(B)) = \text{cl}_{\mathbb{R}^n}(\mathbb{D}_r(\mathbf{0}))$.

This says B “sits nicely” inside a larger coordinate ball.

Remark $\text{cl}_M(B) \cong_{\text{Top}} \text{cl}_{\mathbb{R}^n}(\mathbb{D}_r(\mathbf{0}))$ which is closed and bounded and thus compact, so $\text{cl}_M(B)$ is compact. Thus every regular coordinate ball in M is precompact.

Proposition 3.2.

Every smooth manifold has a countable basis of regular coordinate balls.

Remark There is only one 0-dimensional smooth manifold, up to equivalence of smooth structures.

Definition 3.2.1 (Standard Smooth Structure on \mathbb{R}^n).

Define the atlas $\mathcal{A}_0 = \{(\mathbb{R}^n, \text{id}_{\mathbb{R}^n})\}$ and take the smooth structure it generates, this is the *standard smooth structure* on \mathbb{R}^n .

Proposition 3.3.

There are at least two distinct smooth structures on \mathbb{R}^n .

Proof.

Define $\psi(x) = x^3$; then $\mathcal{A}_1 := \{(\mathbb{R}^n, \varphi)\}$ defines a smooth structure.

Then $\mathcal{A}_1 \neq \mathcal{A}_0$, which follows because $(\text{id}_{\mathbb{R}^n} \circ \varphi^{-1})(x) = x^{\frac{1}{3}}$, which is not smooth at $\mathbf{0}$. ■

4 Chapter 1 Problems

4.1 Recommended Problems

Note: helpful theorem, two smooth structures induced by two smooth atlases $\mathcal{A}_1, \mathcal{A}_2$ are equivalent iff $\mathcal{A}_1 \cup \mathcal{A}_2$ is again a smooth atlas. So it suffices to check pairwise compatibility of charts.

Exercise (Problem 1.6) Show that if $M^n \neq \emptyset$ is a topological manifold of dimension $n \geq 1$ and M has a smooth structure, then it has uncountably many distinct ones.

Hint: show that for any $s > 0$ that $F_s(x) := |x|^{s-1}x$ defines a homeomorphism $F_x : \mathbb{D}^n \rightarrow \mathbb{D}^n$ which is a diffeomorphism iff $s = 1$.

Recommended problem

Solution:

Define

$$F_s : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$\mathbf{x} \mapsto \|\mathbf{x}\|^{s-1} \mathbf{x}.$$

Claim: F_s restricted to \mathbb{D}^n is a continuous map $\mathbb{D}^n \rightarrow \mathbb{D}^n$.

- Note that if $\|\mathbf{x}\| \leq \varepsilon < 1$ then

$$\|F_s(\mathbf{x})\| = \| \|\mathbf{x}\|^{s-1} \mathbf{x} \| = \|\mathbf{x}\|^s \leq \|\mathbf{x}\| \leq \varepsilon < 1,$$

so $F_s(\mathbb{D}^n) \subseteq \mathbb{D}^n$ and moreover $F_s(\mathbb{D}_\varepsilon^n) \subseteq \mathbb{D}_\varepsilon^n$.

- We'll use the fact that $F_s^{-1} = F_{\frac{1}{s}}$ is of the same form, and thus $F_s^{-1}(\mathbb{D}^n) \subseteq \mathbb{D}^n$, forcing $F_s(\mathbb{D}^n) = \mathbb{D}^n$.

- This is a continuous function on the punctured disc $\mathbb{D}_0^n := \mathbb{D}^n \setminus \{\mathbf{0}\}$, since it can be written as a composition of smooth functions:

$$\begin{aligned} \mathbb{D}_0^n &\xrightarrow{\Delta} \mathbb{D}_0^n \times \mathbb{D}_0^n \xrightarrow{(\|\cdot\|, \text{id}_{\mathbb{D}_0^n})} \mathbb{D}_0^n \times \mathbb{D}_0^n \xrightarrow{((\cdot)^{s-1}, \text{id}_{\mathbb{D}_0^n})} \mathbb{D}_0^1 \times \mathbb{D}_0^n \xrightarrow{(a,b) \mapsto ab} \mathbb{D}_0^n \\ \mathbf{x} &\longrightarrow (\mathbf{x}, \mathbf{x}) \longrightarrow (\|\mathbf{x}\|, \mathbf{x}) \longrightarrow (\|\mathbf{x}\|^{s-1}, \mathbf{x}) \longrightarrow \|\mathbf{x}\|^{s-1} \mathbf{x} \end{aligned}$$

For any $s \geq 0$, continuity at zero follows from the fact that $\|F_s(\mathbf{x})\| \leq \|\mathbf{x}\| \rightarrow 0$, so $\lim_{\mathbf{x} \rightarrow \mathbf{0}} F_s(\mathbf{x}) = \mathbf{0}$ and the sequential definition of continuity applies. So F_s is continuous on \mathbb{D}^n for every s .

Here we are taking for granted the fact that taking norms, exponentiating, and multiplying are all smooth functions away from zero.

Claim: F_s is a bijection $\mathbb{D}^n \setminus \mathbf{0} \hookrightarrow \mathbb{D}^n \setminus \mathbf{0}$ that extends to a bijection $\mathbb{D}^n \hookrightarrow \mathbb{D}^n$.

We can note that

$$F_s(\mathbf{x}) = \begin{cases} \|\mathbf{x}\|^s \frac{\mathbf{x}}{\|\mathbf{x}\|} := \|\mathbf{x}\|^s \widehat{\mathbf{x}} & \text{if } \|\mathbf{x}\| \neq 0 \\ \mathbf{0} & \text{if } \|\mathbf{x}\| = 0 \end{cases}$$

This follows because we can construct a two-sided inverse that composes to the identity, namely $F_{\frac{1}{s}}$, for $\mathbf{x} \neq \mathbf{0}$, and note that $F_s(\mathbf{0}) = \mathbf{0}$. Using the fact that $\|t\mathbf{x}\| = t\|\mathbf{x}\|$ for any scalar t , we can check that

$$\begin{aligned} (F_s \circ F_{\frac{1}{s}})(\mathbf{x}) &= F_s(\|\mathbf{x}\|^{\frac{1}{s}} \widehat{\mathbf{x}}) \\ &= \left\| \|\mathbf{x}\|^{\frac{1}{s}} \widehat{\mathbf{x}} \right\|^s \cdot \widehat{\|\mathbf{x}\|^{\frac{1}{s}} \widehat{\mathbf{x}}} \\ &= \left(\|\mathbf{x}\|^{\frac{1}{s}} \right)^s \cdot \|\widehat{\mathbf{x}}\|^s \cdot \frac{\|\mathbf{x}\|^{\frac{1}{s}} \widehat{\mathbf{x}}}{\left\| \|\mathbf{x}\|^{\frac{1}{s}} \widehat{\mathbf{x}} \right\|} \\ &= \|\mathbf{x}\| \cdot 1^s \cdot \left(\frac{\|\mathbf{x}\|^{\frac{1}{s}}}{\|\mathbf{x}\|^{\frac{1}{s}}} \right) \cdot \frac{\widehat{\mathbf{x}}}{\|\widehat{\mathbf{x}}\|} \\ &= \|\mathbf{x}\| \widehat{\mathbf{x}} \\ &= \mathbf{x}. \end{aligned}$$

and similarly

$$\begin{aligned} (F_{\frac{1}{s}} \circ F_s)(\mathbf{x}) &= F_{\frac{1}{s}}(\|\mathbf{x}\|^s \widehat{\mathbf{x}}) \\ &= \left\| \|\mathbf{x}\|^s \widehat{\mathbf{x}} \right\|^{\frac{1}{s}} \cdot \widehat{\|\mathbf{x}\|^s \widehat{\mathbf{x}}} \\ &= \left(\|\mathbf{x}\|^s \right)^{\frac{1}{s}} \|\widehat{\mathbf{x}}\|^{\frac{1}{s}} \cdot \frac{\|\mathbf{x}\|^s \widehat{\mathbf{x}}}{\left\| \|\mathbf{x}\|^s \widehat{\mathbf{x}} \right\|} \\ &= \|\mathbf{x}\| \cdot 1^{1-s} \cdot \left(\frac{\|\mathbf{x}\|^s}{\|\mathbf{x}\|^s} \right) \cdot \frac{\widehat{\mathbf{x}}}{\|\widehat{\mathbf{x}}\|} \\ &= \|\mathbf{x}\| \widehat{\mathbf{x}} \\ &= \mathbf{x}. \end{aligned}$$

Claim: F_s is a homeomorphism for all s .

This follows from the fact that the domain \mathbb{D}^n is compact and the codomain \mathbb{D}^n is Hausdorff, and a continuous bijection between such spaces is a homeomorphism.

Claim: F_s is a diffeomorphism iff $s = 1$.

If $s = 1$, $F_s = \text{id}_{\mathbb{D}^n}$ which is clearly a diffeomorphism.

Otherwise, we claim that F_s is not a diffeomorphism because either F_s or F_s^{-1} will fail to be smooth at $\mathbf{x} = \mathbf{0}$.

- If $0 \leq s < 1$, then F_s fails to be differentiable at zero.
- If $1 < s < \infty$ then $0 \leq \frac{1}{s} < 1$ and the same argument applies to $F_s^{-1} := F_{\frac{1}{s}}$.

Why? Should boil down to $x \mapsto x^t$ for $0 \leq t < 1$ failing to be differentiable at 0 in \mathbb{R}

We now show that we can produce infinitely many distinct maximal atlases on M . Let \mathcal{A} be any smooth atlas on M and fix $p_0 \in M$.

Claim: We can modify \mathcal{A} to obtain an atlas \mathcal{A}' where p_0 is in exactly one chart (V, ψ) with $\psi(p_0) = \mathbf{0} \in \mathbb{R}^n$.

- Pick a chart containing p_0 , say (U, φ) where $\varphi(p_0) := \mathbf{p}$
- Since $\varphi(U) \subseteq \mathbb{R}^n$ is open, find a disc containing \mathbf{p} , say $\mathbb{D}_R(\mathbf{p}) \subset \varphi(U)$.
- Define $V \subseteq M$ as $V := \varphi^{-1}(\mathbb{D}_R(\mathbf{p}))$.
- Define $\psi : U \rightarrow \mathbb{R}^n$ by

$$\begin{aligned} \psi : U &\longrightarrow \mathbb{R}^n \\ x &\mapsto \frac{\varphi(x) - \varphi(p_0)}{R}. \end{aligned}$$

- Note: this is constructed precisely so that $\psi(V) = \mathbb{D}_1(\mathbf{0}) \in \mathbb{R}^n$ and $\psi(p) = 0$.
- This is a homeomorphism onto its image since we can write

$$\psi = \delta_{\frac{1}{R}} \circ \tau_{\mathbf{p}} \circ \varphi$$

is a composition of continuous functions, where δ, τ are dilations/translations in \mathbb{R}^n which are known to be continuous, and

$$\psi^{-1} = \varphi^{-1} \circ \tau_{-\mathbf{p}} \circ \delta_R$$

is again a composition of smooth (and in particular, continuous) functions.

- Define $\mathcal{A}^1 := \mathcal{A} \cup \{(V, \psi|_V)\}$
 - This is a smooth atlas: any pair of charts coming from \mathcal{A} are smoothly compatible, so it suffices to check that an arbitrary chart from \mathcal{A} is smoothly compatible with the new chart.
 - Let (T, ξ) be any other chart, then if $T \cap V \neq \emptyset$, the transition function

$$\psi \circ \xi^{-1} = \delta_{\frac{1}{R}} \tau_{\mathbf{p}} \circ \varphi \circ \xi^{-1}$$

is a composition of smooth functions and thus smooth, and similarly for $\xi \circ \psi^{-1}$.

- Since the charts from \mathcal{A} cover M , so do the charts of \mathcal{A}^1 since $\mathcal{A} \subseteq \mathcal{A}^1$.
- For every $(U_\alpha, \varphi_\alpha) \in \mathcal{A}^1$, define a new chart $(U_\alpha \setminus \{p\}, \varphi_\alpha|_{U_\alpha \setminus \{p\}})$ and define this set of charts as \mathcal{A}^2 .
 - This still covers M : p is in the chart $(V, \psi|_V)$, and if $q \neq p$, then $q \in U_\alpha$ for some α since \mathcal{A} was an atlas, and $q \in U_\alpha \setminus \{p\}$.
 - The coordinate maps are still homeomorphisms onto their images, because the restriction of a homeomorphism is again a homeomorphism.

- The transition functions are still smooth because the restriction of a smooth function is again smooth.

Claim: We can define a new atlas \mathcal{A}_s from \mathcal{A}^2 by only replacing the single chart (V, ψ) with $(V, F_s \circ \psi)$.

- \mathcal{A}_s still covers M , since we haven't changed the coordinate domains
- All coordinate functions are still a homeomorphisms onto their images, since the only change is ψ is replaced with $F_s \circ \psi$ and we've shown that F_s is a homeomorphism; a composition of homeomorphisms is again a homeomorphism.
- The chart $(V, F_s \circ \psi)$ is still a valid chart, since $F_s : \mathbb{D}_n \hookrightarrow \mathbb{D}_n$ and $\psi(V) \cong \mathbb{D}^n$ by construction.
- All charts in \mathcal{A}_s are still smoothly compatible:
 - It suffices to check compatibility between an arbitrary $(U_\alpha, \varphi_\alpha)$ and $(V, F_s \circ \psi)$, so we consider $F_s \circ \psi \circ \varphi_\alpha^{-1}$
 - By construction, $p \notin U_\alpha$, and we know F_s is smooth away from $\mathbf{0}$, so this is a smooth function.

Claim: If $s \neq t$ then \mathcal{A}_s and \mathcal{A}_t are not smoothly compatible, and thus generate distinct maximal smooth atlases.

- If $\mathcal{A}_s, \mathcal{A}_t$ define the same smooth structure, then in particular $(V, F_s \circ \psi)$ must be smoothly compatible with $(V, F_t \circ \psi)$.
- We can compute the transition function

$$(F_s \circ \psi) \circ (F_t \circ \psi)^{-1} = F_s \circ \psi \circ \psi^{-1} \circ F_t^{-1} = F_s \circ F_t^{-1} = F_s \circ F_t^{-1} = F_s \circ F_t^{-1} = F_{\frac{s}{t}}.$$

- From above, we know this is smooth iff $\frac{s}{t} = 1$, i.e. $s = t$.
- So if $s \neq t$, then the maximal atlases correspond to $\mathcal{A}_s, \mathcal{A}_t$ each contain a chart that is not smoothly compatible with the other, and so these are distinct smooth structures.

■

Exercise (Problem 1.7) Let $N := [0, \dots, 1] \in S^n$ and $S := [0, \dots, -1]$ and define the stereographic projection

$$\sigma : S^n \setminus N \longrightarrow \mathbb{R}^n$$

$$[x^1, \dots, x^{n+1}] \mapsto \frac{1}{1 - x^{n+1}} [x^1, \dots, x^n]$$

and set $\tilde{\sigma}(x) = -\sigma(-x)$ for $x \in S^n \setminus S$ (projection from the South pole)

Recommended
problem

Note that the figure should say $\{x^{n+1} = 0\}$ instead of x^n .




Fig. 1.13 Stereographic projection

1. For any $x \in S^n \setminus N$ show that $\sigma(x) = \mathbf{u}$ where $(\mathbf{u}, 0)$ is the point where the line through N and x intersects the linear subspace $H_{n+1} := \{x^{n+1} = 0\}$.

Similarly show that $\tilde{\sigma}(x)$ is the point where the line through S and x intersects H_{n+1} .

2. Show that σ is bijective and

$$\sigma^{-1}(\mathbf{u}) = \sigma^{-1}\left([u^1, \dots, u^n]\right) = \frac{1}{\|\mathbf{u}\|^2 + 1} [2u^1, \dots, 2u^n, \|\mathbf{u}\|^2 - 1].$$

3. Compute the transition map $\tilde{\sigma} \circ \sigma^{-1}$ and verify that the atlas

$$\mathcal{A} := \{(S^n \setminus N, \sigma), (S^n \setminus S, \tilde{\sigma})\}$$

define a smooth structure on S^n .

4. Show that this smooth structure is equivalent to the standard smooth structure: Put graph coordinates on S^n as outlined in 2.2 to obtain $\{(U_i^\pm, \varphi_i^\pm)\}$.

For indices $i < j$, show that

$$\varphi_i^\pm \circ (\varphi_j^\pm)^{-1} [u^1, \dots, u^n] = [u^1, \dots, \hat{u}^i, \dots, \pm \sqrt{1 - \|\mathbf{u}\|^2}, \dots, u^n]$$

where the square root appears in the j th position. Find a similar formula for $i > j$. Show that if $i = j$, then

$$\varphi_i^\pm \circ (\varphi_j^\pm)^{-1} = \varphi_i^- \circ (\varphi_i^+)^{-1} = \text{id}_{\mathbb{D}^n}.$$

Show that these yield a smooth atlas.

Solution (1):

- Parameterize the line through $\mathbf{x} \in S^n$ and \mathbf{N} :

$$\begin{aligned}\ell_{N,\mathbf{x}}(t) &= t\mathbf{x} + (1-t)\mathbf{N} \\ &= t[x^1, \dots, x^n, x^{n+1}] + (1-t)[0, \dots, 1] \\ &= [tx^1, \dots, x^n, tx^{n+1} + (1-t)] \\ &= [tx^1, \dots, x^n, 1 - t(1 - x^{n+1})]\end{aligned}$$

- Evaluate at $t = \frac{1}{1 - x^{n+1}}$ to obtain $\frac{1}{x^{n+1}}[x^1, \dots, x^n, 0] = [\sigma(\mathbf{x}), 0]$.
- For $\tilde{\sigma}(\mathbf{x})$: Todo.

Todo

Solution (2):

- How to derive this formula: no clue.
 - Start with $\mathbf{u} \in \mathbb{R}^n$, parameterize the line $\ell_{N,\mathbf{u}}(t)$, solve for where $\|\ell_{N,\mathbf{u}}(t)\| = 1$ and $\mathbf{u} \neq \mathbf{N}$
 - Should yield $t^2\|\mathbf{u}\| + (1-t)^2 = 1$, solve for nonzero t ; should get $t = \frac{2}{\|\mathbf{u}\| + 1}$, so $x^i = 2u^i/(\|\mathbf{u}\| + 1)$ and $x^{n+1} = \left(\frac{2}{\|\mathbf{u}\| + 1}\right) - 1$.
- Compute compositions $\sigma \circ \sigma^{-1}$: Todo.

Figure out how to invert.

Messy computations that didn't work out.

Solution (3):

- Computing the transition maps:

$$\begin{aligned}(\tilde{\sigma} \circ \sigma^{-1})(\mathbf{u}) &= -\sigma\left(\left(\frac{-1}{\|\mathbf{u}\|^2 + 1}\right)[2u^1, \dots, 2u^n, \|\mathbf{u}\|^2 - 1]\right) \\ &= -1 \cdot \left[\frac{\frac{-2u^1}{\|\mathbf{u}\|^2 + 1}}{1 - \frac{1 - \|\mathbf{u}\|^2}{1 + \|\mathbf{u}\|^2}}, \dots, n\right] \\ &= \left[\frac{2u^1}{\|\mathbf{u}\|^2 + 1} \cdot \frac{1 + \|\mathbf{u}\|^2}{1 + \|\mathbf{u}\|^2 - (1 - \|\mathbf{u}\|^2)}, \dots, n\right] \\ &= \left[\frac{2u^1}{2\|\mathbf{u}\|^2}, \dots, n\right] \\ &= \frac{\mathbf{u}}{\|\mathbf{u}\|^2} \\ &:= \hat{\mathbf{u}},\end{aligned}$$

which is a smooth function on $\mathbb{R}^n \setminus \{\mathbf{0}\}$.

- Todo: computing $(\sigma \circ \tilde{\sigma}^{-1})(\mathbf{u}) = \hat{\mathbf{u}}$
- Todo: argue that it suffices that these are smooth on $\mathbb{R}^n \setminus \{\mathbf{0}\}$

Computation.

What are the actual domains and ranges of the transition functions? It seems like you pull back \mathbb{R}^n to $S^n \setminus N$, then push $S^n \setminus \{N, S\}$ to $\mathbb{R}^n \setminus 0$, but this yields $\mathbb{R}^n \rightarrow \mathbb{R}^n \setminus 0$ where we haven't deleted zero in the domain (problem: not smooth!).

Solution (4):

We want to argue that these define the same maximal smooth atlas, for which it suffices to the charts from each are pairwise smoothly compatible.

- Define $\varphi_i([x^1, \dots, x^n]) = [x^1, \dots, \hat{x}^i, \dots, x^n]$ and $\varphi_i^{-1}([x^1, \dots, x^{n-1}]) = [x^1, \dots, \sqrt{1 - \|x\|^2}, \dots, x^n]$.
- Compute $(\varphi_i \circ \sigma^{-1})(\mathbf{u}) = \frac{1}{\|\mathbf{u}\| + 1} [2u^1, \dots, \hat{u}^i, \dots, 2u^n, \|\mathbf{u}\|^2 - 1]$, which is (clearly) smooth?
- Compute $(\sigma \circ \varphi_i^{-1})(\mathbf{u}) = \sigma([u^1, \dots, \sqrt{1 - \|\mathbf{u}\|^2}, \dots, u^n])$, which is $\frac{1}{1 - u^n} [u^1, \dots, \sqrt{1 - \|\mathbf{u}\|^2}, \dots, u^{n-1}]$.
 - This is smooth if $u^n \neq 1$, but this corresponds to \mathbf{N} in S^2 , in which case $\varphi_i^{-1}(\mathbf{u})$ isn't in the domain of σ to begin with.

Exercise (Problem 1.8) Define an *angle function* on $U \subset S^1$ as any continuous function $\theta : U \rightarrow \mathbb{R}$ such that $e^{i\theta(z)} = z$ for all $z \in U$.

Show that U admits an angle function iff $U \neq S^1$, and for any such function θ , (U, θ) is a smooth coordinate chart for S^1 with its standard smooth structure.

Note that $f : \mathbb{R} \rightarrow S^1$ given by $f(x) = e^{ix}$ is a covering map (in fact the universal cover).

Some way to do this just with covering spaces?

$\Rightarrow :$

- Suppose there exists an angle function $\theta : U \rightarrow \mathbb{R}$.
- Then $f \circ \theta|_U = \text{id}_U$ by assumption, since $u \xrightarrow{\theta|_U} \theta(u) \xrightarrow{f} e^{i\theta(u)} = u$.
- So θ has a left-inverse and is thus injective.
- Suppose $U = S^1$, which is compact.
- Then θ is an injective continuous map on a compact set, so its image $\theta(S^1) \subseteq \mathbb{R}$ is compact.
- Lemma: a continuous map from a compact space to a Hausdorff space is a closed map.
- Since θ is injective and is surjective onto its image, since it is continuous it is a homeomorphism onto its image and $S^1 \cong \theta(S^1)$.
- Since S^1 is connected, $\theta(S^1)$ is connected, and the only connected subsets of \mathbb{R} are intervals.
- Since $\theta(S^1)$ is compact, it must be a closed and bounded subset, so $\theta(S^1) = [a, b] \subset \mathbb{R}$.
- But this forces $S^1 \cong [a, b]$ is a homeomorphism, which is a contradiction: removing one point from S^1 yields one connected component, while removing $\frac{1}{2}(b - a)$ from $[a, b]$ produces a disconnected set.

$\Leftarrow :$

- Suppose $U \neq S^1$, then there exists a point $p \in S^1 \setminus U$; wlog suppose $p = 1$.
- Then $U \subseteq S^1 \setminus \{1\}$
- Note that $f^{-1}(\{1\}) = \{2k\pi \mid k \in \mathbb{Z}\}$.
- Take the interval $I = [0, 2\pi]$ and set $\tilde{f} = f|_I$.
- Since $U \neq S^1$, $\tilde{f}^{-1}(U) \subsetneq I$.
- Then \tilde{f} restricted to $\tilde{f}^{-1}(U)$ is injective, since \tilde{f} only fails injectivity at $0, 2\pi$.
- Then the restricted map $\hat{f} := \tilde{f}|_{\tilde{f}^{-1}(U)} : \tilde{f}^{-1}(U) \rightarrow U$ is a continuous injection and surjects onto its image, thus a bijection
- Claim: \hat{f} is a homeomorphism
 - Define a candidate inverse $\theta = \hat{f}^{-1} : S^1 \rightarrow \mathbb{R}$.
 - Then $f \circ \theta = \text{id}_{S^1}$ implies $e^{i\theta(x)} = x$ for all $x \in U$.

- Letting $V \subseteq f^{-1}(U)$ be open, we have $\theta^{-1}(V) = \widehat{f}(V)$ which (claim?) is open since ???
- So θ is continuous.

Alternatively:

- Take $I = (0, 2\pi)$.
- Then $\tilde{f}(I) = S^1 \setminus \{1\}$, so $U \subseteq \tilde{f}(I)$.
- Claim: $f : S^1 \setminus \{1\} \rightarrow I$ is a homeomorphism.
- Set $\theta(x) = \tilde{f}|_I^{-1} U(x)$; the claim is that this works.
 - Taking a branch cut $\{x + iy \mid x \in [0, \infty), y = 0\}$ for the complex logarithm defines an inverse.

How to prove?

(U, θ) is a smooth coordinate chart:

- Let θ be arbitrary with $e^{i\theta(z)} = z$ and $\theta \subsetneq S^1$.
- $U \subseteq S^1$ is open by assumption.
- We need to show that $\theta : U \rightarrow \varphi(U)$ is a homeomorphism

Exercise (Problem 1.9) Show that \mathbb{CP}^n is a compact $2n$ -dimensional topological manifold, and show how to equip it with a smooth structure, using the correspondence

Recommended problem

$$\begin{aligned} \mathbb{R}^{2n} &\iff \mathbb{C}^n \\ [x^1, y^1, \dots, x^n, y^n] &\iff [x^1 + iy^1, \dots, x^n + iy^n]. \end{aligned}$$

5 Chapter 2

Definition 5.0.1 (Smooth Functionals on Manifolds).

A function $f : M^n \rightarrow \mathbb{R}^k$ is *smooth* iff for every $p \in M$ there exists a smooth chart (U, φ) about p such that $f \circ \varphi^{-1} : \varphi(U) \rightarrow \mathbb{R}^k$ is smooth as a real function.

Fact: $C^\infty(M) := \{f : M \rightarrow \mathbb{R}\}$ is a vector space

Definition 5.0.2 (Coordinate Representations of Functions).

Given a function $f : M \rightarrow \mathbb{R}^k$, the function $\widehat{f} : \varphi(U) \rightarrow \mathbb{R}^k$ where $\widehat{f}(x) = (f \circ \varphi^{-1})(x)$ is a *coordinate representation* of f .

Fact: f is smooth $\iff f$ is smooth (in the above sense) in *some* smooth chart about each point.

Definition 5.0.3 (Smooth Maps Between Manifolds).

$F : M \rightarrow N$ is *smooth* iff for every $p \in M$ there exists charts $p \in (U, \varphi)$ and $F(p) \in (V, \psi)$ such that $F(U) \subseteq V$ and $\psi \circ F \circ \varphi^{-1} : \varphi(U) \rightarrow \psi(V)$ is smooth.

Fact: taking $N = V = \mathbb{R}^k$ and $\psi = \text{id}$ recovers the previous definition.

Proposition 5.1.

Every smooth map between manifolds is continuous.

Proposition 5.2 (Smoothness is Local).

If $F : M \rightarrow N$, then

1. If every $p \in M$ has a neighborhood $U \ni p$ such that F restricted to U is smooth, then F is smooth.
2. If F is smooth, then its restriction to every open subset is smooth.

Definition 5.2.1.

For $F : M \rightarrow N$ and $(U, \varphi), (V, \psi)$ smooth charts in M, N respectively, then $\hat{F} := \psi \circ F \circ \varphi^{-1}$ is the *coordinate representation* of F .

Proposition 5.3.

1. Constant maps $c : M \rightarrow N, c(x) = n_0$ are smooth
2. The identity is smooth
3. Inclusion of open submanifolds $U \hookrightarrow M$ is smooth
4. $F : M \rightarrow N$ and $G : N \rightarrow P$ smooth implies $G \circ F$ is smooth.

Proposition 5.4.

A map $F : N \rightarrow \prod_{i=1}^k M_i$ with at most one i such that $\partial M_i \neq \emptyset$ is smooth iff each component map $\pi_i \circ F : N \rightarrow M_i$ is smooth.

Proving a map between manifolds is smooth:

1. Write the map as a composition of known smooth functions.
2. Write in *smooth local coordinates* and recognize the component functions as compositions of smooth functions

Fact: projection maps from products are smooth

- Every closed subset $A \subseteq M$ of a smooth manifold is the level set of some smooth nonnegative functional $f : M \rightarrow \mathbb{R}$, i.e. $f^{-1}(0) = A$.

6 Chapter 3

Todo list

Thus graphs of continuous functions $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ are locally Euclidean?	5
Coordinates as numbers vs functions?	5
Seems like f is always the *last* coordinate in the graph	7
How is this map a quotient map?	7
Exercise	9
What does "determine the same collection of smooth functions" mean?	11
Recommended problem	13
Why? Should boil down to $x \mapsto x^t$ for $0 \leq t < 1$ failing to be differentiable at 0 in \mathbb{R}	15
Recommended problem	16
Note that the figure should say $\{x^{n+1} = 0\}$ instead of x^n	16
Todo	18
Figure out how to invert.	18
Messy computations that didn't work out.	18
Computation.	18
What are the actual domains and ranges of the transition functions? It seems like you pull back \mathbb{R}^n to $S^n \setminus N$, then push $S^n \setminus \{N, S\}$ to $R^n \setminus 0$, but this yields $\mathbb{R}^n \rightarrow \mathbb{R}^n \setminus 0$ where we haven't deleted zero in the domain (problem: not smooth!).	18
Some way to do this just with covering spaces?	19
How to prove?	20
Recommended problem	20