

# Fundamentals

## Algebra

- Properties of logs:
  - $\ln(\prod) = \sum \ln$  but  $\prod \ln \neq \ln \sum$
  - $\log_b x = \frac{\ln x}{\ln b}$

Be careful!  $\frac{\ln x}{\ln y} \neq \ln \frac{x}{y} = \ln x - \ln y$

- Completing the square:
  - $p(x) = ax^2 + bx + c \implies p(x) = a(x + \frac{b}{2a})^2 + -\frac{1}{2}\left(\frac{b^2-4ac}{2a}\right)$
- Pascal's Triangle:

$n$	Sequence
3	1, 2, 1
4	1, 3, 3, 1
5	1, 4, 6, 4, 1
6	1, 5, 10, 10, 5, 1
7	1, 6, 15, 20, 15, 6, 1
8	1, 7, 21, 35, 35, 21, 7, 1

Obtain new entries by adding in  $\mathcal{T}$  pattern (e.g. 7 = 1+6, 12 = 6 + 15, etc).

Note that  $\binom{n}{i}$  is given by the entry in the  $n$ -th row,  $i$ - column.

Table of Small Factorials

$n$	$n!$
2	2
3	6
4	24
5	120
6	720
7	5040
8	40320
9	362880
10	3628800

$\pi \approx 3.1415926535$

$e \approx 2.71828$

$\sqrt{2} \approx 1.4142135$

Primes Under 100:

2, 3, 5, 7, 11, 13, 17, 19,  
23, 29, 31, 37, 41, 43, 47,  
53, 59, 61, 67,  
71, 73, 79, 83, 89, 97, 101

Checking Divisibility by Small Primes

$p$	$p \mid n \iff$
2	$p \bmod 10 = 2, 4$
3	$\sum \text{digits} \mid 3$
5	$p \bmod 5 = 0, 5$
7	
11	
13	
17	
23	
27	

Geometry

- Generic Conic Sections

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$$

$$\frac{(x - x_0)^2}{w_0} \pm \frac{(y - y_0)^2}{h_0} = c$$

- Circles:

$Ax^2 + By^2 + C = 0$

$(x - x_0)^2 + (y - y_0)^2 = r^2$

- Defining trait: locus of points at a constant distance from the **center**
- **Center** at  $(x_0, y_0)$

- Parabolas:

$Ax^2 + Bx + Cy + D = 0$

$y = ax^2$

- Defining Trait:
  - Locus of points equidistant from the **focus** (a point) and the **directrix** (a line)
  - #todo add image
- **Focus** at  $(0, \frac{1}{4a})$
- **Directrix** at the line  $y = -\frac{1}{4a}$ 
  - For an arbitrary quadratic: complete the square to write in the form  $y = a(x - w_0)^2 + h_0$ , and translate points of interest by  $(x + w_0, y + h_0)$

- Ellipses:

$$\frac{x^2}{w^2} + \frac{y^2}{h^2} = 1$$

- Defining trait:
    - The locus of points where the *sum* of distances to two **focii** are constant.
  - **Center** at  $(0, 0)$  (can translate easily)
  - **Vertices** at  $(\pm w, 0)$  and  $(0, \pm h)$
  - **Focii** at  $F_1 = (\sqrt{w^2 - h^2}, 0), F_2 = (-\sqrt{w^2 - h^2}, 0)$
  - Another useful shortcut form:
- Hyperbolas:

$$\frac{x^2}{w^2} - \frac{y^2}{h^2} = 1$$

- Defining trait:
    - Locus of points where the *difference* between the distances to two **focii** are constant.
  - **Vertices** at  $(0, \pm h)$  and  $(\pm w, 0)$
  - **Focii** at  $F_1 = (\sqrt{w^2 + h^2}, 0), F_2 = (-\sqrt{w^2 + h^2}, 0)$
- Summary of Traits:
    - One point  $p$ :
      - Distance to  $p$  is constant: circle
    - Two points  $a, b$ :
      - Distance to  $a$  equal to distance to  $b$  equals a constant: a line bisecting the midpoint of the line connecting them
      - Difference of distances constant: ellipse
      - Sum of differences constant: hyperbola
    - Point  $p$  and a line  $l$ :
      - Distance to  $p$  equals distance to  $l$  equals a constant: parabola
  - Areas of certain figures:

Shape	Area / Volume
Circle	$\pi r^2$
Annulus	$\pi(r_0 - r_1)^2$
Cylinder	$2\pi rh$
Ellipse	$\frac{1}{2}wh$
Trapezoid:	$\frac{a+b}{2}h$
Any Triangle:	$\frac{1}{2}bh$
Parallelograms:	$bh$
Cones:	?

- Polar coordinates:  $(x, y) \mapsto (\sqrt{x^2 + y^2}, \tan^{-1}(\frac{y}{x}))$
- Spherical Coordinates:  $[\rho, \phi, \theta]$  where

$$x^2 + y^2 + z^1 = \rho^2$$

$$x = \rho \sin \phi \cos \theta$$

$$y = \rho \sin \phi \sin \theta$$

# Trigonometry

- Trig Values
  - Useful note:  $\frac{1}{2} < \frac{\sqrt{2}}{2} < \frac{\sqrt{3}}{2}$

	sin	cos	tan
0	$\frac{\sqrt{0}}{2}$	$\frac{\sqrt{4}}{2}$	0
$\frac{\pi}{6}$	$\frac{\sqrt{1}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{1}}{2}$	$\frac{\sqrt{3}}{1}$
$\frac{\pi}{2}$	$\frac{\sqrt{4}}{2}$	$\frac{\sqrt{0}}{2}$	$\infty$

- Identities
  - $\sin^2 x + \cos^2 x = 1$  (from Pythagorean theorem)
    - Divide through by  $\cos^2 x$  or  $\sin^2 x$  to obtain:
      - $\tan^2 + 1 = \sec^2$
      - $1 + \cot^2 x = \csc^2 x$
    - Just listing what conclusions you can pull out of these permutations:

$$\sin^2 x = 1 - \cos^2 x$$

$$\cos^2 x = 1 - \sin^2 x$$

$$\tan^2 x = \sec^2 - 1$$

$$\csc^2 x = 1 + \cot^2 x$$

$$\sec^2 x = 1 + \tan^2 x$$

$$\cot^2 x = \csc^2 - 1$$

$$1 + \cos^2 x = \text{nothing!}$$

$$1 + \sin^2 x = \text{nothing!}$$

$$1 - \tan^2 x = \text{nothing!}$$

$$1 - \cot^2 x = \text{nothing!}$$

$$\tan^2 x - 1 = \text{nothing!}$$

$$\cot^2 x - 1 = \text{nothing!}$$

- A derivation with multiple payoffs:

$$\cos(a + b) + i \sin(a + b) = e^{i(a+b)}$$

$$= (\cos a + i \sin a)(\cos b - i \sin b)$$

$$= (\cos a \cos b + \sin a \sin b) + i(\sin a \cos b + \cos a \sin b)$$

- $\sin(a + b) = \sin a \cos b + \cos a \sin b$
- $\cos(a + b) = \cos a \cos b - \sin a \sin b$
- $a = b \implies \sin(2a) = 2 \sin a \cos a$
- $a = b \implies \cos(2a) = \cos^2 a - \sin^2 a$

- $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$
- $\cos^2 x = \frac{1}{2}(1 + \cos 2x)$
- $\sin 2x = 2 \sin x \cos x$
- Less useful:  $\tan(a + b) = \frac{\tan a + \tan b}{1 - \tan a \tan b}$

- $\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$

- Law of Cosines

$$a^2 = b^2 + c^2 - 2bc \cos \theta_A$$

- Totally symmetric under any swap of two symbols
- Derivation: pick the vertex corresponding to  $A$ , label the vectors to the other two vertices  $\mathbf{x}, \mathbf{y}$ , then

$$\begin{aligned} |\mathbf{x} - \mathbf{y}|^2 &= \langle \mathbf{x} - \mathbf{y}, \mathbf{x} - \mathbf{y} \rangle \\ &= |\mathbf{x}|^2 + |\mathbf{y}|^2 - 2\langle \mathbf{x}, \mathbf{y} \rangle \\ &= |\mathbf{x}|^2 + |\mathbf{y}|^2 - 2|\mathbf{x}||\mathbf{y}| \cos \theta \end{aligned}$$

## Polynomials

- Vieta's Formulas: Write  $p(x) = \sum a_k x^k = \prod (x_k - r_k)$  and expand the product to obtain

$$p(x) = a_n x^n - \left(\sum_k r_k\right) x^{n-1} + \left(\sum_{i < j} r_i r_j\right) x^{n-2} + \dots = \sum_{k=1}^n (-1)^k \sigma_{n-k}(\{r_i\}_{i=1}^n) x^k$$

where  $\sigma_i$  is the  $i$ -th elementary symmetric sum.

- Example :

$$p(x) = x^2 + bx + c = x^2 - (r_1 + r_2)x + (r_1 r_2)$$

- Example:

$$\begin{aligned} p(x) &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 \\ &= a_3 x^3 - a_3 (r_1 + r_2 + r_3) x^2 + a_3 (r_1 r_2 + r_1 r_3 + r_2 r_3) x - a_3 (r_1 r_2 r_3) \\ &\implies -\frac{a_2}{a_3} = r_1 + r_2 + r_3 \\ &\implies \frac{a_1}{a_3} = r_1 r_2 + r_1 r_3 + r_2 r_3 \\ &\implies \frac{a_0}{a_3} = r_1 r_2 r_3 \end{aligned}$$

- Quick conclusions:
  - Sum of roots of a monic polynomial is the **negative** coefficient of  $x^{n-1}$
  - Product of roots of a monic polynomial is the constant coefficient.

- Common enough to memorize:

$(a + b)^2 =$	$a^2 + b^2 + 2ab$
$(a - b)^2 =$	$a^2 + b^2 - 2ab$
$a^2 + b^2 =$	$(a + b)^2 + 2ab$
$a^2 - b^2 =$	$(a + b)(a - b)$
$(a + b)^3 =$	$a^3 + b^3 + 3(a^2b + ab^2)$
$(a - b)^3 =$	$a^3 - b^3 + 3(-a^2b + ab^2)$
$a^3 + b^3 =$	$(a + b)(a^2 + b^2 - ab)$
$a^3 - b^3 =$	$(a - b)(a^2 + b^2 + ab)$
$(\sqrt{a} + \sqrt{b})^2 =$	$a + b + 2\sqrt{ab}$
$(\sqrt{a} - \sqrt{b})^2 =$	$a + b - 2\sqrt{ab}$
$(a + \sqrt{b})(a - \sqrt{b}) =$	$a^2 - b$
$(a + i\sqrt{b})(a - i\sqrt{b}) =$	$a^2 + b$
$(a + b)(a - b) =$	$a^2 + b^2$

- Polynomial long division
- Rational roots theorem
- Synthetic Division: #todo

## General Techniques:

- Gather everything and interpret as a polynomial in some variable
  - Example: given  $\sinh(x) = \frac{1}{2}e^x - e^{-x}$ , find  $\sinh^{-1} x$ 
    - Let  $u = e^x$ , and note that  $u > 0$  and  $x = \ln u$ .

$$\begin{aligned}
 x &= \frac{1}{2}u + u^{-1} \implies \\
 u - u^{-1} &= 2x \implies \\
 u - u^{-1} - 2x &= 0 \implies \\
 \mathbf{u^2 - 2xu - 1} &= 0 \implies \\
 u &= \frac{1}{2}(2x \pm \sqrt{4x^2 + 4})
 \end{aligned}$$

so  $\sinh^{-1} x = \ln(x + \sqrt{x^2 + 1})$

- Example: you don't always need the quadratic formula. Given  $\cosh x = e^x + e^{-x}$ , find  $\tanh^{-1}(x)$ .

$$\begin{aligned}
 x &= \frac{u - u^{-1}}{u + u^{-1}} \implies \\
 x(u + u^{-1}) - (u - u^{-1}) &= 0 \implies \\
 x(u^2 + 1) - (u^2 - 1) &= 0 \implies \\
 \mathbf{u^2(x - 1) + (x + 1)} &= 0 \implies \\
 u^2 &= \frac{1 + x}{1 - x}
 \end{aligned}$$

- If you see  $x^2 + y^2$ , try adding  $2xy$  to reduce to  $(x + y)^2$
- Finding the minimal polynomial of a number  $a + b$ : #todo

# Single Variable Calculus

## Big Theorems / Tools:

- The Fundamental Theorem of Calculus:

$$\frac{\partial}{\partial x} \int_a^x f(t)dt = f(x)$$
$$\frac{\partial}{\partial x} \int_{a(x)}^{b(x)} g(t)dt = g(b(x))b'(x) - g(a(x))a'(x)$$

- The generalized Fundamental Theorem of Calculus

$$\frac{\partial}{\partial x} \int_{a(x)}^{b(x)} f(x,t)dt - \int_{a(x)}^{b(x)} \frac{\partial}{\partial x} f(x,t)dt = f(x,\cdot) \frac{\partial}{\partial x}(\cdot) \Big|_{a(x)}^{b(x)}$$
$$= f(x,b(x)) b'(x) - f(x,a(x)) a'(x)$$

- Recover FTC by taking  $a(x) = c, b(x) = x, f(x,t) = f(t)$ .
  - Note that if  $f(x,t) = f(t)$  alone, then  $\frac{\partial}{\partial x} f(t) = 0$  and the second integral vanishes
- Extreme Value Theorem
- Involving the Derivative:
  - Mean Value Theorem:

$$f \in C^0(I) \implies \exists p \in I : f(b) - f(a) = f'(p)(b - a)$$

- Useful variant for integrals and average value:

$$f \in C^0(I) \implies \exists p \in I : \int_a^b f(x) \, dx = f(p)(b - a)$$

- Rolle's Theorem
- L'Hopital's Rule: If
  - $f(x), g(x)$  differentiable on  $I - \{\text{pt}\}$
  - $\lim_{x \rightarrow \text{pt}} f(x) = \lim_{x \rightarrow \{\text{pt}\}} g(x) \in \{0, \pm\infty\}$
  - $\forall x \in I, g'(x) \neq 0$
  - $\lim_{x \rightarrow \{\text{pt}\}} \frac{f'(x)}{g'(x)}$  exists

$$\implies \lim_{x \rightarrow \{\text{pt}\}} \frac{f(x)}{g(x)} = \lim_{x \rightarrow \{\text{pt}\}} \frac{f'(x)}{g'(x)}$$

- Taylor Expansions:

$$T(a,x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$
$$= f(a) + f'(a)(x-a) + \frac{1}{2} f''(a)(x-a)^2$$
$$+ \frac{1}{6} f'''(a)(x-a)^3 + \frac{1}{24} f^{(4)}(a)(x-a)^4 + \dots$$

Bounded error:  $|f(x) - T_k(a,x)| < \left| \frac{1}{(k+1)!} f^{(k+1)}(a) \right|$  where  $T_k(a,x)$  is the Taylor series truncated up to and including the  $x^k$  term.

# Differential

## Limits

- Tools for finding  $\lim_{x \rightarrow a} f(x)$ , in order of difficulty:
  - Plug in: equal to  $f(a)$  if continuous
  - L'Hopital's Rule (only for indeterminate forms  $\frac{0}{0}, \frac{\infty}{\infty}$ )
    - For  $\lim f(x)^{g(x)} = 1^\infty, \infty^0, 0^0$ , let  $L = \lim f^g \implies \ln L = \lim g \ln f$
  - Algebraic rules
  - Squeeze theorem
  - Expand in Taylor series at  $a$
  - Monotonic + bounded
- One-sided limits:  $\lim_{x \rightarrow a^-} f(x) = \lim_{\varepsilon \rightarrow 0} f(a - \varepsilon)$
- Limits at zero or infinity:

$$\lim_{x \rightarrow \infty} f(x) = \lim_{\frac{1}{x} \rightarrow 0} f\left(\frac{1}{x}\right) \text{ and } \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow \infty} f(1/x)$$

- Also useful:

$$\lim_{x \rightarrow \infty} \frac{p(x)}{q(x)} = \begin{cases} 0 & \deg p < \deg q \\ \infty & \deg p > \deg q \\ \frac{p_n}{q_n} & \deg p = \deg q \end{cases}$$

- Be careful: limits may not exist!!
  - Example :  $\lim_{x \rightarrow 0} \frac{1}{x} \neq 0$
- Asymptotes:
  - Vertical asymptotes: at values  $x = p$  where  $\lim_{x \rightarrow p} = \pm \infty$
  - Horizontal asymptotes: given by points  $y = L$  where  $L \lim_{x \rightarrow \pm \infty} f(x) < \infty$
  - Oblique asymptotes: for rational functions, divide - terms without denominators yield equation of asymptote (i.e. look at the asymptotic order or “limiting behavior”).
    - Concretely:  $f(x) = \frac{p(x)}{q(x)} = r(x) + \frac{s(x)}{t(x)} \sim r(x)$
- Limit of a recurrence:  $x_n = f(x_{n-1}, x_{n-2}, \dots)$ 
  - If the limit exists, it is a solution to  $x = f(x)$

## Derivatives

- Chain rule:  $\frac{\partial}{\partial x}(f \circ g)(x) = f'(g(x))g'(x)$
- Product rule:  $\frac{\partial}{\partial x} f(x)g(x) = f'g + g'f$ 
  - Note for all rules: always prime the first thing!
- Quotient rule:  $\frac{\partial}{\partial x} \frac{f(x)}{g(x)} = \frac{f'g - g'f}{g^2}$
- Implicit differentiation:  $(f(x))' = f'(x) dx, (f(y))' = f'(y) dy$ 
  - Often able to solve for  $\frac{\partial y}{\partial x}$  this way.
- Obtaining derivatives of inverse functions: if  $y = f^{-1}(x)$  then write  $f(y) = x$  and implicitly differentiate.
- Approximating change:  $\Delta y \approx f'(x)\Delta x$

## Related Rates

General series of steps: want to know some unknown rate  $y_t$

- Lay out known relation that involves  $y$



- Take derivative implicitly (say w.r.t  $t$ ) to obtain a relation between  $y_t$  and other stuff.
- Isolate  $y_t =$  known stuff
- Example: ladder sliding down wall
  - Setup:  $l$ ,  $x_t$  and  $x(t)$  are known for a given  $t$ , want  $y_t$ .
  - $x(t)^2 + y(t)^2 = l^2 \implies 2xx_t + 2yy_t = 2l_t = 0$  (noting that  $l$  is constant)
  - So  $y_t = -\frac{x(t)}{y(t)}x_t$
  - $x(t)$  is known, so obtain  $y(t) = \sqrt{l^2 - x(t)^2}$  and solve.

## Integral

- Average values:

$$f_{\text{avg}}(x) = \frac{1}{b-a} \int_a^b f(t) dt$$

- Proof: apply MVT to  $F(x)$ .
- Area Between Curves
  - Area in polar coordinates:

$$A = \int_{r_1}^{r_2} \frac{1}{2} r^2(\theta) d\theta$$

- Solids of Revolution
  - Disks:  $A = \int \pi r(t)^2 dt$
  - Cylinders:  $A = \int 2\pi r(t)h(t) dt$

## • Arc lengths

$$\begin{aligned} ds &= \sqrt{dx^2 + dy^2} & L &= \int ds \\ &= \int_{x_0}^{x_1} \sqrt{1 + \frac{\partial y}{\partial x}} dx \\ &= \int_{y_0}^{y_1} \sqrt{\frac{\partial x}{\partial y} + 1} dy \end{aligned}$$

- $SA = \int 2\pi r(x) ds$

## Big List of Integration Techniques

Given  $f(x)$ , we want to find an antiderivative  $F(x) = \int f$  satisfying  $\frac{\partial}{\partial x} F(x) = f(x)$

- Guess and check: look for a function that differentiates to  $f$ .
- $u$ - substitution
- Integration by Parts:
  - The standard form:

$$\int u dv = uv - \int v du$$

- A more general form for repeated applications: let  $v^{-1} = \int v$ ,  $v^{-2} = \int \int v$ , etc.

$$\begin{aligned} \int_a^b uv &= uv^{-1} \Big|_a^b - \int_a^b u^1 v^{-1} \\ &= uv^{-1} - u^1 v^{-2} \Big|_a^b + \int_a^b u^2 v^{-2} \\ &= uv^{-1} - u^1 v^{-2} + u^2 v^{-3} \Big|_a^b - \int_a^b u^3 v^{-3} \\ &\vdots \\ \implies \int_a^b uv &= (-1)^n \int_a^b u^n v^{-n} + \sum_{k=1}^n (-1)^k u^{k-1} v^{-k} \Big|_a^b \end{aligned}$$

- Generally useful when one term’s  $n$ th derivative is a constant.
- Shoelace method:
  - Note: you can choose  $u$  or  $v$  equal to 1! Useful if you know the derivative of the integrand.
- Differentiating under the integral

$$\begin{aligned} \frac{\partial}{\partial x} \int_{a(x)}^{b(x)} f(x,t) dt - \int_{a(x)}^{b(x)} \frac{\partial}{\partial x} f(x,t) dt &= f(x,\cdot) \frac{\partial}{\partial x} (\cdot) \Big|_{a(x)}^{b(x)} \\ &= f(x,b(x)) \, b'(x) - f(x,a(x)) \, a'(x) \end{aligned}$$

- Proof: let  $F(x)$  be an antiderivative and compute  $F'(x)$  using the chain rule.
- #todo for constants, this should allow differentiating under the integral when  $f, f_x$  are "jointly continuous"

Derivatives	Integrals	Signs	Result
$u$	$v$	NA	NA
$u'$	$\int v$	+	$u \int v$
$u''$	$\int \int v$	−	$-u' \int \int v$
$\vdots$	$\vdots$	$\vdots$	$\vdots$

Fill out until one column is zero (alternate signs). Get the result column by multiplying diagonally, then sum down the column.

- Trigonometric Substitution

$$\begin{array}{lll} \sqrt{a^2 - x^2} & \Rightarrow & x = a \sin(\theta) \qquad dx = a \cos(\theta) \, d\theta \\ \sqrt{a^2 + x^2} & \Rightarrow & x = a \tan(\theta) \qquad dx = a \sec^2(\theta) \, d\theta \\ \sqrt{x^2 - a^2} & \Rightarrow & x = a \sec(\theta) \qquad dx = a \sec(\theta) \tan(\theta) \, d\theta \end{array}$$

- Partial Fractions
- Completing the Square #todo
- Trig Formulas
  - Double angle formulas:

$$\begin{aligned} \sin^2(x) &= \frac{1}{2}(1 - 2 \cos x) \\ &= \\ &= \\ &= \\ &= \end{aligned}$$

- Products of trig functions
  - Setup:  $\int \sin^a(x) \cos^b(x) dx$ 
    - Both  $a, b$  even:  $\sin(x) \cos(x) = \frac{1}{2} \sin(2x)$
    - $a$  odd:  $\sin^2 = 1 - \cos^2$ ,  $u = \cos(x)$
    - $b$  odd:  $\cos^2 = 1 - \sin^2$ ,  $u = \sin(x)$
  - Setup:  $\int \tan^a(x) \sec^b(x) dx$ 
    - $a$  odd:  $\tan^2 = \sec^2 - 1$ ,  $u = \sec(x)$
    - $b$  even:  $\sec^2 = \tan^2 + 1$ ,  $u = \tan(x)$

## Big Derivative / Integral Table

$\frac{\partial f}{\partial x} \Leftarrow$	$f$	$\Rightarrow \int f dx$
$\frac{1}{2\sqrt{x}}$	$\sqrt{x}$	$\frac{2}{3}x^{\frac{3}{2}}$
$nx^{n-1}$	$x^n, n \neq -1$	$\frac{1}{n+1}x^{n+1}$
$-nx^{-(n+1)}$	$\frac{1}{x^n}, n \neq 1$	$-\frac{1}{n-1}x^{-(n-1)}$
$\frac{1}{x}$	$\ln(x)$	$x \ln(x) - x$
$a^x \ln(a)$	$a^x$	$\frac{a^x}{\ln a}$
$\cos(x)$	$\sin(x)$	$-\cos(x)$
$-\csc(x) \cot(x)$	$\csc(x)$	$\ln  \csc(x) - \cot(x) $
$-\sin(x)$	$\cos(x)$	$\sin(x)$
$\sec(x) \tan(x)$	$\sec(x)$	$\ln  \sec(x) + \tan(x) $
$\sec^2(x)$	$\tan(x)$	$\ln \left  \frac{1}{\cos x} \right $
$-\csc^2(x)$	$\cot(x)$	$\ln  \sin x $
$\frac{1}{1+x^2}$	$\tan^{-1}(x)$	$x \tan^{-1} x - \frac{1}{2} \ln(1+x^2)$
$\frac{1}{\sqrt{1-x^2}}$	$\sin^{-1}(x)$	$x \sin^{-1} x + \sqrt{1-x^2}$
$-\frac{1}{\sqrt{1-x^2}}$	$\cos^{-1}(x)$	$x \cos^{-1} x - \sqrt{1-x^2}$
$\frac{1}{\sqrt{x^2+a}}$	$\ln  x + \sqrt{x^2+a} $	.
$-2 \sin x \cos x$	$\cos^2(x)$	$\frac{1}{2}(x + \sin x \cos x)$
$2 \sin x \cos x$	$\sin^2(x)$	$\frac{1}{2}(x - \sin x \cos x)$
$2 \csc^2(x) \cot(x)$	$\csc^2(x)$	$-\cot(x)$
$2 \sec^2(x) \tan(x)$	$\sec^2(x)$	$\tan(x)$
?	?	?
?	?	?
?	?	?
?	?	?
?	?	?
?	?	?
?	?	?
$(ax+1)e^{ax}$	$xe^{ax}$	$\frac{1}{a^2}(ax-1)e^{ax}$
?	$e^{ax} \sin(bx)$	$\frac{1}{a^2+b^2}e^{ax}(a \sin bx - b \cos bx)$
?	$e^{ax} \cos(bx)$	$\frac{1}{a^2+b^2}e^{ax}(a \sin bx + b \cos bx)$
?	?	?

Other small but useful facts:

$$\int_0^{2\pi} \sin \theta \, d\theta = \int_0^{2\pi} \cos \theta \, d\theta = 0$$

## Optimization

- Critical points: boundary points and wherever  $f'(x) = 0$
- Second derivative test:
  - $f''(p) > 0 \implies p$  is a min
  - $f''(p) < 0 \implies p$  is a max

# Multivariable Calculus

## Notation


$$\phi : \mathbb{R}^n \rightarrow \mathbb{R}, \quad \phi(x_1, x_2, \dots) = \dots$$

$$\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad \mathbf{F}(x_1, x_2, \dots) = [\mathbf{F}_1(x_1, x_2, \dots), \mathbf{F}_2(x_1, x_2, \dots), \dots, \mathbf{F}_n(x_1, x_2, \dots)]$$

$$\vec{v} = [v_1, v_2, \dots]$$

## Partial Derivatives

- Chain Rule: Write out tree of dependent variables:



Then sum each possible path, e.g.

$$\begin{aligned} \left( \frac{\partial z}{\partial x} \right)_y &= \left( \frac{\partial z}{\partial x} \right)_{u,y,v} \\ &+ \left( \frac{\partial z}{\partial v} \right)_{x,y,u} \left( \frac{\partial v}{\partial x} \right)_y \\ &+ \left( \frac{\partial z}{\partial u} \right)_{x,y,v} \left( \frac{\partial u}{\partial x} \right)_{v,y} \\ &+ \left( \frac{\partial z}{\partial u} \right)_{x,y,v} \left( \frac{\partial u}{\partial v} \right)_{x,y} \left( \frac{\partial v}{\partial x} \right)_y \end{aligned}$$

Where the subscripts denote which variables are held constant.

## Approximation and Optimization

- Linear Approximation:
  - $z = f(x, y)$  : use Tangent plane formulation to obtain

$$f(x, y) \approx f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$

- Optimization
  - Critical points of  $f(\vec{x})$  given by points  $\vec{p}_0$  such that  $\nabla f|_{\vec{p}_0} = 0$

- Second derivative test: compute  $H_f(p_0) := \begin{vmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{vmatrix}(\vec{p}_0)$ .
- By cases:
  - $H(\mathbf{p}_0) = 0$ : No conclusion
  - $H(\mathbf{p}_0) < 0$ : Saddle point
  - $H(\mathbf{p}_0) > 0$ :
    - $f_{xx}(\mathbf{p}_0) > 0 \implies$  local min
    - $f_{xx}(\mathbf{p}_0) < 0 \implies$  local max
- Mnemonic: make matrix with  $\nabla f$  as the columns, and then differentiate variables left to right.
- Constrained by domain:
  - Extrema occur on boundaries, so parametrize each boundary to obtain a function in one less variable and apply standard optimization techniques to yield critical points. Test all critical points to find extrema.
- Constrained by an equation:
  - If possible, use constraint to just reduce equation to one dimension and optimize like single-variable case. Otherwise,
  - **Lagrange Multipliers**. The setup:

$$\begin{aligned} &\text{Optimize } f(\mathbf{x}) \\ &\text{subject to } g(\mathbf{x}) = c \\ &\implies \nabla f = \lambda \nabla g \end{aligned}$$

1. Use this formula to obtain a system of equations in the components of  $x$  and the parameter  $\lambda$ .
2. Use  $\lambda$  to obtain a relation involving only components of  $\mathbf{x}$ .
3. Substitute relations **back into constraint** to obtain a collection of critical points.
4. Evaluate  $f$  at critical points to find max/min.

# Geometry in $\mathbb{R}^3$

## Plane Geometry

- Useful to know: rotation matrices

$$\mathbf{R}_\theta = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \implies \mathbf{R}_{\frac{\pi}{2}} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \implies \mathbf{R}_{\frac{\pi}{2}} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ x \end{bmatrix}$$

- Example use: given  $\mathbf{v}$ ,  $\mathbf{R}_{\frac{\pi}{2}} \mathbf{v} \perp \mathbf{v}$ , so useful to obtain normals or other perpendicular vectors in the plane.
- Useful trick: given  $\mathbf{v} = [a, b, c]$ , one perpendicular vector is  $\mathbf{v}^\perp = [c, c, -(a+b)]$  as long as  $\mathbf{v} \neq [-1, -1, 0]$  - in this case, choose  $\mathbf{v}^\perp = [-(b+c), a, a]$ .
- Slope of a line in  $\mathbb{R}^2$ :

$$\mathbf{v} = [x, y] \in \mathbb{R}^2 \implies m = \frac{y}{x}$$

- Normal to a line in  $\mathbb{R}^2$ :

$$m^\perp = \frac{-1}{m} \implies \mathbf{v}^\perp = [-y, x]$$

Important Equations

\$\$

# Lines

$$Ax + By + C = 0 \qquad \mathbf{x} = \mathbf{p} + t\mathbf{v}$$

$$\mathbf{x} \in L \iff \langle \mathbf{x}, \mathbf{n} \rangle = 0?$$

- Determined by a point  $\mathbf{p}$  and a vector  $\mathbf{v}$  on the line.
  - Also determined by two points  $\mathbf{p}_0, \mathbf{p}_1$  by taking  $\mathbf{v} = \mathbf{p}_1 - \mathbf{p}_0$
- Symmetric Equation (sometimes useful)
  - Obtained by isolating  $t$  in each component and setting results equal:

$$(x, y, z) \in L \iff \frac{x - p_x}{v_x} = \frac{y - p_y}{v_y} = \frac{z - p_z}{v_z}$$

(Note that the denominators are just the coefficients of  $t$  in the parametric equation.)

# Planes

$$Ax + By + Cz + D = 0 \qquad ax + by + cz = d \qquad \mathbf{x}(t, s) = \mathbf{p} + t\mathbf{v}_1 + s\mathbf{v}_2$$

$$\mathbf{x} \in P \iff \langle \mathbf{n}, \mathbf{x} - \mathbf{p}_0 \rangle = 0$$

- Determined by a point  $\mathbf{p}_0$  and a normal vector  $\mathbf{n}$ 
  - Also determined by two points  $\mathbf{p}_0, \mathbf{p}_1$  using  $\mathbf{n} = \mathbf{p}_0 \times \mathbf{p}_1$
- **Normal vector to a plane**
  - Can read normal off of equation:  $\mathbf{n} = [a, b, c]$
- Other Facts

$$d = \langle \mathbf{n}, \mathbf{p}_0 \rangle = n_1 p_1 + n_2 p_2 + n_3 p_3$$

- Useful trick: once you compute  $\mathbf{n}$ , you can compute  $d = \langle \mathbf{n}, \mathbf{p} \rangle$  for *any* point in the plane (don't necessarily need to use the one you started with, so pick any point that's convenient to calculate)

# Surfaces

$$S = \{(x, y, z) \mid f(x, y, z) = 0\} \qquad z = f(x, y)$$

- **Tangent plane to a surface:**
  - Need a point  $\mathbf{p}$  and a normal  $\mathbf{n}$ . By cases:
    - $f(x, y, z) = 0$ 
      - $\nabla f$  is a normal vector.
      - Write the tangent plane equation  $\langle \mathbf{n}, \mathbf{x} - \mathbf{p}_0 \rangle$ , done.
    - $z = g(x, y)$ :
      - Let  $f(x, y, z) = g(x, y) - z$ , then  $\mathbf{p} \in S \iff \mathbf{p}$  is in a level set of  $f$ .
      - $\nabla f$  is normal to level sets (and thus the surface), so compute  $\nabla f = [g_x, g_y, -1]$
      - Proceed as in previous case
- **Surfaces of revolution:**
  - Given  $f(x_1, x_2) = 0$ , can be revolved around either the  $x_1$  or  $x_2$  axis.
    - $f(x, y)$  around the  $x$  axis yields  $f(x, \pm\sqrt{y^2 + z^2}) = 0$
    - $f(x, y)$  around the  $y$  axis yields  $f(\pm\sqrt{x^2 + z^2}, y) = 0$
    - Remaining cases proceed similarly - leave the axis variable alone, replace other variable with square root involving missing axis.

- Equations of lines tangent to an intersection of surfaces  $f(x, y, z) = g(x, y, z)$ :
  - Find two normal vectors and take their cross product, e.g.  $\mathbf{n} = \nabla f \times \nabla g$ , then

$$L = \{\mathbf{x} \mid \mathbf{x} = \mathbf{p} + t\mathbf{n}\}$$

- Level curves:
  - Given a surface  $f(x, y, z) = 0$ , the level curves are obtained by looking at e.g.  $f(x, y, c) = 0$ .

## Curves

$$\mathbf{r}(t) = [x(t), y(t), z(t)]$$

- **Tangent line to a curve**
  - Use the fact that  $\mathbf{r}'(t)$  is a tangent vector to  $\mathbf{r}(t)$

$$\mathbf{T}(t) = \mathbf{r}(t_0) + t\mathbf{r}'(t)$$

- **Normal line to a curve**
  - Use the fact that  $\mathbf{r}''(t) \perp \mathbf{r}'(t)$

$$\mathbf{N}(t) = \mathbf{r}(t_0) + t\mathbf{r}''(t)$$

- Special case: Planar Curves and Lines:  $y = f(x)$ ,
  - Let  $g(x, y) = f(x) - y$ , then

$$\nabla g = [f_x(x), -1] \implies m = -\frac{1}{f_x(x)}$$

## Tangent Lines / Planes

- Key insight: just need a point and a normal vector, and the gradient is normal to level sets.  
**The Tangent Plane Equation:** for any locus  $f(\mathbf{x}) = 0$ , we have

$$\mathbf{x} \in T_f(\mathbf{p}_0) \implies \langle \nabla f(\mathbf{p}_0), \mathbf{x} - \mathbf{p}_0 \rangle = 0$$

## Normal Lines

Key insight: the gradient is normal.

To find a normal line, you just need a single point  $\mathbf{p}$  and a normal vector  $\mathbf{n}$ ; then

$$L = \{\mathbf{x} \mid \mathbf{x} = \mathbf{p} + t\mathbf{v}\}$$

## Minimal Distances

Fix a point  $\mathbf{p}$ . Key idea: find a subspace and project onto it.

Key equations: projection and orthogonal projection of  $\mathbf{b}$  onto  $\mathbf{a}$ :

$$\text{proj}_{\mathbf{a}}(\mathbf{b}) = \langle \mathbf{b}, \mathbf{a} \rangle \hat{\mathbf{a}} \qquad \text{proj}_{\mathbf{a}}^{\perp}(\mathbf{b}) = \mathbf{b} - \text{proj}_{\mathbf{a}}(\mathbf{b})$$

- **Point to plane:**
  - Given a plane  $S = \{\mathbf{x} \in \mathbb{R}^3 \mid n_0x + n_1y + n_2z = d\}$ , project onto  $S^{\perp}$  using



$$d = \|\text{proj}_{\mathbf{n}}(\mathbf{p})\|$$

- Given just two vectors  $\mathbf{u}, \mathbf{v}$ : manufacture a normal vector  $\mathbf{n} = \mathbf{u} \times \mathbf{v}$  and continue as above.
- **Point to line:**
  - Given a line  $L : \mathbf{x}(t) = t\mathbf{v}$  for some fixed  $\mathbf{v}$ , use

$$d = \|\text{proj}_{\mathbf{v}}^{\perp}(\mathbf{p})\|$$

- Given a line  $L : \mathbf{x}(t) = \mathbf{w}_0 + t\mathbf{w}$ , let  $\mathbf{v} = \mathbf{x}(1) - \mathbf{x}(0)$  and proceed as above.
- **Line to line:**
  - Given  $\mathbf{r}_1(t) = \mathbf{p}_1 + t\mathbf{v}_1$  and  $\mathbf{r}_2(t) = \mathbf{p}_2 + t\mathbf{v}_2$ :
    - Let  $\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2$ , which is normal to both lines.
    - Then project the vector between any two points onto this normal:

$$d = \|\text{proj}_{\mathbf{n}}(\mathbf{p}_2 - \mathbf{p}_1)\|$$

# Vector Calculus

## Notation

$R$  is a region,  $S$  is a surface,  $V$  is a solid.

$$\oint_{\partial S} \mathbf{F} \cdot d\mathbf{r} = \oint_{\partial S} [\mathbf{F}_1, \mathbf{F}_2, \mathbf{F}_3] \cdot [dx, dy, dz] = \oint_{\partial S} \mathbf{F}_1 dx + \mathbf{F}_2 dy + \mathbf{F}_3 dz$$

## Big Theorems:

- Green’s Theorem:

$$\oint_{\partial R} (L \, dx + M \, dy) = \iint_R \left( \frac{\partial M}{\partial x} - \frac{\partial L}{\partial y} \right) dx dy$$

- Divergence Theorem:

$$\iint_{\partial V} \mathbf{F} \cdot d\mathbf{S} = \iiint_V (\nabla \cdot \mathbf{F}) \, dV$$

- Stokes’ Theorem:

$$\oint_{\partial S} \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S}$$

- Equals zero if  $S$  is a closed surface ( $\partial S = \emptyset$ )
- Computing Areas with Green's Theorem:

$$A(R) = \oint_{\partial R} x \, dy = - \oint_{\partial R} y \, dx = \frac{1}{2} \oint_{\partial R} -y \, dx + x \, dy$$

- $\nabla \times (\nabla \phi) = 0$
- $\nabla \cdot (\nabla \times \mathbf{F}) = 0$

## Definitions

$$\mathbf{a} \cdot \mathbf{b} = \langle \mathbf{x}, \mathbf{y} \rangle = \sum_{i=1}^n x_i y_i = x_1 y_1 + x_2 y_2 + \cdots$$
inner/dot product

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} = \sqrt{\sum_{i=1}^n x_i^2} = \sqrt{x_1^2 + x_2^2 + \cdots}$$
norm

$$\mathbf{a} \times \mathbf{b} = \hat{\mathbf{n}} \|\mathbf{a}\| \|\mathbf{b}\| \sin \theta_{\mathbf{a}, \mathbf{b}} = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$
cross product

$$\nabla := \sum_{i=1}^n \frac{\partial}{\partial x_i} \mathbf{e}_i = \left[ \frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \cdots, \frac{\partial}{\partial x_n} \right]$$
del operator

$$\nabla \phi := \sum_{i=1}^n \frac{\partial \phi}{\partial x_i} \mathbf{e}_i = \left[ \frac{\partial \phi}{\partial x_1}, \frac{\partial \phi}{\partial x_2}, \cdots, \frac{\partial \phi}{\partial x_n} \right]$$
gradient

$$D_{\mathbf{u}}(\phi) = \nabla \phi \cdot \hat{\mathbf{u}}$$
directional derivative

$$\nabla \cdot \mathbf{F} := \sum_{i=1}^n \frac{\partial \mathbf{F}_i}{\partial x_i} = \frac{\partial \mathbf{F}_1}{\partial x_1} + \frac{\partial \mathbf{F}_2}{\partial x_2} + \cdots + \frac{\partial \mathbf{F}_n}{\partial x_n}$$
divergence

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \mathbf{F}_1 & \mathbf{F}_2 & \mathbf{F}_3 \end{vmatrix} = [\mathbf{F}_{3y} - \mathbf{F}_{2z}, \mathbf{F}_{1z} - \mathbf{F}_{3x}, \mathbf{F}_{2x} - \mathbf{F}_{1y}]$$
curl

- Note that the directional derivative uses a normalized direction vector!
- Function Types

$$\nabla : (\mathbb{R}^n \rightarrow \mathbb{R}) \rightarrow (\mathbb{R}^n \rightarrow \mathbb{R}^n)$$

$$\phi \mapsto \nabla \phi := \sum_{i=1}^n \frac{\partial \phi}{\partial x_i} \mathbf{e}_i$$

$$\text{div}(\mathbf{F}) : (\mathbb{R}^n \rightarrow \mathbb{R}^n) \rightarrow (\mathbb{R}^n \rightarrow \mathbb{R})$$

$$\mathbf{F} \mapsto \nabla \cdot \mathbf{F} := \sum_{i=1}^n \frac{\partial \mathbf{F}_i}{\partial x_i}$$

$$\text{curl}(\mathbf{F}) : (\mathbb{R}^3 \rightarrow \mathbb{R}^3) \rightarrow (\mathbb{R}^3 \rightarrow \mathbb{R}^3)$$

$$\mathbf{F} \mapsto \nabla \times \mathbf{F}$$

- Some terminology:

Scalar Field
Vector Field
Gradient Field

$\phi : X \rightarrow \mathbb{R}$ 
 $\mathbf{F} : X \rightarrow \mathbb{R}^n$ 
 $\mathbf{F} : X \rightarrow \mathbb{R}^n \mid \exists \phi : X \rightarrow \mathbb{R} \mid \nabla \phi = F$

- The Gradient: lifts scalar fields on  $\mathbb{R}^n$  to vector fields on  $\mathbb{R}^n$
- Divergence: drops vector fields on  $\mathbb{R}^n$  to scalar fields on  $\mathbb{R}^n$
- Curl: takes vector fields on  $\mathbb{R}^3$  to vector fields on  $\mathbb{R}^3$
- Spherical Coordinates:

$$x = r \cos \theta = \rho \sin \phi \cos \theta$$

$$y = r \sin \theta = \rho \sin \phi \sin \theta$$

# Computations

- Line Integrals Of Curves

- Parametrize the path  $C$  as  $\{\mathbf{r}(t) : t \in [a, b]\}$ , then

$$\begin{aligned}\int_C f \, ds &:= \int_a^b (f \circ \mathbf{r})(t) \|\mathbf{r}'(t)\| \, dt \\ &= \int_a^b f(x(t), y(t), z(t)) \sqrt{x_t^2 + y_t^2 + z_t^2} \, dt\end{aligned}$$

- **Line Integrals of Vector Fields**

- If exact:

$$\frac{\partial}{\partial y} \mathbf{F}_1 = \frac{\partial}{\partial x} \mathbf{F}_2 \implies \int \mathbf{F}_1 \, dx + \mathbf{F}_2 \, dy = \phi(\mathbf{p}_1) - \phi(\mathbf{p}_0)$$

The function  $\phi$  can be found using the same method from ODEs.

- Parametrize the path  $C$  as  $\{\mathbf{r}(t) : t \in [a, b]\}$ , then

$$\begin{aligned}\int_C \mathbf{F} \cdot d\mathbf{r} &:= \int_a^b (\mathbf{F} \circ \mathbf{r})(t) \cdot \mathbf{r}'(t) \, dt \\ &= \int_a^b [\mathbf{F}_1(x(t), y(t), \dots), \mathbf{F}_2(x(t), y(t), \dots)] \cdot [x_t, y_t, \dots] \, dt \\ &= \int_a^b \mathbf{F}_1(x(t), y(t), \dots) x_t + \mathbf{F}_2(x(t), y(t), \dots) y_t + \dots \, dt\end{aligned}$$

- Equivalently written:

$$\int_a^b \mathbf{F}_1 \, dx + \mathbf{F}_2 \, dy + \dots := \int_C \mathbf{F} \cdot d\mathbf{r}$$

in which case  $[dx, dy, \dots] := [x_t, y_t, \dots] = \mathbf{r}'(t)$ .

- Remember to substitute  $dx$  back into the integrand!!

- **Flux Integrals:**

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S \mathbf{F} \cdot \hat{\mathbf{n}} \, dS$$

- **Computing Areas with Green's Theorem**

- Given  $R$  and  $f(x, y) = 0$ 
  - Compute

$$\begin{aligned}\oint_{\partial R} x \, dy &= - \oint_{\partial R} y \, dx \\ &= \frac{1}{2} \oint_{\partial R} -y \, dx + x \, dy = \frac{1}{2} \iint_R 1 - (-1) \, dA = \iint_R 1 \, dA\end{aligned}$$

- Steps:
  - Parametrize  $C$

## Other Results

- $\nabla \cdot \mathbf{F} = 0 \not\implies \exists G : \mathbf{F} = \nabla \times G$ 
  - Counterexample

$$\mathbf{F}(x,y,z) = \frac{1}{\sqrt{x^2+y^2+z^2}}[x,y,z] \text{ , } \quad S = S^2 \subset \mathbb{R}^3$$

$$\implies \nabla \mathbf{F} = 0 \text{ but } \iint_{S^2} \mathbf{F} \cdot d\mathbf{S} = 4\pi \neq 0$$

Where by Stokes’ theorem,

$$\mathbf{F} = \nabla \times \mathbf{G} \implies \iint_{S^2} \mathbf{F} = \iint_{S^2} \nabla \times \mathbf{G} \stackrel{\text{Stokes}}{=} \oint_{\partial S^2} \mathbf{G} \, d\mathbf{r} = 0$$

since  $\partial S^2 = \emptyset$ .

- Sufficient condition:  $\mathbf{F}$  is everywhere  $C^1$

$$\exists \mathbf{G} : \mathbf{F} = \nabla \times \mathbf{G} \iff \forall \text{ closed } S, \iint_S \mathbf{F} \cdot d\mathbf{S} = 0$$

- Recovering Green’s Theorem from Stokes’ Theorem:
  - Let  $\mathbf{F} = [L, M, 0]$ , then  $\nabla \times \mathbf{F} = [0, 0, \frac{\partial M}{\partial x} - \frac{\partial L}{\partial y}]$

# Ordinary Differential Equations

## Techniques Overview

$$p(y)y' = q(x) \qquad \qquad \qquad \text{separable}$$

$$y' + p(x)y = q(x) \qquad \qquad \qquad \text{integrating factor}$$

$$y' = f(x,y), f(tx,ty) = f(x,y) \qquad \qquad y = xV(x) \text{ COV reduces to separable}$$

$$y' + p(x)y = q(x)y^n \qquad \text{Bernoulli, divide by } y^n \text{ and COV } u = y^{1-n}$$

$$M(x,y)dx + N(x,y)dy = 0 \qquad \qquad M_y = N_x : \phi(x,y) = c(\phi_x = M, \phi_y = N)$$

$$P(D)y = f(x,y) \qquad \qquad \qquad x^k e^{rx} \text{ for each root}$$

Where  $e^{zx}$  yields  $e^{ax} \cos bx, e^{ax} \sin bx$

## Ordinary Differential Equations

- Separable equations:

$$p(y)\frac{dy}{dx} - q(x) = 0 \implies \int p(y)dy = \int q(x)dx + C$$

$$\frac{dy}{dx} = f(x)g(y) \implies \int \frac{1}{g(y)}dy = \int f(x)dx + C$$

- Population growth:

$$\frac{dP}{dt} = kP \implies P = P_0 e^{kt}$$

- Logistic growth:

- General form:  $\frac{dP}{dt} = (B(t) - D(t))P(t)$
- Assume birth rate is constant  $B(t) = B_0$  and death rate is proportional to instantaneous population  $D(t) = D_0 P(t)$ . Then let  $r = B_0, C = B_0/D_0$  be the *carrying capacity*:

$$\frac{dP}{dt} = r \left(1 - \frac{P}{C}\right) P \implies P(t) = \frac{P_0}{\frac{P_0}{C} + e^{-rt} \left(1 - \frac{P_0}{C}\right)}$$

- First order linear:

$$\frac{dy}{dx} + p(x)y = q(x) \implies I(x) = e^{\int p(x)dx}, \quad y(x) = \frac{1}{I(x)} \left( \int q(x)I(x)dx + C \right)$$

- Exact:

- $M(x, y)dx + N(x, y)dy = 0$  is exact  $\iff \exists \phi : \frac{\partial \phi}{\partial x} = M(x, y), \frac{\partial \phi}{\partial y} = N(x, y)$

$$\iff \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

- General solution:

$$\phi(x, y) = \int^x M(s, y)ds + \int^y N(x, t)dt - \int^y \frac{\partial}{\partial t} \left( \int^x M(s, t)ds \right) dt$$

(where  $\int^x f(t)dt$  means take the antiderivative of  $f$  and consider it a function of  $x$ )

- Cauchy Euler: #todo
- Bernoulli: \$todo

## Linear Homogeneous

General form:

$$y^{(n)} + c_{n-1}y^{(n-1)} + \dots + c_2y'' + cy' + cy = 0$$

$$p(D)y = \prod (D - r_i)^{m_i} y = 0$$

where  $p$  is a polynomial in the differential operator  $D$  with roots  $r_i$ :

- Real roots: contribute  $m_i$  solutions of the form

$$e^{rx}, xe^{rx}, \dots, x^{m_i-1}e^{rx}$$

- Complex conjugate roots: for  $r = a + bi$ , contribute  $2m_i$  solutions of the form

$$e^{(a \pm bi)x}, xe^{(a \pm bi)x}, \dots, x^{m_i-1}e^{(a \pm bi)x}$$

$$= e^{ax} \cos(bx), e^{ax} \sin(bx), xe^{ax} \cos(bx), xe^{ax} \sin(bx), \dots,$$

Example: by cases, second order equation of the form

$$ay'' + by' + cy = 0$$

- Two distinct roots:  $c_1 e^{r_1 x} + c_2 e^{r_2 x}$
- One real root:  $c_1 e^{rx} + c_2 x e^{rx}$

- Complex conjugates  $\alpha \pm i\beta$ :  $e^{\alpha x}(c_1 \cos \beta x + c_2 \sin \beta x)$

## Linear Inhomogeneous

General form:

$$y^{(n)} + c_{n-1}y^{(n-1)} + \dots + c_2y'' + cy' + cy = F(x)$$

$$p(D)y = \prod (D - r_i)^{m_i} y = 0$$

Then solutions are of the form  $y_c + y_p$ , where  $y_c$  is the solution to the associated homogeneous system and  $y_p$  is a particular solution.

Methods of obtaining particular solutions

## Undetermined Coefficients

- Find an operator  $p(D)$  the annihilates  $F(x)$  (so  $q(D)F = 0$ )
- Find solution of  $q(D)p(D) = 0$ , subtract of known solutions from homogeneous part to obtain the form of the trial solution  $A_0f(x)$ , where  $A_0$  is the undetermined coefficient
- Substitute trial solution into original equation to determine  $A_0$

Useful Annihilators:

$F(x) = p(x) :$	$D^{\deg(p)+1}$
$F(x) = p(x)e^{ax} :$	$(D - a)^{\deg(p)+1}$
$F(x) = \cos(ax) + \sin(ax) :$	$D^2 + a^2$
$F(x) = e^{ax}(a_0 \cos(bx) + b_0 \sin(bx)) :$	$(D - z)(D - \bar{z}) = D^2 - 2aD + a^2 + b^2$
$F(x) = p(x)e^{ax} \cos(bx) + p(x)e^{ax} \cos(bx) :$	$((D - z)(D - \bar{z}))^{\max(\deg(p), \deg(q))+1}$

## Variation of Parameters

#todo

## Reduction of Order

#todo

## Systems of Differential Equations

General form:

$$\frac{\overrightarrow{\partial x(t)}}{\partial t} = A\overrightarrow{x(t)} + \overrightarrow{b(t)} \text{ or } \mathbf{x}'(t) = A\mathbf{x}(t) + \mathbf{b}(t)$$

General solution to homogeneous equation:

$$c_1\overrightarrow{x_1(t)} + c_2\overrightarrow{x_2(t)} + \dots + c_n\overrightarrow{x_n(t)} = X(t)\vec{c}$$

If  $A$  is a matrix of constants:  $\overrightarrow{x(t)} = e^{\lambda_i t} \vec{v}_i$  is a solution for each eigenvalue/eigenvector pair  $(\lambda_i, v_i)$

- If  $A$  is defective: #todo generalized eigenvectors...?

Nonhomogeneous Equation: particular solutions given by  $\vec{x}_p(t) = X(t) \int^t X^{-1}(s) \vec{b}(s) \, ds$

# Laplace Transforms

Definitions:

$$H_a(t) = \begin{cases} 0, & 0 \leq t < a \\ 1, & t \geq a \end{cases}$$
$$\delta(t) : \int_{\mathbb{R}} \delta(t-a) f(t) \, dt = f(a), \quad \int_{\mathbb{R}} \delta(t-a) \, dt = 1$$
$$(f * g)(t) = \int_0^t f(t-s) g(s) \, ds$$

Useful property: for  $a \leq b$ ,  $H_a(t) - H_b(t) = \mathbb{1}_{[a,b]}$ .

$t^n, n \in \mathbb{N}$	$\iff$	$n! \frac{1}{s^{n+1}}, \quad s > 0$
$t^{-\frac{1}{2}}$	$\iff$	$\sqrt{\pi} s^{-\frac{1}{2}}, \quad s > 0$
$e^{at}$	$\iff$	$\frac{1}{s-a}, \quad s > a$
$\cos(bt)$	$\iff$	$\frac{s}{s^2 + b^2}, \quad s > 0$
$\sin(bt)$	$\iff$	$\frac{b}{s^2 + b^2}, \quad s > 0$
$\delta(t-a)$	$\iff$	$e^{-as}$
$H_a(t)$	$\iff$	$s^{-1} e^{-as}$
$e^{at} f(t)$	$\iff$	$F(s-a)$
$H_a(t) f(t-a)$	$\iff$	$e^{-as} F(s)$
$f'(t)$	$\iff$	$sL(f) - f(0)$
$f''(t)$	$\iff$	$s^2 L(f) - sf(0) - f'(0)$
$f^{(n)}(t)$	$\iff$	$s^n L(f) - \sum_{i=0}^{n-1} s^{n-1-i} f^{(i)}(0)$
$f(t)g(t)$	$\iff$	$F(s) * G(s)$

- For  $f$  periodic with period  $T$ ,  $L(f) = \frac{1}{1+e^{-sT}} \int_0^T e^{-st} f(t) \, dt$

# Linear Algebra

Assume everywhere that  $A$  is an  $m \times n$  matrix that represents a linear transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## General Notes

- Rank: number of nonzero rows in RREF
- $\text{Trace}(A) = \sum_{i=1}^m A_{i,i}$
- Elementary row operations / matrices:
  - Permute rows
  - Multiple a row by a scalar

- Add any row to another
- $A(m \times n), B(n \times p), AB = C \implies c_{ij} = \sum_{k=1}^n a_{ik}b_{kj} = \langle \mathbf{a}_i^T, \mathbf{b}_j \rangle$ 
  - i.e., the  $c_{ij}$  entry is just dotting row  $i$  of  $A$  with column  $j$  of  $B$ .

## Systems of Linear Equations

Notation:  $A\vec{x} = \vec{b}$  a linear system,  $r = \text{rank}(A)$  and  $r' = \text{rank}(A \mid \vec{b})$  an augmented matrix.

- Consistent: A system of linear equations is *consistent* when it has at least one solution.
- Inconsistent: A system of linear equations is *inconsistent* when it has no solutions.
- Tall matrices: more equations than unknowns
- Wide matrices: more unknowns than equations
- Three possibilities for a system of linear equations:
  - No solutions
  - One unique solution
  - Infinitely many solutions
- Possibilities:
  - $r < r'$ : the system is inconsistent.
  - $r = r'$ : the system is consistent, and
    - $r' = n \implies$  there is a unique solution (square, tall)
    - $r' < n \implies$  there are infinitely many solutions (wide)
- Homogeneous systems are **always** consistent.

## The Determinant

- Properties of the Determinant  $A : m \times n$ 
  - $\det(AB) = \det(A) \det(B)$
  - Permute two Rows:  $\det A' = -\det A$
  - Factor a scalar  $t$  out of one row:  $\det A' = t \det A$ 
    - $\det(tA) = t^m \det(A)$
  - Add one row to another:  $\det(A') = \det(A)$
  - $\det(L) = \det(U) = \prod_{i=1}^n a_{ii}$  for upper or lower triangular matrices.
  - $\det(A^{-1}) = \frac{1}{\det(A)}$
  - $\det A^k = k \det A$
  - $\det A^T = \det A$
  - $\det(\mathbf{a}_1 + \mathbf{a}_2, \dots) = \det(\mathbf{a}_1, \dots) + \det(\mathbf{a}_2, \dots)$
  - If any row of  $A$  is all zeros,  $\det(A) = 0$ .
  - Take  $A = \begin{pmatrix} \vec{a} \rightarrow \\ \vec{b} \rightarrow \\ \vdots \end{pmatrix}$ , then in  $\mathbb{R}^3$ ,  $\det(A)$  is the volume of the parallelepiped spanned by  $\vec{a}, \vec{b}, \vec{c}$ .

## The Spaces of a Matrix / Linear Map


- Finding bases for various spaces of  $A$ :
  - Row space: reduce to RREF, and take nonzero rows of RREF ( $\subseteq \mathbb{R}^n$ )
  - Column space: reduce to RREF, and take columns with pivots from original  $A$  ( $\subseteq \mathbb{R}^m$ )
  - Nullspace: reduce to RREF, zero rows are free variables, convert back to equations and pull free variables out as scalar multipliers.



# Eigenvalues and Eigenvectors

- Defining equation:  $\lambda \in E(A) \iff \forall x \in \mathbb{R}^m, A\vec{x} = \lambda\vec{x}$
- Finding: solve  $A - I\lambda_i = 0$  for each  $i$ .
- $\lambda \in E(A) \implies \lambda^2 \in E(A^2)$  (with the same eigenvector).
- Eigenvectors corresponding to distinct eigenvalues are **always** linearly independent
- $A$  has  $n$  distinct eigenvalues  $\implies A$  has  $n$  linearly independent eigenvectors.
- Similar matrices have identical eigenvalues and multiplicities.
- A matrix  $A$  is diagonalizable  $\iff A$  has  $n$  linearly independent eigenvectors.
- Useful Facts
  - $\prod \lambda_i = \det A$
  - $\sum \lambda_i = \text{Tr } A$

# Misc

- $|\text{rowspace}(A)| = |\text{colspace}(A)|$
- Proof of Cauchy-Schwarz: See Goode page 346.
- Distance from a point  $p$  to a line  $\vec{a} + t\vec{b}$ : let  $\vec{w} = \vec{p} - \vec{a}$ , then:  $\|w - P(w, v)\|$ 
  -  distance from line to point
- Computing change of basis matrices: #todo
- Two step vector subspace test:
  - Ensure it contains the zero vector
  - Ensure it's closed under scalar multiplication and vector addition
- Any set of two vectors  $\{\vec{v}, \vec{w}\}$  is linearly dependent  $\iff \exists \lambda : \vec{v} = \lambda\vec{w}$ .
- A set of functions  $\{f_i\}$  is linearly independent on  $I \iff \exists x_0 \in I : W(x_0) \neq 0$  (where  $W$  is the Wronskian)
  - NOTE:  $W \equiv 0$  on  $I \not\implies \{f_i\}$  is linearly dependent!
  - Counterexample:  $\{x, x + x^2, 2x - x^2\}$  where  $W \equiv 0$  but  $x + x^2 = 3(x) + (2x - x^2)$ .
  - Sufficient condition: each  $f_i$  is the solution to a linear homogeneous ODE  $L(y) = 0$ .
- Every square matrix is similar to a matrix in jordan canonical form.
- Projection onto column space of  $A$ : given by  $P(\vec{x}) = A(A^t A)^{-1} A^T \vec{x}$
- Normal equations:  $\vec{x}$  is a least squares solution to  $A\vec{x} = \vec{b} \iff A^T A \vec{x} = A^T \vec{b}$

# Gram-Schmidt Process

Extending  $\{\mathbf{x}_i\}$  to an orthonormal basis

$$\begin{aligned} \mathbf{v}_1 &= \mathbf{x}_1 \\ \mathbf{v}_2 &= \mathbf{x}_2 - P(\mathbf{x}_2, \mathbf{v}_1) \\ \mathbf{v}_3 &= \mathbf{x}_3 - P(\mathbf{x}_3, \mathbf{v}_1) - P(\mathbf{x}_3, \mathbf{v}_2) \\ &\dots \\ \mathbf{v}_i &= \mathbf{x}_i - \sum_{k=1}^{i-1} P(\mathbf{x}_i, \mathbf{v}_k) = \mathbf{x}_i - \sum_{k=1}^{i-1} \frac{\langle \mathbf{x}_i, \mathbf{v}_k \rangle}{\|\mathbf{v}_k\|^2} \mathbf{v}_k \end{aligned}$$

# Inverting a Matrix

Equivalent formulas for  $A^{-1}$ :

- Adjoins:  $A^{-1} = \frac{\text{adjugate}(A)}{\det(A)}$
- Gauss Jordan:  $[A \mid I] \sim [I \mid A^{-1}]$

- Cramer's Rule:  $A\vec{x} = \vec{b} \implies x_k = \frac{\det(B_k)}{\det(A)}$  where  $B_k$  is  $A$  where the  $k$ -th column is replaced by  $\vec{b}$

## Big List of Equivalent Properties

Let  $A$  be an  $m \times n$  matrix. TFAE:

- $A$  is invertible and has a unique inverse  $A^{-1}$
- $A^T$  is invertible
- $\det(A) \neq 0$
- The linear system  $A\vec{x} = \vec{b}$  has a unique solution for every  $b \in \mathbb{R}^m$
- The homogeneous system  $A\vec{x} = 0$  has only the trivial solution  $\vec{x} = 0$
- $\text{rank}(A) = m$  (i.e.  $A$  is full rank)
- $\text{nullity}(A) := \dim \text{nullspace}(A) = 0$
- $A = \prod_{i=1}^k E_i$  for some finite  $k$ , where each  $E_i$  is an elementary matrix.
- $A$  is row-equivalent to the identity matrix  $I_n$
- $A$  has exactly  $n$  pivots
- The columns of  $A$  are a basis for  $\mathbb{R}^n$ 
  - i.e.  $\text{colspace}(A) = \mathbb{R}^n$
- The rows of  $A$  are a basis for  $\mathbb{R}^m$ 
  - i.e.  $\text{rowspace}(A) = \mathbb{R}^m$
- $(\text{colspace } A)^\perp = (\text{rowspace } A)^\perp = \{\vec{0}\}$
- Zero is not an eigenvalue of  $A$ .
- $A$  has  $n$  linearly independent eigenvectors
- The rows of  $A$  are coplanar.

As a consequence, all of the following negations are equivalent:

- $A$  is not invertible/singular
- At least one row of  $A$  is a linear combination of the others
- The *RREF* of  $A$  has a row of all zeros.

Reformulated in terms of linear maps  $T$ , TFAE:

- $T^{-1} : \mathbb{R}^m \rightarrow \mathbb{R}^n$  exists
- $\text{im}(T) = \mathbb{R}^n$
- $\ker(T) = 0$
- $T$  is injective
- $T$  is surjective
- $T$  is an isomorphism
- The system  $A\vec{x} = 0$  has infinitely many solutions

## Complex Analysis

- Properties of modulus:
  - $z = a + ib \implies |z| = \sqrt{a^2 + b^2}$
  - $|z|^2 = z\bar{z} = a^2 + b^2$
  - $\frac{z\bar{z}}{|z|^2} = \frac{(a+ib)(a-ib)}{a^2+b^2} = 1$
- $\frac{1}{a+ib} = \frac{1}{z} = \frac{\bar{z}}{|z|^2} = \frac{a-ib}{a^2+b^2}$
- $e^{zx} = e^{(a+ib)x} = e^{ax}(\cos(bx) + i\sin(bx))$

- Complex exponential:  $x^z := e^{z \ln x}$
- $n$ -th roots:  $e^{\frac{ki}{2\pi n}}$
- For  $z = a + bi$ ,  $(x - z)(x - \bar{z}) = x^2 - 2\operatorname{Re}(z)x + (a^2 + b^2)$

# Real Analysis

Summary for GRE exam:

- limits,
- continuity,
- boundedness,
- compactness,
- definitions of topological spaces,
- Lipschitz continuity
- sequences and series of functions.

Notation used throughout:  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ ,  $K$  is a compact set, and "integrable" or  $L_R(K)$  denotes "Riemann integrable on  $K$ ".

## Big Theorems / Formulas

- **Generalized Mean Value Theorem**

$$f, g \text{ differentiable on } [a, b] \implies \exists c \in [a, b] : [f(b) - f(a)] g'(c) = [g(b) - g(a)] f'(c)$$

- Recover MVT: #todo
- **Bolzano-Weierstrass**: every bounded sequence has a convergent subsequence.
- **Heine-Borel**: in  $\mathbb{R}^n$ ,  $X$  is compact  $\iff X$  is closed and bounded.

## Big Examples

- A function continuous and discontinuous at infinitely many points:

$$f(x) = \begin{cases} 0 & x \in \mathbb{R} - \mathbb{Q} \\ \frac{1}{q} & x = \frac{p}{q} \in \mathbb{Q} \end{cases}$$

- Then  $f$  is discontinuous on  $\mathbb{Q}$  and continuous on  $\mathbb{R} - \mathbb{Q}$ . Proof
- Fix  $\varepsilon$ , let  $x_0 \in \mathbb{R} - \mathbb{Q}$ , choose  $n : \frac{1}{n} < \varepsilon$  using Archimedean property.
  - Define  $S = \{x \in \mathbb{Q} : x \in (0, 1), x = \frac{m}{n'}, n' < n\}$
  - Then  $|S| \leq 1 + 2 + \cdots (n - 1)$ , so choose  $\delta = \min_{s \in S} |s - x_0|$
  - Then

$$x \in N_\delta(x_0) \implies f(x) < \frac{1}{n} < \varepsilon \quad \blacksquare$$

- #todo, revisit and spell out more
- Let  $x_0 = \frac{p}{q} \in \mathbb{Q}$  and  $\{x_n\} = \{x - \frac{1}{n\sqrt{2}}\}$ . Then

$$x_n \uparrow x_0 \text{ but } f(x_n) = 0 \rightarrow 0 \neq \frac{1}{q} = f(x_0) \quad \blacksquare$$

# Motivation: Commuting Limit Operations

- Suppose  $f_n \rightarrow f$  (pointwise, not necessarily uniformly)
- Let  $F(x) = \int f(t)$  be an antiderivative of  $f$
- Let  $f'(x) = \frac{\partial f}{\partial x}(x)$  be the derivative of  $f$ .

Then consider the following possible ways to commute various limiting operations:

Does taking the derivative of the integral of a function always return the original function?

$$\left[\frac{\partial}{\partial x}, \int dx\right] : \qquad \frac{\partial}{\partial x} \int f(x,t)dt \stackrel{?}{=} \int \frac{\partial}{\partial x} f(x,t)dt$$

**Answer:** Sort of (but possibly not).

**Counterexample:**

$$f(x) = \begin{cases} 1 & x > 0 \\ -1 & x \leq 0 \end{cases} \implies \int f \approx |x|,$$

which is not differentiable. (This is remedied by the so-called "weak derivative")

**Sufficient Condition:** If  $f$  is continuous, then both are always equal to  $f(x)$  by the FTC.

Is the derivative of a continuous function always continuous?

$$\left[\frac{\partial}{\partial x}, \lim_{x_i \rightarrow x}\right] : \qquad \lim_{x_i \rightarrow x} f'(x_n) \stackrel{?}{=} f'(\lim_{x_i \rightarrow x} x)$$

**Answer:** No.

**Counterexample:**

$$f(x) = \begin{cases} x^2 \sin(1/x) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases} \implies f'(x) = \begin{cases} 2x \sin\left(\frac{1}{x}\right) - \cos\left(\frac{1}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

which is discontinuous at zero.

**Sufficient Condition:** There doesn't seem to be a general one (which is perhaps why we study  $C^k$  functions).

Is the limit of a sequence of differentiable functions differentiable **and** the derivative of the limit?

$$\left[\frac{\partial}{\partial x}, \lim_{f_n \rightarrow f}\right] : \qquad \lim_{f_n \rightarrow f} \frac{\partial}{\partial x} f_n(x) \stackrel{?}{=} \frac{\partial}{\partial x} \lim_{f_n \rightarrow f} f_n(x)$$

**Answer:** *Super* no – even the uniform limit of differentiable functions need not be differentiable!

**Counterexample:**  $f_n(x) = \frac{\sin(nx)}{\sqrt{n}} \Rightarrow f = 0$  but  $f'_n \not\rightarrow f' = 0$

**Sufficient Condition:**  $f_n \Rightarrow f$  and  $f_n \in C^1$ .

Is the limit of a sequence of integrable functions integrable **and** the integral of the limit?

$$\left[\int dx, \lim_{f_n \rightarrow f}\right](f) : \qquad \lim_{f_n \rightarrow f} \int f_n(x)dx =? \int \lim_{f_n \rightarrow f} f_n(x)dx$$

**Answer:** No.

**Counterexample:** Order  $\mathbb{Q} \cap [0, 1]$  as  $\{q_i\}_{i \in \mathbb{N}}$ , then take

$$f_n(x) = \sum_{i=1}^n \mathbb{1}_{[q_n]} \rightarrow \mathbb{1}_{\left[\mathbb{Q} \cap [0, 1]\right]}$$

where each  $f_n$  integrates to zero (only finitely many discontinuities) but  $f$  is not Riemann-integrable.

**Sufficient Condition:** \$

- $f_n \Rightarrow f$ , or
- $f$  integrable and  $\exists M : \forall n, |f_n| < M$  ( $f_n$  uniformly bounded)

Is the integral of a continuous function also continuous?

$$\left[\int dx, \lim_{x_i \rightarrow x}\right] : \qquad \lim_{x_i \rightarrow x} F(x_i) =? F(\lim_{x_i \rightarrow x} x_i)$$

**Answer:** Yes.

**Proof:**  $|f(x)| < M$  on  $I$ , so given  $c$  pick a sequence  $x \rightarrow c$ . Then

$$|f(x)| < M \implies \left|\int_c^x f(t)dt\right| < \int_c^x Mdt \implies |F(x) - F(c)| < M(b-a) \rightarrow 0$$

Is the limit of a sequence of continuous functions also continuous?

$$\left[\lim_{x_i \rightarrow x}, \lim_{f_n \rightarrow f}\right] : \qquad \lim_{f_n \rightarrow f} \lim_{x_i \rightarrow x} f(x_i) =? \lim_{x_i \rightarrow x} \lim_{f_n \rightarrow f} f_n(x_i)$$

**Answer:** No.

**Counterexample:**  $f_n(x) = x^n \rightarrow \delta(1)$

**Sufficient Condition:**  $f_n \Rightarrow f$

Does a sum of differentiable functions necessarily converge to a differentiable function?

$$\left[\frac{\partial}{\partial x}, \sum_{f_n}\right] : \qquad \frac{\partial}{\partial x} \sum_{k=1}^{\infty} f_k =? \sum_{k=1}^{\infty} \frac{\partial}{\partial x} f_k$$

Answer: No.

**Counterexample:**  $f_n(x) = \frac{\sin(nx)}{\sqrt{n}} \Rightarrow 0 := f$ , but  $f'_n = \sqrt{n} \cos(nx) \not\rightarrow 0 = f'$  (at, say,  $x = 0$ )

**Sufficient Condition:** When  $f_n \in C^1, \exists x_0 : f_n(x_0) \rightarrow f(x_0)$ , and  $\sum \|f'_n\|_\infty < \infty$  (continuously differentiable, converges at a point, and the derivatives absolutely converge)

## Continuity

$$f \text{ cts} \iff \lim_{x \rightarrow p} f(x) = f(p)$$

Example of a discontinuous function:  $\sin(\frac{1}{x})$  at  $x = 0$ .

Uniform continuity #todo

## Differentiability

$$f'(p) := \frac{\partial f}{\partial x}(p) = \lim_{x \rightarrow p} \frac{f(x) - f(p)}{x - p}$$

- For multivariable functions: existence and continuity of  $\frac{\partial \mathbf{f}}{\partial x_i} \forall i \implies \mathbf{f}$  differentiable
  - Necessity of continuity: example of a continuous functions with all partial and directional derivatives that is not differentiable:

$$f(x, y) = \begin{cases} \frac{y^3}{x^2 + y^2} & (x, y) \neq (0, 0) \\ 0 & \text{else} \end{cases}$$

## Properties, strongest to weakest

$$C^\infty \subsetneq C^k \subsetneq \text{differentiable} \subsetneq C^0 \subset L_R(K)$$

- Example showing  $f \in C^0 \not\implies f$  is differentiable **and**  $f$  not differentiable  $\not\implies f \notin C^0$ .
  - Take  $f(x) = |x|$  at  $x = 0$ .
- Example showing that  $f$  differentiable  $\not\implies f \in C^1$ :
  - Take

$$f(x) = \begin{cases} x^2 \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases} \implies f'(x) = \begin{cases} -\cos(\frac{1}{x}) + 2x \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$$

but  $\lim_{x \rightarrow 0} f'(x)$  does not exist and thus  $f'$  is not continuous at zero.

Proof that  $f$  differentiable  $\implies f \in C^0$ :

$$f(x) - f(p) = \frac{f(x) - f(p)}{x - p} (x - p) \stackrel{\text{hypothesis}}{=} f'(p)(x - p) \xrightarrow{x \rightarrow p} 0$$

## Giant Table of Relations

Bold are assumed hypothesis, regular text is the strongest conclusion you can reach, strikethrough denotes implications that aren't necessarily true.

$f'$	$f$	$\therefore f$	$F$
<del>exists</del>	<b>continuous</b>	K-integrable	exists
<del>continuous</del>	<b>differentiable</b>	continuous	exists
<del>exists</del>	<b>integrable</b>	<del>continuous</del>	differentiable

Explanation of items in table:

- K-integrable: compactly integrable.
- $f$  integrable  $\implies F$  differentiable  $\implies F \in C^0$ 
  - By definition and FTC, and differentiability  $\implies$  continuity
- $f$  differentiable and  $K$  compact  $\implies f$  integrable on  $K$ .
  - In general,  $f$  differentiable  $\not\implies f$  integrable. Necessity of compactness:

$$f(x) = e^x \in C^\infty(\mathbb{R}) \text{ but } \int_{\mathbb{R}} e^x dx \rightarrow \infty$$

- $f$  integrable  $\not\implies f$  differentiable
  - An integrable function that is not differentiable:  $f(x) = |x|$  on  $\mathbb{R}$
- $f$  differentiable  $\implies f$  continuous a.e.

## Integrability

- Sufficient criteria for integrability:
  - $f$  continuous, montone, bounded, finitely many discontinuities, or
  - Uniformly continuous, or
  - Finitely many discontinuities
- $f$  integrable  $\iff$  bounded and continuous a.e.
  - Prime example of a non-integrable function:  $f = \mathbb{1} \left[ \mathbb{Q} \right]$
- FTC for the Riemann Integral.
  - If  $F$  is a differentiable function on the interval  $[a, b]$ , and  $F'$  is bounded and continuous a.e., then  $F' \in L_R([a, b])$  and

$$\forall x \in [a, b] : \int_a^x F'(t) \, dt = F(x) - F(a)$$

- Suppose  $f$  bounded and continuous a.e. on  $[a, b]$ , and define  $F(x) := \int_a^x f(t) \, dt$ . Then  $F$  is absolutely continuous on  $[a, b]$ , and for  $p \in [a, b]$ ,

$$f \in C^0(p) \implies F \text{ differentiable at } p, \, F'(p) = f(p), \text{ and } F' \stackrel{\text{a.e.}}{=} f.$$

## List of Free Conclusions:

- $f$  integrable on  $U$ :
  - $f$  is bounded
  - $f$  is continuous a.e. (finitely many discontinuities)
  - $F$  is continuous
  - $F$  is differentiable
- $f$  continuous on  $U$ :
  - $f$  is integrable on compact subsets of  $U$
  - $F$  exists
- $f$  differentiable at a point  $p$ :
  - $f$  is continuous

- $f$  is differentiable in  $U$ 
  - $f$  is continuous a.e.
- Defining the Riemann integral: #todo

## Pointwise convergence

$$f_n \rightarrow f = \lim_{n \rightarrow \infty} f_n$$

Summary:

$$\lim_{f_n \rightarrow f} \lim_{x_i \rightarrow x} f_n(x_i) \neq \lim_{x_i \rightarrow x} \lim_{f_n \rightarrow f} f_n(x_i)$$

$$\lim_{f_n \rightarrow f} \int_I f_n \neq \int_I \lim_{f_n \rightarrow f} f_n$$

- Pointwise convergence is strictly weaker than uniform convergence.
  - Example of a function that converges pointwise but not uniformly:  $f_n(x) = x^n$  on  $[0, 1]$
  - Proof: towards a contradiction let  $\varepsilon = \frac{1}{2}$ . Then let  $n = N(\frac{1}{2})$  and  $x = \left(\frac{3}{4}\right)^{\frac{1}{n}}$ : then  $f(x) = 0$  but  $|f_n(x) - f(x)| = x^n = \frac{3}{4} > \frac{1}{2}$ .



- $f_n$  continuous  $\not\implies f$  is continuous
  - i.e. “the pointwise limit of continuous functions is not necessarily continuous”
  - Take

$$f_n(x) = x^n, \quad f_n(x) \rightarrow \mathbb{1}_{[x = 1]}$$

- $f_n$  differentiable  $\not\implies f'_n$  converges
  - Take

$$f_n(x) = \frac{1}{n} \sin(n^2 x) \rightarrow 0, \quad f'_n = n \cos(n^2 x) \text{ does not converge}$$

- $f_n$  integrable  $\not\implies \lim_{f_n \rightarrow f} \int_I f_n \neq \int_I \lim_{f_n \rightarrow f} f_n$ 
  - May fail to converge to same value, take

$$f_n(x) = \frac{2n^2 x}{(1 + n^2 x^2)^2} \rightarrow 0, \quad \int_0^1 f_n = 1 - \frac{1}{n^2 + 1} \rightarrow 1$$

◦

## Uniform Convergence

$$f_n \rightrightarrows f = \lim_{n \rightarrow \infty} f_n \text{ and } \sum_{n=1}^{\infty} f_n \rightrightarrows S$$

Summary:

$$\lim_{x_i \rightarrow x} \lim_{f_n \rightarrow f} f_n(x_i) = \lim_{f_n \rightarrow f} \lim_{x_i \rightarrow x} f_n(x_i) = \lim_{f_n \rightarrow f} f_n(\lim_{x_i \rightarrow x} x_i)$$

$$\lim_{f_n \rightarrow f} \int_I f_n = \int_I \lim_{f_n \rightarrow f} f_n$$



$$\sum_{n=1}^{\infty} \int_I f_n = \int_I \sum_{n=1}^{\infty} f_n$$

"The uniform limit of a(n)  $x$  function is  $x$ ", for  $x \in \{\text{continuous, bounded}\}$

- Equivalent to convergence in the uniform metric on the metric space of bounded functions on  $X$ :

$$f_n \rightrightarrows f \iff \sup_{x \in X} |f_n(x) - f(x)| \rightarrow 0$$

- $(B(X, Y), \|\cdot\|_{\infty})$  is a metric space and  $f_n \rightrightarrows f \iff \|f_n - f\|_{\infty} \rightarrow 0$   
(where  $B(X, Y)$  are bounded functions from  $X$  to  $Y$  and  $\|f\|_{\infty} = \sup_{x \in I} \{f(x)\}$ )
- $f_n \rightrightarrows f \implies f_n \rightarrow f$  pointwise
- $f_n$  continuous  $\implies f$  continuous
  - i.e. "the uniform limit of continuous functions is continuous"
- $f_n \in C^1$ ,  $\exists x_0 : f_n(x_0) \rightarrow f(x_0)$ , and  $f'_n \rightrightarrows g \implies f$  differentiable and  $f' = g$  (i.e.  $f'_n \rightarrow f'$ )
  - Necessity of  $C^1$  – look at failures of  $f'_n$  to be continuous:
    - Take  $f_n(x) = \sqrt{\frac{1}{n^2} + x^2} \rightrightarrows |x|$ , not differentiable
    - Take  $f_n(x) = n^{-\frac{1}{2}} \sin(nx) \rightrightarrows 0$  but  $f'_n \not\rightarrow f' = 0$  and  $f' \neq g$
- $f_n$  integrable  $\implies f$  integrable and  $\int f_n \rightarrow \int f$
- $f_n$  bounded  $\implies f$  bounded
- $f_n \rightrightarrows f_n \not\implies f'_n$  converges
  - Says nothing about it general
- $f'_n \rightrightarrows f' \not\implies f_n \rightrightarrows f$ 
  - Unless  $f$  converges at one or more points.

## Sequences and Metric Spaces

- Big Theorems:
  - **Bolzano-Weierstrass**: every bounded sequence has a convergent subsequence.
  - **Heine-Borel**: in  $\mathbb{R}^n$ ,  $X$  is compact  $\iff X$  is closed and bounded.
    - Necessity of  $\mathbb{R}^n$ :  $X = (\mathbb{Z}, d(x, y) = 1)$  is closed, complete, bounded, but not compact since  $\{1, 2, \dots\}$  has no convergent subsequence
    - Converse holds iff bounded is replaced with totally bounded
  - $X$  compact  $\iff X$  sequentially compact
- $\{x_i\} \rightarrow p \implies$  every subsequence also converges to  $p$ .
- $\{x_i\} \rightarrow p \implies \{x_i\}$  is Cauchy
  - Converse holds in complete metric spaces. Example of a Cauchy sequence that doesn't converge:  $x_i = \pi$  truncated to  $i$  decimal places in  $\mathbb{Q} \subset \mathbb{R}$ .
- $X$  complete and  $X \subset Y \implies X$  closed in  $Y$ 
  - Necessity of completeness:  $\mathbb{Q} \subset \mathbb{Q}$  is closed but  $\mathbb{Q} \subset \mathbb{R}$  is not.
- $X$  compact  $\implies X$  complete and bounded.
  - Holds for any metric space, converse generally does not
- $X$  compact and  $Y \subset X \implies Y$  compact  $\iff Y$  closed.

## Series

- Define  $s_n(x) = \sum_{k=1}^n f_k(x)$  and  $S(x) = \lim_{n \rightarrow \infty} s_n(x)$ , which can converge pointwise or uniformly.

## Sequences and Series of Functions

Notation:  $\sum_{k \in \mathbb{N}} f_k$  is a "series"

- $\limsup |f_k(x)| \neq 0 \implies$  not convergent-  $\limsup |f_k(x)| \neq 0 \implies$  not convergent

# Topology

todo

# Number Theory

- Totient Function

$$\begin{aligned}\phi(p) &= p - 1 \\ \phi(p^k) &= p^{k-1}(p - 1) \\ n = pq, (p, q) &= 1 \implies \phi(n) = \phi(p)\phi(q)\end{aligned}$$

- With these properties, the following formulas can be derived:

$$\begin{aligned}\phi(n) &= \phi\left(\prod_i p_i^{k_i}\right) = \prod_i p_i^{k_i-1}(p_i - 1) \\ &= n \left(\frac{\prod_i (p_i - 1)}{\prod_i p_i}\right) \\ &= n \prod_i \left(1 - \frac{1}{p_i}\right)\end{aligned}$$

- Fermat's Little Theorem

$$\begin{aligned}x^n - x &= 0 \mod n \\ x^{p-1} - 1 &= 0 \mod p\end{aligned}$$

- The Euclidean Algorithm
- The Jacobi symbol

# Abstract Algebra

## To Sort

- Fermat's Little Theorem
- The Euclidean Algorithm
- Burnside's Lemma
- The Sylow Theorems
- Galois Theory
- <http://mathroughguides.wikidot.com/glossary:abstract-algebra>

## Big List of Notation

$C(x) =$	$\{g \in G : gxg^{-1} = x\}$	$\subseteq G$	Centralizer
$C_G(x) =$	$\{gxg^{-1} : g \in G\}$	$\subseteq G$	Conjugacy Class
$G_x =$	$\{g.x : x \in X\}$	$\subseteq X$	Orbit
$x_0 =$	$\{g \in G : g.x = x\}$	$\subseteq G$	Stabilizer
$Z(G) =$	$\{x \in G : \forall g \in G, gxg^{-1} = x\}$	$\subseteq G$	Center
$\text{Inn}(G) =$	$\{\phi_g(x) = gxg^{-1}\}$	$\subseteq \text{Aut}(G)$	Inner Aut.
$\text{Out}(G) =$	$\text{Aut}(G)/\text{Inn}(G)$	$\hookrightarrow \text{Aut}(G)$	Outer Aut.
$N(H) =$	$\{g \in G : gHg^{-1} = H\}$	$\subseteq G$	Normalizer

## Group Theory

Notation:  $H < G$  a subgroup,  $N < G$  a normal subgroup, concatenation is a generic group operation.

- $\mathbb{Z}_n$  the unique cyclic group of order  $n$
- $\mathbf{Q}$  the quaternion group
- $G^n = G \times G \times \dots \times G$
- $Z(G)$  the center of  $G$
- $o(G)$  the order of a group
- $S_n$  the symmetric group
- $A_n$  the alternating group
- $D_n$  the dihedral group of order  $2n$
- Group Axioms
  - Closure:  $a, b \in G \implies ab \in G$
  - Identity:  $\exists e \in G \mid a \in G \implies ae = ea = a$
  - Associativity:  $a, b, c \in G \implies (ab)c = a(bc)$
  - Invertibility:  $a \in G \implies \exists b \in G \mid ab = ba = e$
- Definitions:
  - Order
    - Of a group:  $o(G) = |G|$ , the cardinality of  $G$
    - Of an element:  $o(g) = \min\{n \in \mathbb{N} : g^n = e\}$
  - Index
  - Center: the elements that commute with everything
  - Centralizer: all elements that commute with a given element/subgroup.
  - Group Action: a function  $f : X \times G \rightarrow G$  satisfying
    - $x \in X, g_1, g_2 \in G \implies g_1.(g_2.x) = (g_1g_2).x$
    - $x \in X \implies e.x = x$
  - Orbits partition any set
  - Transitive Action
  - Conjugacy Class:  $C \subset G$  is a conjugacy class  $\iff$ 
    - $x \in C, g \in G \implies gxg^{-1} \in C$
    - $x, y \in C \implies \exists g \in G : gxg^{-1} = y$
    - i.e. subsets that are closed under  $G$  acting on itself by conjugation and on which the action is transitive
    - i.e. orbits under the conjugation action
    - The order of any conjugacy class divides the order of  $G$
  - $p$ -group: Any group of order  $p^n$ .
  - Simple Group: no nontrivial normal subgroups
  - Normal Series:  $0 \trianglelefteq H_0 \trianglelefteq H_1 \dots \trianglelefteq G$
  - Composition Series: The successive quotients of the normal series
  - Solvable:  $G$  is solvable  $\iff G$  has an abelian composition series.
- One step subgroup test:

$$a, b \in H \implies ab^{-1} \in H$$

- Useful isomorphism invariants:
  - Order profile of elements:  $n_1$  elements of order  $p_1$ ,  $n_2$  elements of order  $p_2$ , etc
    - Useful to look at elements of order 2!
  - Order profile of subgroups
  - $Z(A) \cong Z(B)$
  - Number of generators (generators are sent to generators)
  - Number and size of conjugacy classes
  - Number of Sylow- $p$  subgroups.
  - Commutativity
  - “Being cyclic”
  - Automorphism Groups
  - Solvability
  - Nilpotency

## Big Theorems

- Isomorphism Theorems

$$\begin{aligned} \phi : G \rightarrow G' &\implies \frac{G}{\ker \phi} \cong \phi(G) \\ H \trianglelefteq G, K < G &\implies \frac{K}{H \cap K} \cong \frac{HK}{H} \\ H, K \trianglelefteq G, K < H &\implies \frac{G/K}{H/K} \cong \frac{G}{H} \end{aligned}$$

- Lagrange's Theorem:  $H < G \implies o(H) \mid o(G)$ 
  - Converse is false:  $o(A_4) = 12$  but has no order 6 subgroup.
- The  $GZ$  Theorem:  $G/Z(G)$  cyclic  $\implies G \in \mathbf{Ab}$
- Orbit Stabilizer Theorem:  $G/x_0 \cong Gx$
- The Class Equation
  - Let  $G \curvearrowright X$  and  $\mathcal{O}_i \subseteq X$  be the nontrivial orbits, then

$$|X| = |X_0| + \sum_{[x_i] \in X/G} |Gx|$$

- The right hand side is the number of fixed points, plus a sum over all of the orbits of size greater than 1, where any representative within the orbit is chosen and we look at the index of its stabilizer in  $G$ .
- Let  $G \curvearrowright G$  and for each nontrivial conjugacy class  $C_G$  choose a representative  $[x_i] = C_G = C_G(x_i)$  to obtain

$$|G| = |Z(G)| + \sum_{[x_i] = C_G(x_i)} [G : [x_i]]$$

- Useful facts:
  - $H < G \in \mathbf{Ab} \implies H \trianglelefteq G$ 
    - Converse doesn't hold, even if all subgroups are normal. Counterexample:  $\mathbf{Q}$
  - $G/Z(G) \cong \text{Inn}(G)$
  - $H, K < G$  with  $H \cong K \not\implies G/H \cong G/K$ 
    - Counterexample:  $G = \mathbb{Z}_4 \times \mathbb{Z}_2, H = \langle (0, 1) \rangle, K = \langle (2, 0) \rangle$ . Then  $G/H \cong \mathbb{Z}_4 \not\cong \mathbb{Z}_2^2 \cong G/K$

- $G \in \mathbf{Ab} \implies$  for each  $p$  dividing  $o(G)$ , there is an element of order  $p$
- Any surjective homomorphism  $\phi : A \twoheadrightarrow B$  where  $o(A) = o(B)$  is an isomorphism
- Sylow Subgroups:
  - Todo
- Big List of Interesting Groups
  - $\mathbb{Z}_4, \mathbb{Z}_2^2$
  - $D_4$
  - $Q = \langle a, b | a^4 = 1, a^2 = b^2, ab = ba^3 \rangle$  the quaternion group
  - $S^3$ , the smallest nonabelian group
- Chinese Remainder Theorem:

$$\mathbb{Z}_{pq} \cong \mathbb{Z}_p \oplus \mathbb{Z}_q \iff (p, q) = 1$$

- Fundamental Theorem of Finitely Generated Abelian Groups:
  - $G = \mathbb{Z}^n \oplus \bigoplus \mathbb{Z}_{q_i}$
- Finding all of the unique groups of a given order: #todo

## Cyclic Groups

- Generated by ?
- For each  $d$  dividing  $o(G)$ , there exists a subgroup  $H$  of order  $d$ .
  - If  $G = \langle a \rangle$ , then take  $H = \langle a^{\frac{n}{d}} \rangle$

## The Symmetric Group

- Generated by:
  - Transpositions
  - #todo
- Cycle types: characterized by the number of elements in the cycle.
  - Two elements are in the same conjugacy class  $\iff$  they have the same cycle type.
- Inversions: given  $\tau = (p_1 \cdots p_n)$ , a pair  $p_i, p_j$  is *inverted* iff  $i < j$  but  $p_j < p_i$
- Can count inversions  $N(\tau)$ 
  - Equal to minimum number of transpositions to obtain non-decreasing permutation
- Sign of a permutation:  $\sigma(\tau) = (-1)^{N(\tau)}$
- Parity of permutations  $\cong (\mathbb{Z}, +)$ 
  - even  $\circ$  even = even
  - odd  $\circ$  odd = even
  - even  $\circ$  odd = odd

## Ring Theory

- Ring Axioms: #todo
  - Examples:
  - Non-Examples:
- Definition of an Ideal
- Definitions of types of rings:
  - Field
  - Unique Factorization Domain (UFD)
  - Principal Ideal Domain (PID)
  - Euclidean Domain:
  - Integral Domain
  - Division Ring

$$\text{field} \implies \text{Euclidean Domain} \implies \text{PID} \implies \text{UFD} \implies \text{integral domain}$$

Counterexamples to inclusions are strict:

- An ED that is not a field:
- A PID that is not an ED:  $\mathbb{Q}[\sqrt{19}]$
- A UFD that is not a PID:
- An integral domain that is not a UFD:

- Integral Domains
- Unique Factorization Domains
- Prime Elements
- Prime Ideals
- Field Extensions
- The Chinese Remainder Theorem for Rings
- Polynomial Rings
  - Irreducible Polynomials
    - Over  $\mathbb{Z}_2 : x, x + 1, x^2 + x + 1, x^3 + x + 1, x^3 + x^2 + 1$
    - Eisenstein’s Criterion
- Gauss’ Lemma
- When is  $\mathbb{Q}[\sqrt{d}]$  a field? #todo

# Combinatorics

- Choosing:  $\binom{n}{k}$

# Probability

Summary for GRE:

- Calculating Mean, standard deviation, and variance from PDF,
- Bernoulli trials.

# Numerical Analysis

- Euler’s Method:
  - To solve  $\frac{dy}{dx} = f(x, y), y(x_0) = y_0$ , choose a step size  $\varepsilon$ , and let  $x_{n+1} = x_0 + n\varepsilon$ . Then

$$y_{n+1} = y_n + \varepsilon f(x_n, y_n)$$

- Decompositions of Matrices:
  - $LU$
  - Cholesky
  - Singular Value

# Appendix

# Neat Tricks

- Commuting differentials and integrals:

$$\frac{d}{dx} \int_{a(x)}^{b(x)} f(x,t)dt = f(x,b(x))\frac{d}{dx}b(x) - f(x,a(x))\frac{d}{dx}a(x) + \int_{a(x)}^{b(x)} \frac{\partial}{\partial x} f(x,t)dt$$

- Need  $f, \frac{df}{dx}$  to be continuous in both variables. Also need  $a(x), b(x) \in C_1$ .
- If  $a, b$  are constant, boundary terms vanish.
- Recover the fundamental theorem with  $a(x) = a, b(x) = b, f(x,t) = f(t)$ .

## Useful Series and Sequences

Notation:  $\uparrow, \downarrow$ : monotonically converges from below/above.

- Taylor Series:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!}(x-x_0)^n$$

- Cauchy Product:

$$\left(\sum_{k=0}^{\infty} a_k x^k\right) \left(\sum_{k=0}^{\infty} b_i x^n\right) = \sum_{k=0}^{\infty} \left(\sum_{i=0}^k a_n b_n\right) x^k$$

- Differentiation:

$$\frac{\partial}{\partial x} \sum_{k=i}^{\infty} a_k x^k = \sum_{k=i+1}^{\infty} k a_k x^{k-1}$$

- Common Series

$$\sum_{k=0}^N x^k$$

$$= \frac{1 - x^{N+1}}{1 - x}$$

$$\sum_{k=1}^{\infty} x^k$$

$$= \frac{1}{1 - x} \quad \text{for } |x| < 1$$

$$\sum_{k=1}^{\infty} kx^{k-1}$$

$$= \frac{1}{(1 - x)^2} \quad \text{for } |x| < 1$$

$$\sum_{k=2}^{\infty} k(k - 1)x^{k-2}$$

$$= \frac{2}{(1 - x)^3} \quad \text{for } |x| < 1$$

$$\sum_{k=3}^{\infty} k(k - 1)(k - 2)x^{k-3}$$

$$= \frac{6}{(1 - x)^4} \quad \text{for } |x| < 1$$

$$\sum_{k=1}^{\infty} \binom{n}{k} x^k y^{n-k}$$

$$= (x + y)^n$$

$$\sum_{k=1}^{\infty} \frac{x^k}{k}$$

$$= -\log(1 - x)$$

$$\sum_{k=0}^{\infty} \frac{x^k}{k!}$$

$$= e^x$$

$$\sum_{n=0}^{\infty} \frac{(-1)^k}{(2n + 1)!} x^{2k+1}$$

$$= x - \frac{x^3}{3!} + \frac{x^5}{5!}$$

$$\sum_{k=0}^{\infty} \frac{(-1)^k}{(2n)!} x^{2k}$$

$$= 1 - \frac{x^2}{2!} + \frac{x^4}{4!}$$

$$\sum_{k=0}^{\infty} \frac{(-1)^k}{2n + 1} x^{2k+1}$$

$$= x - \frac{x^3}{3} + \frac{x^5}{5}$$

$$\sum_{k=0}^{\infty} \frac{1}{(2k + 1)!} x^{2n+1}$$

$$= x + \frac{x^3}{3!} + \frac{x^5}{5!} + \cdots$$

$$\sum_{k=0}^{\infty} \frac{1}{(2k)!} x^{2k}$$

$$= 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \cdots$$

$$\sum_{k=0}^{\infty} \frac{x^{2k+1}}{2k + 1}$$

$$= \operatorname{arctanh} x$$

$$\sum_{k=1}^{\infty} \frac{1}{k}$$

$$= \infty$$

$$\sum_{k=1}^{\infty} (-1)^k \frac{1}{k}$$

$$= \ln(2)$$

$$\sum_{k=1}^N \frac{1}{k}$$

$$\approx \ln(N) + \gamma + \frac{1}{2N}$$

$$\sum_{k=1}^{\infty} \frac{1}{k^2}$$

$$= \frac{\pi^2}{6}$$

Rational Roots Theorem

Partial Fraction Decomposition

Given  $R(x) = \frac{p(x)}{q(x)}$ , factor  $q(x)$  into  $\prod q_i(x)$ .



- Linear factors of the form  $q_i(x) = (ax + b)^n$  contribute

$$r_i(x) = \sum_{k=1}^n \frac{A_k}{(ax + b)^k} = \frac{A_1}{ax + b} + \frac{A_2}{(ax + b)^2} + \dots$$

- Irreducible quadratics of the form  $q_i(x) = (ax^2 + bx + c)^n$  contribute

$$r_i(x) = \sum_{k=1}^n \frac{A_k x + B_k}{(ax^2 + bx + c)^k} = \frac{A_1 x + B_1}{ax^2 + bx + c} + \frac{A_2 x + B_2}{(ax^2 + bx + c)^2} + \dots$$

- Note:  $ax^2 + bx + c$  is irreducible  $\iff b^2 < 4ac$
- Write  $R(x) = \frac{p(x)}{\prod q_i(x)} = \sum r_i(x)$ , then solve for the unknown coefficients  $A_k, B_k$ .
  - IMPORTANT SHORTCUT: don't try to solve the resulting linear system: for each  $q_i(x)$ , multiply through by that factor and evaluate at its root to zero out many terms!
  - For linear terms  $q_i(x) = (ax + b)^n$ , define  $P(x) = (ax + b)^n R(x)$ ; then

$$A_k = \frac{1}{(n - k)!} P^{(n-k)}(a), \quad k = 1, 2, \dots, n$$

$$\implies A_n = P(a), \quad A_{n-1} = P'(a), \quad \dots, \quad A_1 = \frac{1}{(n - 1)!} P^{(n-1)}(A)$$

- Note: #todo check, might need to evaluate at  $-b/a$  instead, extend to quadratics.