

Title

D. Zack Garza

Saturday 19th September, 2020

Contents

1	Chapter 9	1
2	9.1 and Review	1
2.1	Review Last Time	1
3	9.2	2
3.1	Three steps to gluing theorem	2
4	9.3: Pre-gluing	3

1 | Chapter 9

2 | 9.1 and Review

- (M, ω) a symplectic manifold, $H \in ?$ a Hamiltonian, X_H its ?
- $\int_{S^2} u^* \omega = \sigma_1$ where $u \in C^\infty(S^2, W)$.
- $\langle c_1(TW), \pi_2(TW) \rangle = 0$?
- $C_k(H) := \mathbb{Z}/2\mathbb{Z}[S]$ where S is the set of periodic orbits of X_H of Maslov index k .
- x, y critical points of \mathcal{A}_H with $\mathcal{M}(x, y)$ the moduli space of contractible solutions of finite energy connecting x, y .

2.1 Review Last Time

- $\mathbb{R} \curvearrowright \mathcal{M}_{x,y}$, so we quotient to define $\mathcal{L}(x, y) =: \mathcal{M}_{x,y}/\mathbb{R}$ with the quotient topology.
- Topology defined by when sequences converge:

$$\tilde{u}_n \xrightarrow{n \rightarrow \infty} \tilde{u} \iff \exists \{s_n\} \subseteq \mathbb{R} \text{ such that } u_n(s_n + s, \cdot) \xrightarrow{n \rightarrow \infty} u(s, \cdot).$$

Proposition 2.1(?)

$\mathcal{L}(x, y)$ is Hausdorff.

- Want to show $\mathcal{L}(x, y)$ is a compact 0-dimensional manifold.
- Have a differential

$$\partial : C_k(H) \longrightarrow C_{k-1}(H)$$

$$\partial(x) = \sum_{\text{Ind}(y)=k-1} n(x, y)y.$$

with $n(x, y)$ the number (mod 2) of trajectories of grad \mathcal{A}_H connecting x, y , i.e solutions to the Floer equation.

- Want to prove that the following is a 1-dimensional manifold:

$$M := \overline{\mathcal{L}}(x, z) = \mathcal{L}(x, z) \cup_{\mu(y)=\mu(x)+1} \mathcal{L}(x, y) \times \mathcal{L}(y, z).$$

and show that M is compact with ∂M equal to the last union.

- Last time: closure of space of trajectories connecting x, y contains “broken” trajectories.
- Last time: toward proving that M is compact

3 | 9.2

- Wanted to compactify $\mathcal{L}(x, y)$, needed to go to space of broken trajectories.
- Main theorem of chapter 9: 9.2.1.

Theorem 3.1(9.2.1).

Let (H, J) be a regular pair with H nondegenerate.

Let x, z be two periodic trajectories of H such that $\mu(x) = \mu(z) + 2$.

Then $\overline{\mathcal{L}}(x, y)$ is a compact 1-manifold with boundary satisfying

$$\partial \overline{\mathcal{L}}(x, y) = \bigcup_{\mu(x) < \mu(y) < \mu(z)} \mathcal{L}(x, y) \times \mathcal{L}(y, z).$$

As a corollary, $\partial^2 = 0$.

- Know $\overline{\mathcal{L}}(x, y)$ is compact and $\mathcal{L}(x, y)$ is a 1-manifold
- Now suffices to study in a neighborhood of boundary points (“gluing theorem”)

3.1 Three steps to gluing theorem

1. Pre-gluing: Get a function w_p which interpolates between u and v (not exactly a solution itself, but will be approximated by one later).

2. Constructing ψ a “true solution” from w_p using the Newton-Picard method. We’ll have

$$\psi(p) = \exp_{w_p}(\gamma(p)) \quad \gamma(p) \in W^{1,p}(w_p^* TW) = T_{w_p} \mathcal{P}(x, z).$$

where $\mathcal{P} = ?$.

3. Get a lift $\hat{\psi} = \pi \circ \psi$ where $\pi = ?$ satisfying

- $\hat{\psi}(p) \xrightarrow{n \rightarrow \infty} (\hat{u}, \hat{v})$
- $\hat{\varphi}$ is an embedding
- $\hat{\psi}$ is unique in the following sense (the last point)

Theorem 3.2(9.2.3 (Gluing Theorem)).

Let x, y, z be critical points of the action functional \mathcal{A}_H such that $\mu(x) = \mu(y) + 1 = \mu(z) + 2$. Let $(u, v) \in \mathcal{M}(x, y) \times \mathcal{M}(y, z)$ be trajectories, inducing $(\bar{u}, \bar{v}) \in \mathcal{L}(x, y) \times \mathcal{L}(y, z)$.

- There exist a differentiable map $\psi : (\rho_0, \infty) \rightarrow \mathcal{M}(x, z)$ for some $\rho > 0$ such that
- $\pi \circ \psi : (\rho_0, \infty) \rightarrow \mathcal{L}(x, z)$ is an embedding
- $\hat{\psi} \xrightarrow{\rho \rightarrow \infty} (\bar{u}, \bar{v}) \in \overline{\mathcal{L}(x, z)}$.
- If $\ell_n \in \mathcal{L}(x, z)$ with $\ell_n \xrightarrow{n \rightarrow \infty} (\bar{u}, \bar{v})$, then for $n \gg 1$ we have $\ell \in \mathfrak{F}(\hat{\psi})$.

4 | 9.3: Pre-gluing

- Choose a bump function β on $\{0\}^c \subset \mathbb{R} \rightarrow [0, 1]$ which is 1 on $|x| \geq 1$ and 0 on $|x| < \varepsilon$
- Split into positive and negative parts β^\pm :




Figure 1: Figure 9.3

- Define w in the following way: