

Weil Conjectures

D. Zack Garza

Sunday 19th April, 2020

Contents

1	Notes from Daniel's Office Hours	1
1.1	Definition of Zeta Function	1
1.1.1	Simple but Useful Example: A Point	3
1.1.2	Aside: Why call it a Zeta function?	3
1.1.3	More Examples	4
1.2	Statement of Weil Conjectures	6
1.3	Hard Example: An Elliptic Curve	6

1 Notes from Daniel's Office Hours

0. Definition of Zeta functions
1. Statement of the conjectures
2. Easy examples: $\mathbb{P}_{\mathbb{A}^1}^n, \text{Gr}_{\mathbb{A}^1}(k, n) = \text{GL}(n, \mathbb{A}^1)/P$ the stabilizer of an \mathbb{A}^1 -point in $\mathbb{C}^n, \mathbb{F}_{p^n}$.
3. Medium example: E/\mathbb{A}^1 an elliptic curve.
4. Work out a harder example as in Weil

References

- http://www-personal.umich.edu/~mmustata/zeta_book.pdf

1.1 Definition of Zeta Function

Fix q a prime and $\mathbb{F} := \mathbb{F}_q$ the finite field with q elements, along with its unique degree n extensions

$$\mathbb{F}_n := \mathbb{F}_{q^n} = \left\{ x \in \overline{\mathbb{F}_p} \mid x^{q^n} - x = 0 \right\} \quad \forall n \in \mathbb{Z}^{\geq 2}$$

Definition 1.0.1.

Let

$$J = \langle f_1, \dots, f_N \rangle \trianglelefteq k[x_0, \dots, x_n]$$

be an ideal, then a *projective algebraic variety* $X \hookrightarrow \mathbb{P}_{\mathbb{F}}^{\infty}$ can be given by

$$X = V(J) = \left\{ \mathbf{x} \in \mathbb{P}_{\mathbb{F}}^{\infty} \mid f_1(\mathbf{x}) = \dots = f_N(\mathbf{x}) = \mathbf{0} \right\}$$

where an ideal generated by *homogeneous* polynomials in $n + 1$ variables, i.e. there is some fixed $d \in \mathbb{Z}^{\geq 1}$ such that

$$f(\mathbf{x}) = \sum_{\substack{\mathbf{I}=(i_1, \dots, i_n) \\ \sum_j i_j = d}} \alpha_{\mathbf{I}} \cdot x_0^{i_1} \cdots x_n^{i_n} \quad \text{and} \quad f(\lambda \cdot \mathbf{x}) = \lambda^d f(\mathbf{x}).$$

For the experts: we can take a reduced (possibly reducible) scheme of finite type over a field \mathbb{F}_p . We will be thinking of K -valued points for K/\mathbb{F}_p algebraic extensions. From the audience: what condition do we need to put on such a scheme to guarantee an embedding into \mathbb{P}^∞ ?

Examples:

- Dimension 1: Curves
- Dimension 2: Surfaces
- Codimension 1: Hypersurfaces

Example: Take $f_1(x) = x \in \mathbb{F}[x]$, consider $V(\langle f_1 \rangle) \subset \mathbb{P}_{\mathbb{F}_n}^1$. This is given by the single point $x = \mathbf{0}$.

Fix X/\mathbb{F} an N -dimensional projective algebraic variety. Note that it then has points in any finite extension L/K .

Definition 1.0.2.

Define the *local zeta function* (or *Hasse-Weil zeta function*) of X the following formal power series:

$$\zeta_X(z) = \exp \left(\sum_{n=1}^{\infty} \frac{\alpha_n}{n} z^n \right) \in \mathbb{Q}[[z]] \quad \text{where} \quad \alpha_n := \#X(\mathbb{F}_n).$$

Note the following two properties:

$$\begin{aligned} \zeta_X(0) &= 1 \\ z \left(\frac{\partial}{\partial z} \right) \log \zeta_X(z) &= t \left(\frac{\zeta'_X(z)}{\zeta_X(z)} \right) = \sum_{n=1}^{\infty} \alpha_n z^n = \alpha_1 z + \alpha_2 z^2 + \cdots, \end{aligned}$$

which is an *ordinary generating function* for the sequence (α_n) .

Thus if we define $G(x)$ to be the OGF for (α_n) , we have $\zeta_X(t) = \exp$

Todo: why not an OGF.

Remark: Note that for an OGF $F(x) = \sum_{n=0}^{\infty} f_n x^n$, we can extract coefficients in the following way:

$$[x^n]F(x) = [x^n]T_{F,0}(x) = \frac{1}{n!} \left(\frac{\partial}{\partial x} \right)^n F(x) \Big|_{x=0}.$$

Using the Residue theorem, we can also extract in the following way:

$$[x^n]F(x) = \frac{1}{2\pi i} \oint_{\mathbb{S}^1} \frac{F(z)}{z^{n+1}}.$$

1.1.1 Simple but Useful Example: A Point

Take $X = \{x = 0\} / \mathbb{F}$ a single point over \mathbb{F} , then

$$\begin{aligned}\#X(\mathbb{F}) &:= \alpha_1 = 1 \\ \#X(\mathbb{F}_2) &:= \alpha_2 = 1 \\ &\vdots \\ \#X(\mathbb{F}_n) &:= \alpha_n = 1 \\ &\vdots\end{aligned}$$

Recall that by integrating a geometric series we can derive

$$\begin{aligned}\log(1+t) &= \sum_{n=1}^{\infty} (-1)^{n-1} \frac{t^n}{n} \\ \implies \log(1-t) &= - \sum_{n=1}^{\infty} \frac{t^n}{n} \\ \implies -\log(1-t) &= \sum_{n=1}^{\infty} \frac{t^n}{n} \\ &= 1 \cdot t + 1 \cdot t^2 + 1 \cdot t^3 + \dots\end{aligned}$$

and so

$$\zeta_{\{\text{pt}\}}(t) = \exp(-\log(1-t)) = \frac{1}{1-t}.$$

1.1.2 Aside: Why call it a Zeta function?

Knowing the above calculation, we can now make a precise analogy.

Suppose

$$\mathbb{A}_{\mathbb{Z}}^n \supseteq X = V(\langle f_1, \dots, f_d \rangle) \quad \text{where} \quad f_i \in \mathbb{Z}[x_0, \dots, x_{n-1}].$$

Then for every prime, we can reduce the equations mod p and consider

$$\mathbb{A}_{\mathbb{F}_p}^n \supseteq X_p := V(\langle f_1 \bmod p, \dots, f_d \bmod p \rangle) \quad \text{where} \quad f_i \bmod p \in \mathbb{F}_p[x_0, \dots, x_{n-1}]$$

Then define the “local at p ” zeta function:

$$L_X(s) = \prod_{p \text{ prime}} \zeta_{X_p}(p^{-s}).$$

Note: the index set for the product may require some minor adjustment over \mathbb{Q} in general. There are also potentially modifications needed to extend to schemes.

Taking $X = \operatorname{Spec} \mathbb{Q}$ and $X_p = \operatorname{Spec} \mathbb{F}_p$ (which is a single point since \mathbb{F}_p is a field) and noting that

$$\begin{aligned}\zeta_{X_p}(z) &= \exp\left(\sum_{n=1}^{\infty} \frac{1}{n} z^n\right) \\ &= \exp(-\log(1-z)) \\ &= \frac{1}{1-z},\end{aligned}$$

we find that

$$\begin{aligned}L_X(s) &= \prod_{p \text{ prime}} \zeta_{X_p}(p^{-s}) \\ &= \prod_{p \text{ prime}} \left(\frac{1}{1-p^{-s}}\right) \\ &= \zeta(s),\end{aligned}$$

the classical Riemann Zeta function.

Moreover, it is a theorem (difficult, not proved here!) that for any variety X/\mathbb{F}_p , we have

$$\zeta_X(t) = \prod_{x \in X_{\text{cl}}} \left(\frac{1}{1-t^{\deg(x)}}\right) \xrightarrow{t=p^{-s}} \zeta_X(s) = \prod_{x \in X_{\text{cl}}} \left(\frac{1}{1-p^{-s \deg(x)}}\right).$$

Note that this immediately implies that $\zeta_X(t) \in \mathbb{Z}[[t]]$ is a *rational* function.

Note for experts: $\zeta_X(z)$ an honest generating function for the 0-cycles on X ($F(X_{\text{cl}})$) where are effective (nonnegative coefficients).

1.1.3 More Examples

Example (Affine Line): $X = \mathbb{A}^1/\mathbb{F}$ the affine line over \mathbb{F} , then Note that we can write

$$\mathbb{A}^1(\mathbb{F}_n) = \left\{ \mathbf{x} = [x_1] \mid x_1 \in \mathbb{F}_n \right\}$$

as the set of one-component vectors with entries in \mathbb{F}_n , so

$$\begin{aligned}X(\mathbb{F}) &= q \\ X(\mathbb{F}_2) &= q^2 \\ &\vdots \\ X(\mathbb{F}_n) &= q^n.\end{aligned}$$

Thus

$$\zeta_X(z) = \exp\left(\sum_{n=1}^{\infty} \frac{q^n}{n} z^n\right) = \frac{1}{1-qt}.$$

Example (Affine Space): Set $X = \mathbb{A}^m/\mathbb{F}$, affine m -space over \mathbb{F} , so we can just repeat with now m coordinates

$$\mathbb{A}^1(\mathbb{F}_n) = \left\{ \mathbf{x} = [x_1, \dots, x_m] \mid x_i \in \mathbb{F}_n \right\}$$

Counting yields

$$\begin{aligned} X(\mathbb{F}) &= q^m \\ X(\mathbb{F}_2) &= (q^2)^m \\ &\vdots \\ X(\mathbb{F}_n) &= (q^n)^m. \end{aligned}$$

Thus

$$\zeta_X(z) = \exp \left(\sum_{n=1}^{\infty} \frac{q^{nm}}{n} z^n \right) = \frac{1}{1 - q^m z}.$$

Example (Projective Line): $X = \mathbb{P}^1/\mathbb{F}$ the projective line over \mathbb{F} , then

$$\begin{aligned} X(\mathbb{F}) &= q + 1 \\ X(\mathbb{F}_2) &= q^2 + 1 \\ &\vdots \\ X(\mathbb{F}_n) &= q^n + 1 \\ &\cdot \end{aligned}$$

where we write $\mathbb{P}^1_{\mathbb{F}} = \mathbb{A}^1_{\mathbb{F}} \coprod \{\infty\}$ is the affine line with a point at infinity. We can also count by coordinates:

$$\mathbb{P}^1(\mathbb{F}^n) = \left\{ [x_1, x_2] \mid x_1, x_2 \neq 0 \in \mathbb{F}^n \right\} / \sim = \left\{ [x_1, 1] \mid x_1 \in \mathbb{F}^n \right\} \coprod \{[1, 0]\}.$$

Example (Affine Space): Take $X = \mathbb{A}^n/\mathbb{F}$, then $\alpha_n = q^m + 1$ for a point at infinity, so

$$X(\mathbb{F}) = .$$

Thus

$$\zeta_X(t) = \frac{1}{(1 - q^{-t})(1 - q^{1-t})}.$$

Example (Projective Space): Take $X = \mathbb{P}^n_{\mathbb{F}}$, then $\alpha_n = 1 + q^m + (q^m)^2 + \dots + (q^m)^n$, so

$$\zeta_X(t) = \left(\frac{1}{1 - q^{-t}} \right) \left(\frac{1}{1 - q^{1-t}} \right) \left(\frac{1}{1 - q^{2-t}} \right) \cdots \left(\frac{1}{1 - q^{n-t}} \right),$$

1.2 Statement of Weil Conjectures

or equivalently, take your favorite curve $\gamma \in \mathbb{C}$ homotopic to \mathbb{S}^1 .

Note: this is extremely amenable to numerical approximation if you have a closed form for F or even just a black-box numerical version of F ! I.e. easy to throw at a computer.

Todo: how to manually count points in \mathbb{P}^n !

Example: Take $X = \text{Gr}_{\mathbb{F}}(k, n)$, then ????? so

$$\zeta_X(t) = ?.$$

Questions about properties

- $\zeta_{X \coprod Y}(t) = ? \zeta_X(t) \zeta_Y(t)$?
- $\zeta_{X \times Y} = ?$

1.2 Statement of Weil Conjectures

1. (Rationality)

$$\zeta_X(t) = \frac{p_1(t)p_3(t) \cdots p_{2N-1}(t)}{p_0(t)p_2(t) \cdots p_N(t)} \in \mathbb{Z}(t), \quad \text{i.e.} \quad p_i(t) \in \mathbb{Z}[t]$$

$$\begin{aligned} P_0(t) &= 1 - t \\ P_{2n}(t) &= 1 - q^n t \\ P_i(t) &= \prod_j (1 - a_{ij}t), \quad a_{ij} \in \mathbb{C}. \end{aligned}$$

2. (Functional Equation and Poincare Duality)

$$\zeta_X(n-t) = \pm q^{\frac{1}{2}(nE) - Et} \zeta(x, t).$$

3. (Riemann Hypothesis)
4. (Betti Numbers)

1.3 Hard Example: An Elliptic Curve

Take $X = E/\mathbb{F}$, then $\alpha_n = q^n - (a^n + \bar{a}^n - 1)$ where $|a|_{\mathbb{C}} = |\bar{a}|_{\mathbb{C}} = \sqrt{q}$. Then

$$\zeta_X(t) = \frac{(1 - aq^{-t})(1 - \bar{a}q^{-t})}{(1 - q^{-t})(1 - q^{1-s})}.$$