

# Title

*D. Zack Garza*

# Table of Contents

## Contents

|   |   |
|---|---|
| Table of Contents                               | 2 |
| 1 Lecture 08: Computing Étale Cohomology        | 3 |
| 1.1 Čech to Derived Spectral Sequence . . . . . | 3 |
| 1.2 Mayer-Vietoris . . . . .                    | 4 |

# 1 | Lecture 08: Computing Étale Cohomology

Recall the definition of  $\check{C}(U/X, \mathcal{F})$  and  $\check{C}(X_{\text{ét}}, \mathcal{F})$  (??).

## ⚠ Warning 1.0.1

$\check{H}^i(X_{\text{ét}}, \mathcal{F}) \neq H^i(X_{\text{ét}}, \mathcal{F})$  in general, but by a theorem of Milne this is true if  $X$  is quasicompact any finite subset is contained in an affine open. This is true if  $X$  is quasiprojective.

**Remark 1.0.2:** There is a version for which this does always work where Čech covers are replaced with **hypercovers**.

## 1.1 Čech to Derived Spectral Sequence

Since we have enough injectives, take an injective resolution  $\mathcal{F} \rightarrow \mathcal{I}^0 \rightarrow \dots$  which is exact and each  $\mathcal{I}^j$  is injective. We can apply the Čech complex functor to obtain a double complex

$$\check{C}^\bullet(U/X, \mathcal{I}^0) \rightarrow \check{C}^\bullet(U/X, \mathcal{I}^1) \rightarrow \dots,$$

where the horizontal differentials come from the resolution and the vertical come from the Čech complex. To any double complex, one can associate two spectral sequences. First consider taking horizontal cohomology:

$$\begin{array}{ccccccc} & \vdots & & \vdots & & \ddots & \\ & \uparrow & & \uparrow & & & \\ \check{C}(U/X, \mathcal{I}^1) = \mathcal{I}^0(U \times_X U) & \longrightarrow & \mathcal{I}^1(U \times_X U) & \longrightarrow & \dots & & \\ & \uparrow & & \uparrow & & & \\ \check{C}(U/X, \mathcal{I}^0) = \mathcal{I}^0(U) & \longrightarrow & \mathcal{I}^1(U) & \longrightarrow & \dots & & \end{array}$$

Taking the vertical cohomology yields

$$E_2^{i,j} = \check{H}^i(U, \mathcal{H}^j(\mathcal{F})),$$

where  $\mathcal{H}^j$  is the presheaf  $V \mapsto H_{\text{ét}}^j(V, \mathcal{F})$ .

Now we take cohomology in the other order: taking the vertical cohomology collapses to the bottom row, which are global sections, and so

$$E_2^{i,j} = H^i(\Gamma(X, \mathcal{I})) E_\infty,$$

which is the derived functor cohomology. The spectral sequence thus converges in the following way:

$$\check{H}^i(U, \mathcal{H}^j(\mathcal{F})) \Rightarrow H^{i+j}(X_{\text{ét}}, \mathcal{F}).$$

**Exercise 1.1.1 (Good for getting used to spectral sequences):** Show that if  $\check{C}(X_{\text{ét}}, \cdot)$  is exact on  $\text{Sh}^{\text{ab}}(X_{\text{ét}})$ , then  $\check{H}^* \cong H^*$ . See Tohoku or Hartshorne, and prove this using the Čech to derived functor spectral sequence.

## 1.2 Mayer-Vietoris

**Proposition 1.2.1 (?)**.

Let  $U = U_0 \cup U_1$  with each  $U_i$  a Zariski open subset. Then there exists a functorial LES

$$\cdots \rightarrow H^s(U, \mathcal{F}) \xrightarrow{\text{Res}} H^s(U_0, \mathcal{F}) \oplus H^s(U_1, \mathcal{F}) \xrightarrow{\text{Res}} H^s(U_0 \cap U_1, \mathcal{F}) \xrightarrow{\delta} H^{s+1}(U, \mathcal{F}) \rightarrow \cdots.$$

*Proof (?)*.

Apply the Čech to derived spectral sequence to the cover  $\mathcal{U} := U_0 \amalg U_1 \rightarrow U$ . This says take

$$\mathcal{F}(\mathcal{U}) \rightarrow \mathcal{F}((\mathcal{U})^{\times_U 2}) \rightarrow \mathcal{F}((\mathcal{U})^{\times_U 3}) \rightarrow \cdots.$$


None of these objects are empty, which doesn't happen with the usual Čech complex of an open cover, where the alternating complex is taken which doesn't see all of these.

**Claim:** This complex is quasi-isomorphic to the 2-term complex

$$\mathcal{F}(U_0 \amalg U_1) = \mathcal{F}(U_0) \times \mathcal{F}(U_1) \rightarrow \mathcal{F}(U_0 \cap U_1).$$

**Exercise 1.2.2 (?)**: Prove this. This uses the fact that we have a Zariski cover instead of a general étale cover, since it's not true in general: a counterexample is  $\mathbb{G}_m \rightarrow \mathbb{G}_m$  where  $x \mapsto x^2$ . The double intersection won't make sense, since it won't be connected and there's not a distinguished component.

Given this,  $E_2$  vanishes outside of 2 columns, and considering computing  $H^1$  we have the following situation:



This is a general phenomenon: a spectral sequence collapsing onto two columns is the same data as a long exact sequence.

**Exercise 1.2.3 (?)**: Check this. ■

**Theorem 1.2.4(?)**.

Suppose  $X$  is a scheme and  $\mathcal{F} \in \mathrm{QCoh}(X)$  (for example,  $\mathcal{F} := \mathcal{O}_X$ ). Then there is a canonical isomorphism

$$H^i(X, \mathcal{F}) \xrightarrow{\sim} H^i(X_{\text{ét}}, \mathcal{F}^{\text{ét}}) \xrightarrow{\sim} H^i(X_{\text{fppf}}, \mathcal{F}^{\text{fppf}}),$$

The first term is the usual Zariski cohomology of a quasicoherent sheaf, the second is the étale cohomology where  $\mathcal{F}^{\text{ét}}$  is the associated sheaf on the étale site given by pulling back to an étale morphism, and the third is the same on fppf site, since the categories  $\mathrm{QCoh}$  are canonically isomorphic.

**Remark 1.2.5:** The derived functor cohomology

$$H^i(\mathrm{Sh}(X_{\text{zar}}), \mathcal{F}) := \mathrm{Ext}_{\mathrm{Sh}(X_{\text{zar}})}^i(\mathbb{Z}, \mathcal{F})$$

since it only depends on the category of sheaves. This is because we're taking the derived functors of  $\Gamma$ , which is the same as  $\mathrm{Hom}(\mathbb{Z}, \cdot)$ . This is equal to

$$H^i(\mathrm{QCoh}(X), \mathcal{F}) := \mathrm{Ext}_{\mathrm{QCoh}(X)}^i(\mathcal{O}_X, \mathcal{F})$$

for the same reason, since it's true for any sheaf of  $\mathcal{O}_X$  modules. This isomorphism is not just formal, since  $\mathrm{QCoh}$  is much smaller than  $\mathrm{Sh}$ . The reason is that injective quasicoherent sheaves are flasque (?), so it also computes derived functor cohomology. In general, it's much harder to be an injective object in the  $\mathrm{Sh}$  than it is in  $\mathrm{QCoh}$ , since it has to satisfy a lifting property with respect to more maps.

The main takeaway: we already showed  $\mathrm{QCoh}(X_{\text{ét}}) \cong \mathrm{QCoh}(X_{\text{zar}})$ , but now we have this isomorphism in a much larger category.

*Proof (?)*.

We'll prove this in a special case: if  $X$  is quasicompact and separated, Čech cohomology computes derived functor cohomology.

**Claim 1:** Every cover can be refined to a *finite* cover by affines, using quasicompactness.

**Claim 2:** Supposing  $X$  is affine and  $U \rightarrow X$  is an fppf affine cover, then  $\check{C}U/X, \mathcal{F}$  is exact if  $\mathcal{F} = \widehat{M}$  is the quasicoherent sheaf associated to some module  $M$ .

*Proof (of claim 2).*

Let  $U = \mathrm{Spec} B$ ,  $X = \mathrm{Spec} A$ , and  $M \in A\text{-mod}$ .

Then we get the complex

$$M \rightarrow M \otimes_A B \rightarrow M \otimes_A B^{\otimes_A 2} \rightarrow M \otimes_A B^{\otimes_A 4} \rightarrow \dots,$$

which is the *Amitsur complex*. We showed that  $M$  was the kernel of the first map, and we argued that since  $B$  was an fppf  $A$ -algebra, we can check this after tensoring with  $B$ , in which case we had a section which yielded a nullhomotopy. This complex is exact as in the argument used in the proof of descent. ■

By this claim, we know that

$$\check{H}^i(U/X, \mathcal{F}) = \begin{cases} \mathcal{F}(X) & i = 0 \\ 0 & i > 0. \end{cases}$$

for  $\mathcal{F}$  quasicoherent and  $U, X$  affine.

**Claim 3:** When  $\mathcal{F}$  is quasicoherent and  $X$  affine,

$$\check{H}^i(X_{\text{ét}}, \mathcal{F}) = \begin{cases} \mathcal{F}(X) & i = 0 \\ 0 & i > 0. \end{cases}$$

*Proof (?)*.

Affine covers are cofinal in the diagram of covers. We're taking a direct limit and every time we have something zero, we kill it by refining the cover. ■

■