Linearization Continued

D. Zack Garz

Linearization Continued Section 8.4 Follow-Up

D. Zack Garza

April 2020

Linearization Continued

D. Zack Garza

The Floer equation is given by

$$\frac{\partial u}{\partial s} + J(u)\frac{\partial u}{\partial t} + \operatorname{grad} H_t(u) = 0.$$

– We fixed a solution and lifted it to a sphere:

$$u \in C^{\infty}(S^1 \times \mathbb{R}; W) \quad \mapsto \quad \tilde{u} \in C^{\infty}(S^2; W)$$

- We use the assumption: For every $w \in C^{\infty}(S^2, W)$ there exists a symplectic trivialization of the fiber bundle w^*TW , i.e. $\langle c_1(TW), \pi_2(W) \rangle =$ 0 where c_1 denotes the first Chern class of the bundle TW.
- We use this trivialize the pullback \tilde{u}^*TW to obtain an orthonormal unitary frame

$$\{Z_i\}_{i=1}^{2n} \subset T_{u(s,t)}W$$

Linearization Continued

– We used the chosen frame $\{Z_i\}$ to define a chart centered at u of $\mathcal{P}^{1,p}(x,y)$ given by


$$\iota: W^{1,p}\left(\mathbb{R} \times S^1; \mathbb{R}^{2n}\right) \longrightarrow \mathcal{P}^{1,p}(x,y)$$
$$Y = (y_1, \dots, y_{2n}) \longmapsto \exp_u\left(\sum y_i Z_i\right).$$

– We regard Y(s,t) as a tangent vector to W in some Euclidean embedding.

Linearization Continued

D. Zack Garz

- We seek to compute the composite map in charts:



Linearization Continued

D. Zack Garz

Extract the part that is linear in *Y* and collect terms:

$$(d\mathcal{F})_{u}(Y) = \left(\frac{\partial Y}{\partial s} + J(u)\frac{\partial Y}{\partial t}\right) + \left((dJ)_{u}(Y)\frac{\partial u}{\partial t} - (dJ)_{u}(Y)X_{t} - J(u)(dX_{t})_{u}(Y)\right)$$

.