

Problem Set 1

D. Zack Garza

Sunday 30th August, 2020

Contents

Source: Section 1 of Gathmann

Exercise 0.1 (1.19).

Prove that every affine variety $X \subset \mathbb{A}^n/k$ consisting of only finitely many points can be written as the zero locus of n polynomials.

Hint: Use interpolation. It is useful to assume at first that all points in X have different x_1 -coordinates.

Exercise 0.2 (1.21).

Determine \sqrt{I} for

$$I := \langle x_1^3 - x_2^6, x_1x_2 - x_2^3 \rangle \subseteq \mathbb{C}[x_1, x_2].$$

Exercise 0.3 (1.22).

Let $X \subset \mathbb{A}^3/k$ be the union of the three coordinate axes. Compute generators for the ideal $I(X)$ and show that it can not be generated by fewer than 3 elements.

Exercise 0.4 (1.23: Relative Nullstellensatz).

Let $Y \subset \mathbb{A}^n/k$ be an affine variety and define $A(Y)$ by the quotient

$$\pi : k[x_1, \dots, x_n] \longrightarrow A(Y) := k[x_1, \dots, x_n]/I(Y).$$

- Show that $V_Y(J) = V(\pi^{-1}(J))$ for every $J \subseteq A(Y)$.
- Show that $\pi^{-1}(I_Y(X)) = I(X)$ for every affine subvariety $X \subseteq Y$.
- Using the fact that $I(V(J)) \subseteq \sqrt{J}$ for every $J \subseteq k[x_1, \dots, x_n]$, deduce that $I_Y(V_Y(J)) \subseteq \sqrt{J}$ for every $J \subseteq A(Y)$.

Conclude that there is an inclusion-reversing bijection

$$\left\{ \begin{array}{c} \text{Affine subvarieties} \\ \text{of } Y \end{array} \right\} \iff \left\{ \begin{array}{c} \text{Radical ideals} \\ \text{in } A(Y) \end{array} \right\}.$$

Exercise 0.5 (Extra).

Let $J \trianglelefteq k[x_1, \dots, x_n]$ be an ideal, and find a counterexample to $I(V(J)) = \sqrt{J}$ when k is not algebraically closed.