

CRAG

D. Zack  
Garza

Zeta  
Functions

# CRAG

## The Weil Conjectures

D. Zack Garza

April 2020

CRAG

D. Zack  
Garza

Zeta  
Functions

## Zeta Functions

Fix  $q$  a prime and  $\mathbb{F} := \mathbb{F}_q$  the (unique) finite field with  $q$  elements, along with its (unique) degree  $n$  extensions

$$\mathbb{F}_{q^n} = \left\{ x \in \overline{\mathbb{F}}_q \mid x^{q^n} - x = 0 \right\} \quad \forall n \in \mathbb{Z}^{\geq 2}$$

## Definition (Zeta Function)

Let  $J = \langle f_1, \dots, f_M \rangle \trianglelefteq k[x_0, \dots, x_n]$  be an ideal, then a *projective algebraic variety*  $X \subset \mathbb{P}_{\mathbb{F}}^n$  can be described as

$$X = V(J) = \left\{ \mathbf{x} \in \mathbb{P}_{\mathbb{F}}^n \mid f_1(\mathbf{x}) = \dots = f_M(\mathbf{x}) = \mathbf{0} \right\}$$

where  $J$  is generated by *homogeneous* polynomials in  $n + 1$  variables, i.e. there is a fixed  $d = \deg f_i \in \mathbb{Z}^{\geq 1}$  such that

$$f(\mathbf{x}) = \sum_{\substack{\mathbf{l}=(i_1, \dots, i_n) \\ \sum_j i_j = d}} \alpha_{\mathbf{l}} \cdot x_0^{i_1} \cdots x_n^{i_n} \quad \text{and} \quad f(\lambda \cdot \mathbf{x}) = \lambda^d f(\mathbf{x}), \lambda \in \mathbb{F}^{\times}.$$

- For a fixed variety  $X$ , we can consider its  $\mathbb{F}$ -points  $X(\mathbb{F})$ .
  - Note that  $\#X(\mathbb{F}) < \infty$  is an integer
- For any  $L/\mathbb{F}$ , we can also consider  $X(L)$ 
  - In particular, we can consider  $X(\mathbb{F}_{q^n})$  for any  $n \geq 2$ .
  - We again have  $\#X(\mathbb{F}_{q^n}) < \infty$  and are integers for every such  $n$ .
- So we can consider the sequence

$$[N_1, N_2, \dots, N_n, \dots] := [\#X(\mathbb{F}), \#X(\mathbb{F}_{q^2}), \dots, \#X(\mathbb{F}_{q^n}), \dots].$$

- Idea: associate some generating function (a formal power series) to this sequence, e.g.

$$F(z) = \sum_{n=1}^{\infty} N_n z^n.$$