

# Title

D. Zack Garza

Saturday 26<sup>th</sup> September, 2020

## Contents

1 Saturday, September 26

1

# 1 | Saturday, September 26

**Remark 1.**

There is a natural action of  $\mathrm{MCG}(\Sigma)$  on  $H_1(\Sigma; \mathbb{Z})$ , i.e. a *homology representation* of  $\mathrm{MCG}(\Sigma)$ :

$$\begin{aligned}\rho : \mathrm{MCG}(\Sigma) &\rightarrow \mathrm{Aut}_{\mathrm{Grp}}(H_1(\Sigma; \mathbb{Z})) \\ f &\mapsto f_*.\end{aligned}$$

**Definition 1.0.1** (Special Linear Group).

$$\mathrm{SL}(n, \mathbb{k}) = \left\{ M \in \mathrm{GL}(n, \mathbb{k}) \mid \det M = 1 \right\} = \ker \det_{\mathbb{G}_m}.$$

**Theorem 1.1** (*Mapping Class Group of the Torus*).

The homology representation of the torus induces an isomorphism

$$\sigma : \mathrm{MCG}(\Sigma_2) \xrightarrow{\cong} \mathrm{SL}(2, \mathbb{Z})$$

**Remark 2.**

$$\mathrm{SL}(2, \mathbb{Z}) = \left\langle S = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, T = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \right\rangle.$$

Note that  $S^2 = 1$  and

$$T^n = \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}$$

---

Moreover, if  $\mathbf{x} = [x_1, x_2] \in \mathbb{Z} \oplus \mathbb{Z}$  and  $A \in \mathrm{SL}(2, \mathbb{Z})$ , we have  $A\mathbf{x} \in \mathbb{Z} \oplus \mathbb{Z}$ , i.e. this preserves t

*Proof .*

- For  $f$  any automorphism, the induced map  $f_* : \mathbb{Z}^2 \rightarrow \mathbb{Z}^2$  is a group automorphism, so we can consider the group morphism

$$\begin{aligned}\tilde{\sigma} : (\mathrm{Map}(X, X), \circ) &\rightarrow (\mathrm{GL}(2, \mathbb{Z}), \circ) \\ f &\mapsto f_*.\end{aligned}$$

- This will descend to the quotient  $\mathrm{MCG}(X)$  iff  $\mathrm{Map}^0(X, X) \subseteq \ker \tilde{\sigma} = \tilde{\sigma}^{-1}(\mathrm{id})$ 
  - This holds because any map in the identity component is homotopic to the identity, and homotopic maps induce the equal maps on homology.
- So we have a (now injective) map

$$\begin{aligned}\tilde{\sigma} : \mathrm{MCG}(X) &\rightarrow \mathrm{GL}(2, \mathbb{Z}) \\ f &\mapsto f_*.\end{aligned}$$

**Claim:**  $\mathrm{im}(\tilde{\sigma}) \subseteq \mathrm{SL}(2, \mathbb{Z})$ .

- We can thus freely restrict the codomain to define the map

$$\begin{aligned}\sigma : \mathrm{MCG}(X) &\rightarrow \mathrm{SL}(2, \mathbb{Z}) \\ f &\mapsto f_*.\end{aligned}$$

**Claim:**  $\sigma$  is surjective.

•

■