

# Title

D. Zack Garza

August 21, 2019

## Contents

0.1 Exercises . . . . .	1
-------------------------	---

### 0.1 Exercises

#### Problem 1.

Let  $C$  denote the Cantor set.

1. Show that  $C$  contains point that is not an endpoint of one of the removed intervals.
2. Show that  $C$  is nowhere dense, meager, and has measure zero.
3. Show that  $C$  is uncountable.

#### Solution 1.

1. First we will characterize the endpoints of the removed intervals. Let  $C_n$  be the  $n$ th stage of the deletion process that is used to define the Cantor set; then what remains is a union of intervals:

$$C_n = [0, \frac{1}{3^n}] \cup [\frac{2}{3^n}, \frac{3}{3^n}] \cup \cdots \cup [\frac{3^n - 1}{3^n}, 1],$$

and so the endpoints are precisely the numbers of the form  $\frac{k}{3^n}$  where  $0 \leq k \leq 3^n$ . Moreover, any endpoint appearing in  $C_n$  is never removed in any later step, and so all endpoints remaining in  $C$  are of this form where we allow  $0 \leq n < \infty$ .

Thus, our goal is to produce a number  $x \in [0, 1]$  such that  $x \neq \frac{k}{3^n}$  for any  $k$  or  $n$ , but also satisfies  $x \in C$ .

Claim: If  $x \in C$ , then one can find a ternary expansion for which all of the digits are either 0 or 2, i.e.

$$x = \sum_{k=1}^{\infty} a_k 3^{-k} \quad \text{where } a_k \in \{0, 2\}.$$

Proof: By induction on the index  $n$  in  $C_n$ , first consider  $C_1 = [0, 1] \setminus [\frac{1}{3}, \frac{2}{3}] = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$ . So if  $x \in C_1$ , then  $x \notin [\frac{1}{3}, \frac{2}{3}]$ . But note that  $a_1$  is computed in the following way:

$$a_1 = \begin{cases} 0 & 0 \leq x < \frac{1}{3}, \\ 1 & \frac{1}{3} \leq x < \frac{2}{3}, \\ 2 & \frac{2}{3} \leq x < 1. \end{cases}$$

Since the interval  $(\frac{1}{3}, \frac{2}{3})$  is deleted in  $C_1$ , we find that  $a_1 = 1$  iff  $x = \frac{1}{3}$ . In this case, however, we can write

$$x = \frac{1}{3} = \sum_{k=1}^{\infty} a_k 3^{-k} \quad \text{where } a_1 = 1, a_{k>1} = 0,$$

which can be expanded as

$$x = \sum_{k=1}^{\infty} b_k 3^{-k} \quad \text{where } b_1 = 0, b_{k>1} = 2$$

which follows from the fact that this is a geometric sum that can be written in closed form,

$$x' = \sum_{k=2}^{\infty} 2 \cdot 3^{-k}$$