

Linearization and Transversality

Sections 8.3 and 8.4

D. Zack Garza

April 2020

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
 H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Review 8.2

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
 H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Section 8.3: The Space of Perturbations of H

Goal

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Goal: Given a fixed Hamiltonian $H \in C^\infty(W \times S^1; \mathbb{R})$, perturb it (without modifying the periodic orbits) so that $\mathcal{M}(x, y)$ are manifolds of the expected dimension.

Goal

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
 H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Start by trying to construct a subspace $\mathcal{C}_\varepsilon^\infty(H) \subset \mathcal{C}^\infty(W \times S^1; \mathbb{R})$, the space of perturbations of H depending on a certain sequence $\varepsilon = \{\varepsilon_k\}$, and show it is a dense subspace.



Define an Absolute Value

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Idea: similar to how you build $L^2(\mathbb{R})$, define a norm $\|\cdot\|_\varepsilon$ on $C_\varepsilon^\infty(H)$ and take the subspace of finite-norm elements.

- Let $h(\mathbf{x}, t) \in C_\varepsilon^\infty(H)$ denote a perturbation of H .
- Fix $\varepsilon = \{\varepsilon_k \mid k \in \mathbb{Z}^{\geq 0}\} \subset \mathbb{R}^{>0}$ a sequence of real numbers, which we will choose carefully later.
- For a fixed $\mathbf{x} \in W$, $t \in \mathbb{R}$ and $k \in \mathbb{Z}^{\geq 0}$, define

$$|d^k h(\mathbf{x}, t)| = \max \left\{ d^\alpha h(\mathbf{x}, t) \mid |\alpha| = k \right\},$$

the maximum over all sets of multi-indices α of length k .

Note: I interpret this as

$$d^{\alpha_1, \alpha_2, \dots, \alpha_k} h = \frac{\partial^k h}{\partial x_{\alpha_1} \partial x_{\alpha_2} \cdots \partial x_{\alpha_k}},$$

the partial derivatives wrt the corresponding variables.

Define a Norm

- Define a norm on $C^\infty(W \times S^1; \mathbb{R})$:

$$\|h\|_{\infty} = \sum_{k \geq 0} \varepsilon_k \sup_{(x,t) \in W \times S^1} |d^k h(x, t)|.$$

- Since $W \times S^1$ is assumed compact (?), fix a finite covering $\{B_i\}$ of $W \times S^1$ such that

$$\bigcup_i B_i^\circ = W \times S^1.$$

- Choose them in such a way we obtain charts

$$\psi_i : B_i \longrightarrow \overline{B(0, 1)} \subset \mathbb{R}^{2n+1} \quad (?).$$

- Obtain the computable form

$$\|h\|_{\infty} = \sum_{k \geq 0} \varepsilon_k \sup_{(x,t) \in W \times S^1} \sup_{i, z \in B(0,1)} |d^k (h \circ \psi_i^{-1})(z)|.$$

Define a Banach Space

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

- Define

$$C_\varepsilon^\infty = \left\{ h \in C^\infty(W \times S^1; \mathbb{R}) \mid \|h\|_\varepsilon < \infty \right\} \subset C^\infty(W \times S^1; \mathbb{R}),$$

which is a Banach space (normed and complete).

- Show that the sequence $\{\varepsilon_k\}$ can be chosen so that C_ε^∞ is a *dense* subspace for the C^∞ topology, and in particular for the C^1 topology.

Theorem

Such a sequence $\{\varepsilon_k\}$ can be chosen.

Lemma

$C^\infty(W \times S^1; \mathbb{R})$ with the C^1 topology is separable as a topological space (contains a countable dense subset).

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
 H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Section 8.4: Linearizing the Floer Equation: The Differential of F