

# Title

D. Zack Garza

Friday 20<sup>th</sup> March, 2020

## Contents

<b>1 Friday February 21st</b>	<b>1</b>
1.1 Singularities . . . . .	1

## 1 Friday February 21st

### 1.1 Singularities

Recall that there are three types of singularities:

- Removable
- Poles
- Essential

#### **Theorem 1.1(3.2).**

An isolated singularity  $z_0$  of  $f$  is a pole  $\iff \lim_{z \rightarrow z_0} f(z) = \infty$ .

#### **Theorem 1.2(3.3, Casorati-Weierstrass).**

If  $f$  is holomorphic and has an essential singularity  $z_0$ , then there exists a radius  $r$  such that  $f(D_r(\{z_0\}) \setminus \{z_0\})$  is dense in  $\mathbb{C}$ .

*Proof.*

Proceed by contradiction. Suppose there exists a  $w \in \mathbb{C}$  and a  $\delta > 0$  such that

$$D_\delta(w) \cap f(D_r(\{z_0\}) \setminus \{z_0\}) = \emptyset.$$

If  $z \in D_r(w) \setminus z_0$ , then  $|f(z) - w| > \delta$ . Let  $g(z) = \frac{1}{f(z) - w}$ ; then  $|g(z)| < \frac{1}{\delta}$ .

