

Title

D. Zack Garza

Sunday 13th September, 2020

Contents

1	Sunday, September 13	1
1.1	General Notes	1
1.2	1.a	2
1.3	3.a	2
1.4	3.b	3

1 | Sunday, September 13

1.1 General Notes

- Say what you're assuming at the start of the proof.
 - If flipping logic and not using a direct proof (contradiction, contrapositive, etc), then signpost/announce it near the beginning of the proof.
 - Examples: for $P \implies Q$,
 - * Direct proof: "Suppose $P \dots$ "
 - * Contradiction: "Suppose P but not $Q \dots$ " (usually show not P . If you show Q , a direct proof might be simpler.)
 - * Contrapositive: "Suppose not $Q \dots$ "
- Put any important equations (i.e. major steps of the proof) on their own lines or in `displaymath` environments.
- Use some whitespace to separate parts of the proof and increase readability.
- Remember that limits of sequences need not exist, but \liminf s/ \limsup s always do (just may be $\pm\infty$).
- Try to avoid abbreviating the names of major theorems (example: "AP" can stand for many results, not just the Archimedean property!)
- It's not generally true that $a \leq M \implies |a| \leq M$, e.g. take $a = -1$. This only holds $a \geq 0$.

- A generic set may not contain its inf or sup. Example: $\inf \left\{ \frac{1}{n} \right\} = 0$ and $0 \notin \left\{ \frac{1}{n} \right\}$, or $\sup \left\{ 1 - \frac{1}{n} \right\} = 1$ with $1 \notin \left\{ 1 - \frac{1}{n} \right\}$.
- If there exists some element of a set or sequence with a given property, try to say where it comes from and why the property holds for it.
- Similarly, if a property holds for all elements of a set or sequence, try to say why.

1.2 1.a

Proof ($A \implies B$).

- Suppose $\{a_n\}$ is not bounded above.
- Then any $k \in \mathbb{N}$ is not an upper bound for $\{a_n\}$.
- So choose a subsequence $a_{n_k} > k$, then by order-limit laws,

$$a_{n_k} > k \implies \liminf_{k \rightarrow \infty} a_{n_k} > \liminf_{k \rightarrow \infty} k = \infty.$$

■

Proof ($\neg A \implies \neg B$).

- Suppose $\{a_n\}$ is bounded by M , so $a_n < M < \infty$ for all $n \in \mathbb{N}$.
- Then if $\{a_{n_k}\}$ is a subsequence, we have $a_{n_k} \in \{a_n\}$, so $a_{n_k} < M$ for all $k \in \mathbb{N}$.
- But then

$$a_{n_k} < M \implies \limsup_{k \rightarrow \infty} a_{n_k} \leq M,$$

- Now note that if $\lim_{k \rightarrow \infty} a_{n_k}$ exists,

$$\lim_{k \rightarrow \infty} a_{n_k} < \limsup_{k \rightarrow \infty} a_{n_k} \leq M < \infty,$$

so every subsequence is bounded and thus can not converge to ∞ .

■

1.3 3.a

Proof (Using definition (i)).

- Suppose $x_n \leq M$ for all n , we will show that every subsequential limit is also bounded by M .
- Let

$$S := \left\{ x \in \mathbb{R} \mid x \text{ is a subsequential limit of } \{x_n\} \right\}$$

be the set of subsequential limits.

– Note that $\inf S := \liminf_{n \rightarrow \infty} x_n$ by definition (i).

- Let $\{x_{n_k}\} \in S$ be an arbitrary convergent subsequence (since we are only concerned about subsequences with well-defined limits).
- Then for every k we have $x_{n_k} \in \{x_n\}$, so

$$|x_{n_k}| \leq M.$$

- By order limit laws,

$$|x_{n_k}| \leq M \implies \lim_{k \rightarrow \infty} |x_{n_k}| \leq M,$$

- Since the map $x \mapsto |x|$ is continuous, using the sequential definition of continuity we can pass the limit through the absolute value to obtain

$$\left| \lim_{k \rightarrow \infty} x_{n_k} \right| \leq M.$$

- Since the subsequence was arbitrary, we find that M is an upper bound for S and so $\sup S \leq M$.
- But

$$\inf S \leq \sup S \leq M \implies \inf S \leq M.$$

■

Proof (Using definition (ii)).

- Suppose $|x_n| \leq M$ for every n , we will directly show that $\left| \lim_{n \rightarrow \infty} \inf_{k \geq n} x_k \right| \leq M$.
- By order-limit laws, for every fixed n we have

$$|x_n| \leq M \iff -M \leq x_n \leq M \implies -M \leq \inf_{k \geq n} x_k \leq M,$$

where we've used the fact that $x_n \geq -M$ for all n implies that $\inf_{k \geq n} x_k \geq -M$.

- Again applying order-limit laws,

$$-M \leq \inf_{k \geq n} x_k \leq M \implies -M \leq \lim_{n \rightarrow \infty} \inf_{k \geq n} x_k \leq M \iff \left| \lim_{n \rightarrow \infty} \inf_{k \geq n} x_k \right| \leq M.$$

■

1.4 3.b

Proof (Using definition (i)).

Note that here we define S to be the set of all subsequential limits of $\{x_n\}$ and

$$\liminf_n x_n := \inf S.$$

- Suppose toward a contradiction that $\beta < \liminf_n x_n$ **but** there does not exist any N such that $n \geq N \implies x_n > \beta$.
- Then for all N there exists an $n > N$ with $x_n \leq \beta$, so the set

$$B := \{n \in \mathbb{N} \mid x_n \leq \beta\}$$

is countably infinite.

- Then the elements of B define a subsequence x_{n_k} which satisfies

$$x_{n_k} \leq \beta \quad \forall k \implies L := \lim_{k \rightarrow \infty} x_{n_k} \leq \beta$$

where we've used order-limit laws.

- We now have $L \in S$, a subsequential limit satisfying $L \leq \beta$ and since $\inf S$ is a lower bound for S ,

$$\inf S \leq L \leq \beta.$$

which contradicts $\beta < \liminf_n x_n$. ■

Proof .

Note that here we define

$$\liminf_n x_n := \lim_{n \rightarrow \infty} S_n \quad \text{where} \quad S_n := \inf \{x_k \mid k \geq n\}.$$

- Write $L := \lim_{n \rightarrow \infty} S_n$ and suppose $\beta < L$.
- Then we have

$$\forall \varepsilon > 0, \exists N \quad \text{such that} \quad n \geq N \implies |S_n - L| < \varepsilon.$$

- Since $\beta < L \iff L - \beta > 0$, we can set $\varepsilon := L - \beta$ to produce an N such that

$$n \geq N \implies |L - S_n| < L - \beta \iff \beta - L < S_n - L < L - \beta.$$

- Just taking the first part of this composite inequality we have

$$n \geq N \implies \beta - L < S_n - L \iff \beta < S_n := \inf_{k \geq n} x_k \leq x_n,$$

supplying the N for which $n \geq N \implies \beta < x_n$ as desired. ■