Title

D. Zack Garza

Friday 25th September, 2020

Contents

1 Friday, September 25

1

1 | Friday, September 25

• For X, Y topological spaces, consider

$$Y^X = \text{hom}_{\text{Top}}(X, Y) := \left\{ f : X \to Y \mid f \text{ is continuous} \right\}.$$

- Topologize with the *compact-open* topology: $U \in \text{hom}_T(X, X)$ open iff for every $f \in U$, f(K) is open for every compact $K \subseteq X$.
 - * If Y = (Y, d) is a metric space, this is the topology of "uniform convergence on compact sets": for $f_n \to f$ in this topology iff

$$||f_n - f||_{\infty,K} := \sup \left\{ d(f_n(x), f(x)) \mid x \in K \right\} \stackrel{n \to \infty}{\to} 0 \quad \forall K \subseteq X \text{ compact.}$$

In words: $f_n \to f$ uniformly on every compact set.

- Can be used to topologize a lot of interesting spaces:

$$X = I := [0, 1] \leadsto \quad P(X; x_0) := \left\{ f : I \to X \mid f(0) = x_0 \right\}$$
$$X = S^1 \leadsto \quad \Omega(X; x_0) = \mathcal{L}(X; x_0) := \left\{ f : S^1 \to X \right\}.$$

Since these are homeomorphisms, everything is invertible, so equip with function composition to form a group.

_