

# Homework 7

D. Zack Garza

November 5, 2019

## Contents

<b>1</b>	<b>Problem 1</b>	<b>1</b>
1.1	Part 1 . . . . .	1
1.2	Part 2 . . . . .	2
<b>2</b>	<b>Problem 2</b>	<b>3</b>
2.1	Part 1 . . . . .	3
2.2	Part 2 . . . . .	3
<b>3</b>	<b>Problem 3</b>	<b>3</b>
3.1	Part 1 . . . . .	3
3.2	Part 2 . . . . .	4
3.3	Part 3 . . . . .	5
<b>4</b>	<b>Problem 4</b>	<b>6</b>

## 1 Problem 1

### 1.1 Part 1

In order for  $IS$  to be a submodule of  $A$ , we need to show the following implication:

$$x \in IS, a \in A \implies xa, ax \in IS.$$

Suppose  $x \in IS$ . Then by definition,  $x = \sum_{i=1}^n r_i a_i$  for some  $r_i \in R, a_i \in A$ .

But then

$$\begin{aligned} xa &= \left( \sum_{i=1}^n r_i a_i \right) a \\ &= \sum_{i=1}^n r_i a_i a \\ &:= \sum_{i=1}^n r_i a'_i, \end{aligned}$$

where  $a'_i := a_i a$  for each  $i$ , which is still an element of  $A$  since  $A$  itself is a module and thus closed under multiplication.

But this expresses  $xa$  as an element of  $IS$ . Similarly, we have

$$\begin{aligned} ax &= a \left( \sum_{i=1}^n r_i a_i \right) \\ &= \sum_{i=1}^n a r_i a_i a \\ &:= \sum_{i=1}^n r_i a a_i, \\ &:= \sum_{i=1}^n r_i a'_i, \end{aligned}$$

and so  $ax \in IS$  as well.

## 1.2 Part 2

Letting  $R/I \curvearrowright A/IA$  be the action given by  $r + I \curvearrowright +IA := ra + IA$ , we need to show the following:

- $r.(x + y) = r.x + r.y$ ,
- $(r + r').x = r.x + r'.x$ ,
- $(rs).x = r.(s.x)$ , and
- $1.x = x$ .

Letting  $\oplus$  denote the addition defined on cosets, we have

$$\begin{aligned} r \curvearrowright (x + IA \oplus y + IA) &:= r \curvearrowright x + y + IA \\ &:= r(x + y) + IA \\ &= rx + ry + IA \\ &:= rx + IA \oplus ry + IA \\ &:= (r \curvearrowright x + IA) \oplus (r \curvearrowright y + IA). \end{aligned}$$

$$\begin{aligned} (r + s) \curvearrowright x + IA &:= (r + s)x + IA \\ &:= rx + sx + IA \\ &:= rx + IA \oplus sx + IA \\ &:= (rs \curvearrowright IA) \oplus (sx \curvearrowright IA). \end{aligned}$$

$$\begin{aligned}
(rs) \curvearrowright x + IA &:= rsx + IA \\
&= r(sx) + IA \\
&:= r \curvearrowright (sx + IA) \\
&= r \curvearrowright (s \curvearrowright x + IA).
\end{aligned}$$

$$1 \curvearrowright x + IA := 1x + IA = x + IA.$$

## 2 Problem 2

### 2.1 Part 1

We want to show that every simple  $R$ -module  $M$  is cyclic, i.e. if the only ideals of  $M$  are  $(0)$  and  $M$  itself, that  $M = \langle m \rangle$  for some element  $m \in M$ .

Towards a contradiction, let  $M$  be a simple  $R$ -module and suppose  $M$  is not cyclic, so  $M \neq \langle m \rangle$  for any  $m \in M$ . But then let  $a \in M$  be an arbitrary nontrivial element; then  $(a)$  is a non-empty ideal (since it contains  $a$ ), so  $(a) \neq 0$ . Since  $M$  is simple, we must have  $(a) = M$ , a contradiction.

### 2.2 Part 2

Let  $\phi : A \rightarrow A$  be a module endomorphism on a simple module  $A$ . Then  $\text{im } \phi := \phi(A)$  is a submodule of  $A$ . Since  $A$  is simple, we have either  $\text{im } \phi = 0$ , in which case  $\phi$  is the zero map, or  $\text{im } \phi = A$ , so  $\phi$  is surjective. In this case, we can also consider  $\ker \phi$ , which is a submodule of  $A$ . Since  $A$  is simple, we can again only have  $\ker \phi = A$ , which can not happen if  $\phi$  is not the zero map, or  $\ker \phi = 0$ , in which case  $\phi$  is both a surjective and an injective map and thus an isomorphism of modules.

## 3 Problem 3

### 3.1 Part 1

We want to show that if  $A, B$  are  $R$ -modules then  $X = (\text{hom}_{R\text{-mod}}(A, B), +)$  is an abelian group. Let  $f, g, h \in X$ , we then need to show the following:

- a. Closure:  $f + g \in X$
- b. Associativity:  $f + (g + h) = (f + g) + h$
- c. Identity:  $\text{id} \in X$
- d. Inverses:  $f^{-1} \in X$
- e. Commutativity:  $f + g = g + f$

Closure: This follows from the definition, because  $(f + g) \curvearrowright x := f(x) + g(x)$  pointwise, which is well-defined homomorphism  $A \rightarrow B$ .

Associativity: We have

$$\begin{aligned}
f + (g + h) \curvearrowright x &:= f(x) + (g + h)(x) \\
&:= f(x) + (g(x) + h(x)) \\
&= (f(x) + g(x)) + h(x) \\
&= (f + g) + h \curvearrowright x.
\end{aligned}$$

Identity: We can define  $\mathbf{0} : A \rightarrow B$  by  $\mathbf{0}(x) = 0 \in B$ . Then

$$(f + \mathbf{0}) \curvearrowright x = f(x) + 0 = f(x) = 0 + f(x) = (\mathbf{0} + f) \curvearrowright x.$$

Inverses: Given  $f \in X$ , we can define  $-f : A \rightarrow B$  as  $-f(x) = -x$ . Then

$$\begin{aligned}
(f + -f) \curvearrowright x &= f(x) + -f(x) = f(x) - f(x) = x - x = 0 = \mathbf{0} \curvearrowright x \\
(-f + f) \curvearrowright x &= -f(x) + f(x) = -f(x) + f(x) = -x + x = 0 = \mathbf{0} \curvearrowright x.
\end{aligned}$$

Commutativity: Since  $B$  is a module, by definition  $(B, +)$  is an abelian group. Thus

$$(f + g) \curvearrowright x = f(x) + g(x) = g(x) + f(x) = (g + f) \curvearrowright x.$$

### 3.2 Part 2

By part 1,  $(\text{hom}_{R\text{-mod}}(A, A), +)$  is an abelian group, We just need to check that  $(\text{hom}_R(A, A), \circ)$  is a monoid, i.e.:

- Associativity:  $f \circ (g \circ h) = (f \circ g) \circ h$
- Identity:  $\text{id} \circ f = f$
- Closure:  $f \circ g \in \text{hom}_{R\text{-mod}}(A, A)$

Associativity: We have

$$\begin{aligned}
f \circ (g \circ h) \curvearrowright x &:= (f \circ (g \circ h))(x) \\
&= f((g \circ h)(x)) \\
&= f(g(h(x))) \\
&= (f \circ g)(h(x)) \\
&= ((f \circ g) \circ h)(x) \\
&:= (f \circ g) \circ h \curvearrowright x.
\end{aligned}$$

Identity: Take  $\text{id}_A : A \rightarrow A$  given by  $\text{id}_A(x) = x$ , then

$$f \circ \text{id}_A \curvearrowright x = f(\text{id}_A(x)) = f(x) = \text{id}_A(f(x)) = \text{id}_A \circ f \curvearrowright x.$$

Closure: If  $f : A \rightarrow A$  and  $g : A \rightarrow A$  are homomorphisms, then  $f \circ g : A \rightarrow A$  as a set map, and is an  $R$ -module homomorphism because

$$\begin{aligned} f \circ g \curvearrowright (r + s)(x + y) &= f(g((r + s)(x + y))) \\ &= f((r + s)(g(x) + g(y))) \\ &= (r + s)(f(g(x)) + f(g(y))) \\ &= (f \curvearrowright (r + s)(x + y)) \circ (g \curvearrowright (r + s)(x + y)). \end{aligned}$$

### 3.3 Part 3

For arbitrary  $x, y \in A$ , we need to check the following:

- a.  $f \curvearrowright (x + y) = f \curvearrowright x + f \curvearrowright y$
- b.  $(f + g) \curvearrowright x = f \curvearrowright x + g \curvearrowright x$
- c.  $f \circ g \curvearrowright x = f \curvearrowright (g \curvearrowright x)$
- d.  $\text{id}_A \curvearrowright x = x$

For (a):

$$\begin{aligned} f \curvearrowright (x + y) &:= f(x + y) \\ &= f(x) + f(y) \quad \text{since } f \text{ is a homomorphism} \\ &= f \curvearrowright x + f \curvearrowright y \end{aligned}$$

For (b):

$$\begin{aligned} (f + g) \curvearrowright x &= (f + g)(x) \\ &= f(x) + g(x) \\ &= f \curvearrowright x + g \curvearrowright x. \end{aligned}$$

For (c):

$$\begin{aligned} f \circ g \curvearrowright x &= (f \circ g)(x) \\ &= f(g(x)) \\ &= f \curvearrowright g(x) \\ &= f \curvearrowright (g \curvearrowright x). \end{aligned}$$

For (d):

$$\text{id}_A \curvearrowright x = \text{id}_A(x) = x.$$

## 4 Problem 4

We have the following situation:

$$\begin{array}{ccccccc}
 A_1 & \longrightarrow & A_2 & \longrightarrow & A_3 & \longrightarrow & A_4 \\
 \downarrow \alpha_1 & & \downarrow \alpha_2 & & \downarrow f & & \downarrow \alpha_4 \\
 B_1 & \longrightarrow & B_2 & \longrightarrow & B_3 & \longrightarrow & B_4
 \end{array}$$