

Lie Algebras

D. Zack Garza

October 21, 2019

Contents

1	Lecture 1	2
1.1	Overview	3
1.1.1	Chapter 2	3
1.1.2	Chapter 3-4	3
1.2	Classification	3
1.3	Chapters 4-5	4
1.3.1	Chapter 6	4
1.3.2	Chapter 7	4
1.3.3	Topics	4
1.4	Content	4
1.5	Linear Lie Algebras	5
2	Lecture 2	7
2.1	Lie Algebras of Derivations	7
2.2	Abstract Lie Algebras	8
2.3	Ideals	9
3	Lecture 3	9
3.1	Homomorphisms and Representations	10
3.2	Automorphisms	10
4	Lecture 4	11
4.1	Solvability	11
4.2	Nilpotency	12
5	Lecture 5	13
5.1	Chapter 2: Semisimple Lie Algebras	14
6	Lecture 6	14
6.1	4.1: Lie's Theorem	15
6.2	4.2: Jordan-Chevalley Decomposition	16
7	Lecture n+2	17
8	Lecture n+3	21

9 Lecture n+4	21
9.1 Cartan's Criteria	23
9.2 Section 5.2	24
10 Lecture 10?	24
10.1 Inner Derivations	26
11 Wednesday September 4th	27
11.1 4.3: Cartan's Criterion	27
11.2 Killing Form	29
11.2.1 Criterion for Semisimplicity	29
12 Wednesday September 11th	30
13 Friday September 13th	30
14 Monday September 16th	30
14.1 Proof of Weyl's(?) Theorem	30
14.1.1 Step 1:	32
14.1.2 Step 2:	32
14.1.3 Step 3:	32
14.1.4 Step 4: The general case	32
14.2 Preservation of Jordan Decomposition	34
15 Wednesday Lecture	34
15.1 Finite Dimensional Representations of $\mathfrak{sl}(2, \mathbb{C})$	35
15.2 Weights and Maximal Vectors	35
16 Friday Lecture	38
16.1 Root Space Decomposition	38
16.1.1 Maximal Toral subalgebra and roots	38
17 Monday Lecture	40
17.1 The Centralizer of \mathfrak{h}	40
17.2 8.3: Orthogonality Properties	42
18 Lecture Wednesday	42
18.1 Orthogonality Properties	42
18.2 Orthogonality/Integrality Properties	44
19 Lecture (Friday)	45

1 Lecture 1

The material for this class will roughly come from Humphrey, Chapters 1 to 5. There is also a useful appendix which has been uploaded to the ELC system online.

1.1 Overview


Here is a short overview of the topics we expect to cover:

1.1.1 Chapter 2

- Ideals, solvability, and nilpotency
- Semisimple Lie algebras
 - These have a particularly nice structure and representation theory
- Determining if a Lie algebra is semisimple using Killing forms
- Weyl's theorem for complete reducibility for finite dimensional representations
- Root space decompositions


1.1.2 Chapter 3-4


We will describe the following series of correspondences:



1.2 Classification

The classical Lie algebras can be essentially classified by certain classes of diagrams:





1.3 Chapters 4-5

These cover the following topics:

- Conjugacy classes of Cartan subalgebras
- The PBW theorem for the universal enveloping algebra
- Serre relations

1.3.1 Chapter 6

Some important topics include:

- Weight space decompositions
- Finite dimensional modules
- Character and the Harish-Chandra theorem
- The Weyl character formula
 - This will be computed for the specific Lie algebras seen earlier

We will also see the type A_ℓ algebra used for the first time; however, it differs from the other types in several important/significant ways.

1.3.2 Chapter 7

Skip!

1.3.3 Topics

Time permitting, we may also cover the following extra topics:

- Infinite dimensional Lie algebras [Carter 05]
- BGG Cat- \mathcal{O} [Humphrey 08]

1.4 Content

Fix F a field of characteristic zero – note that prime characteristic is closer to a research topic.

Definition 1. A **Lie Algebra** \mathfrak{g} over F is an F -vector space with an operation denoted the Lie bracket,

$$\begin{aligned} [\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} &\rightarrow \mathfrak{g} \\ (x, y) &\mapsto [x, y]. \end{aligned}$$

satisfying the following properties:

- $[\cdot, \cdot]$ is bilinear
- $[x, x] = 0$
- The Jacobi identity:

$$[x, [y, z]] + [y, [x, z]] + [z, [x, y]] = 0.$$

Exercise 1. Show that $[x, y] = -[y, x]$.

Definition 2. Two Lie algebras $\mathfrak{g}, \mathfrak{g}'$ are said to be isomorphic if $\varphi([x, y]) = [\varphi(x), \varphi(y)]$.

1.5 Linear Lie Algebras

Let $V = \mathbb{F}^n$, and define $\text{End}(V) = \{f : V \rightarrow V \mid f \text{ is linear}\}$. We can then define $\mathfrak{gl}(n, V)$ by setting $[x, y] = (x \circ y) - (y \circ x)$.

Exercise 2. Verify that V is a Lie algebra.

Definition 3. Define

$$\mathfrak{sl}(n, V) = \{f \in \mathfrak{gl}(n, V) \mid \text{Tr}(f) = 0\}.$$

(Note the different in definition compared to the lie group $\text{SL}(n, V)$.)

Definition 4. A *subalgebra* of a Lie algebra is a vector subspace that is closed under the bracket.

Definition 5. The symplectic algebra

$$\mathfrak{sp}(2\ell, F) = \left\{ A \in \mathfrak{gl}(2\ell, F) \mid MA - A^T M = 0 \right\} \text{ where } M = \left(\begin{array}{c|c} 0 & I_n \\ \hline -I_n & 0 \end{array} \right).$$


Definition 6. The orthogonal algebra

$$\mathfrak{so}(2\ell, F) = \left\{ A \in \mathfrak{gl}(2\ell, F) \mid MA - A^T M = 0 \right\} \text{ where}$$

$$M = \begin{cases} \left(\begin{array}{c|cc} 1 & 0 & \\ \hline 0 & 0 & I_n \\ & -I_n & 0 \end{array} \right) & n = 2\ell + 1 \text{ odd,} \\ \left(\begin{array}{c|c} 0 & I_n \\ \hline -I_n & 0 \end{array} \right) & \text{else.} \end{cases}$$

Proposition 7. The dimensions of these algebras can be computed;

- The dimension of $\mathfrak{gl}(n, \mathbb{F})$ is n^2 , and has basis $\{e_{i,j}\}$ the matrices if a 1 in the i, j position and



zero elsewhere.

- For type A_ℓ , we have $\dim \mathfrak{sl}(n, \mathbb{F}) = (\ell + 1)^2 - 1$.
- For type C_ℓ , we have $\dim \mathfrak{sp}(n, \mathbb{F}) = \ell^2 + 2 \left(\frac{\ell(\ell+1)}{2} \right)$, and so elements here

$$\begin{pmatrix} A & B = B^t \\ C = C^t & A^t \end{pmatrix}.$$

- For type D_ℓ we have


$$\dim \mathfrak{so}(2\ell, \mathbb{F}) = \dim \left\{ \begin{pmatrix} A & B = -B^t \\ C = -C^t & -A^t \end{pmatrix} \right\},$$

which turns out to be $2\ell^2 - \ell$.

- For type B_ℓ , we have $\dim \mathfrak{so}(2\ell, \mathbb{F}) = 2\ell^2 - \ell + 2\ell = 2\ell^2 + \ell$, with elements of the form

$$\left(\begin{array}{c|cc} 0 & M & N \\ \hline -N^t & A & C = C^t \\ -M^t & B = B^t & -A^t \end{array} \right).$$

Exercise 3. Use the relation $MA = A^{tM}$ to reduce restrictions on the blocks.



Theorem 8. These are *all* of the isomorphisms between any of these types of algebras, in any dimension.

2 Lecture 2

Recall from last time that a Lie Algebra is a vector space with a bilinear bracket, which importantly satisfies the Jacobi identity:

$$[x, [y, z]] + [y, [x, z]] + [z, [x, y]] = \mathbf{0}.$$

Also recall the examples from last time:

- $A_\ell \iff \mathfrak{sl}(\ell + 1, F)$
- $B_\ell \iff \mathfrak{so}(2\ell + 1, F)$
- $C_\ell \iff \mathfrak{sp}(2\ell, F)$
- $D_\ell \iff \mathfrak{so}(2\ell, F)$

Exercise 4. Characterize these matrix subalgebras in terms of basis elements, and compute their dimensions.

2.1 Lie Algebras of Derivations

Definition 9. An F -algebra A is an F -vector space endowed with a bilinear map $A^2 \rightarrow A$, $(x, y) \mapsto xy$.

Definition 10. An algebra is **associative** if $x(yz) = (xy)z$.

Modern interest: simple Lie algebras, which have a good representation theory. Take a look at Erdmann-Wildon (Springer) for an introductory look at 3-dimensional algebras.

Definition 11. Any map $\delta : A^2 \rightarrow A$ that satisfies the Leibniz rule is called a **derivation** of A , where the rule is given by $\delta(xy) = \delta(x)y + x\delta(y)$.

Definition 12. We define $\text{Der}(A) = \{\delta \mid \delta \text{ is a derivation}\}$.

Any Lie algebra \mathfrak{g} is an F -algebra, since $[\cdot, \cdot]$ is bilinear. Moreover, \mathfrak{g} is associative iff $[x, [y, z]] = 0$.

Exercise 5. Show that $\text{Der } \mathfrak{g} \leq \mathfrak{gl}(\mathfrak{g})$ is a Lie subalgebra. One needs to check that $\delta_1, \delta_2 \in \mathfrak{g} \implies [\delta_1, \delta_2] \in \mathfrak{g}$.

Exercise 6 (Turn in). Define the adjoint by $\text{ad } x : \mathfrak{g} \rightarrow \mathfrak{g}, y \mapsto [x, y]$. Show that $\text{ad } x \in \text{Der}(\mathfrak{g})$.

2.2 Abstract Lie Algebras

Fact: Every finite-dimensional Lie algebra is isomorphic to a linear Lie algebra, i.e. a subalgebra of $\mathfrak{gl}(V)$. Each isomorphism type can be specified by certain *structure constants* for the Lie bracket.

Example 13. Any F -vector space can be made into a Lie algebra by setting $[x, y] = 0$; such algebras are referred to as *abelian*.

Attempting to classify Lie algebras of dimension at most 2.

- 1 dimensional: We can write $\mathfrak{g} = Fx$, and so $[x, x] = 0 \implies [\cdot, \cdot] = 0$. So every bracket must be zero, and thus every Lie algebra is abelian.
- 2 dimensional: Write $\mathfrak{g} = Fx \oplus Fy$, the only nontrivial bracket here is $[x, y]$. Some cases:
 - $[x, y] = 0 \implies \mathfrak{g}$ is abelian.
 - $[x, y] = ax + by \neq 0$. Assume $a \neq 0$ and set $x' = ax + by, y' = \frac{y}{a}$. Now compute $[x', y'] = [ax + by, \frac{y}{a}] = [x, y] = ax + by = x'$. Punchline: $\mathfrak{g} \cong Fx' \oplus Fy', [x', y'] = x'$.

We can fill in a table with all of the various combinations of brackets:

$[\cdot, \cdot]$	x'	y'
x'	0	x'
y'	$-x'$	0

Example 14. Let $V = \mathbb{R}^3$, and define $[a, b] = a \times b$ to be the usual cross product.

Exercise 7. Look at notes for basis elements of $\mathfrak{sl}(2, F)$,

$$e = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix},$$

$$h = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix},$$

$$f = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}.$$

Compute the matrices of $\text{ad } (e), \text{ad } (h), \text{ad } (g)$ with respect to this basis.

2.3 Ideals

Definition 15. A subspace $I \subseteq \mathfrak{g}$ is called an **ideal**, and we write $I \trianglelefteq \mathfrak{g}$, if $x, y \in I \implies [x, y] \in I$.

Note that there is no need to distinguish right, left, or two-sided ideals. This can be shown using $[x, y] = [-y, x]$.

Exercise 8. Check that the following are all ideals of \mathfrak{g} :

- $\{0\}, \mathfrak{g}$.
- $\mathfrak{z}(\mathfrak{g}) = \{z \in \mathfrak{g} \mid [x, z] = 0 \quad \forall x \in \mathfrak{g}\}$
- The commutator (or derived) algebra $[\mathfrak{g}, \mathfrak{g}] = \{\sum_i [x_i, y_i] \mid x_i, y_i \in \mathfrak{g}\}$.
– Moreover, $[\mathfrak{gl}(n, F), \mathfrak{gl}(n, F)] = \mathfrak{sl}(n, F)$.

Fact: If $I, J \trianglelefteq \mathfrak{g}$, then

- $I + J = \{x + y \mid x \in I, y \in J\} \trianglelefteq \mathfrak{g}$
- $I \cap J \trianglelefteq \mathfrak{g}$
- $[I, J] = \{\sum_i [x_i, y_i] \mid x_i \in I, y_i \in J\} \trianglelefteq \mathfrak{g}$

Definition 16. A Lie algebra is **simple** if $[\mathfrak{g}, \mathfrak{g}] \neq 0$ (i.e. when \mathfrak{g} is not abelian) and has no non-trivial ideals. Note that this implies that $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$.

Theorem 17. Suppose that $\text{char } F \neq 2$, then $\mathfrak{sl}(2, F)$ is not simple.

Proof. Recall that we have a basis of $\mathfrak{sl}(2, F)$ given by $B = \{e, h, f\}$ where

- $[e, f] = h,$
- $[h, e] = 2e,$
- $[h, f] = -2f.$

So think of $[h, e] = \text{ad } h$, so h is an eigenvector of this map with eigenvalues $\{0, \pm 2\}$. Since $\text{char } F \neq 2$, these are all distinct. Suppose $\mathfrak{sl}(2, F)$ has a nontrivial ideal I ; then pick $x = ae + bh + cf \in I$. Then $[e, x] = 0 - 2be + ch$, and $[e, [e, x]] = 0 - 0 + 2ce$. Again since $\text{char } F \neq 2$, then if $c \neq 0$ then $e \in I$. Now you can show that $h \in I$ and $f \in I$, but then $I = \mathfrak{sl}(2, F)$, a contradiction. So $c = 0$.

Then $x = bh \neq 0$, so $h \in I$, and we can compute

$$\begin{aligned} 2e &= [h, e] \in I \implies e \in I, \\ 2f &= [h, -f] \in I \implies f \in I. \end{aligned}$$

which implies that $I = \mathfrak{sl}(2, F)$ and thus it is simple. □

Note that there is a homework coming due next Monday, about 4 questions.

3 Lecture 3

Last time, we looked at ideals such as $0, \mathfrak{g}, Z(\mathfrak{g})$, and $[\mathfrak{g}, \mathfrak{g}]$.

Definition: If $I \trianglelefteq \mathfrak{g}$ is an ideal, then the quotient \mathfrak{g}/I also yields a Lie algebra with the bracket given by $[x + I, y + I] = [x, y] + I$.

Exercise: Check that this is well-defined, so that if $x + I = x' + I$ and $y + I = y' + I$ then $[x, y] + I = [x', y'] + I$.

3.1 Homomorphisms and Representations

Definition 18. A linear map $\phi : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ is a *Lie homomorphism* if $\phi[x, y] = [\phi(x), \phi(y)]$.

Remark. $\ker \phi \trianglelefteq \mathfrak{g}_1$ and $\text{im } \phi \leq \mathfrak{g}_2$ is a subalgebra.

Fact: There is a canonical way to set up a 1-to-1 correspondence $\{I \trianglelefteq \mathfrak{g}\} \iff \{\text{hom } \phi : \mathfrak{g} \rightarrow \mathfrak{g}'\}$ where $I \mapsto (x \mapsto x + I)$ and the inverse is given by $\phi \mapsto \ker \phi$.

Theorem (Isomorphism theorem for Lie algebras):

- If $\phi : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ is a Lie algebra homomorphism, then $\mathfrak{g}/\ker \phi \cong \text{im } \phi$
- If $I, J \trianglelefteq \mathfrak{g}$ are ideals and $I \subset J$ then $J/I \trianglelefteq \mathfrak{g}/I$ and $(\mathfrak{g}/I)/(J/I) \cong \mathfrak{g}/J$.
- If $I, J \trianglelefteq \mathfrak{g}$ then $(I + J)/J \cong I/(I \cap J)$.

Definition: A *representation* of a Lie algebra \mathfrak{g} is a Lie algebra homomorphism $\phi : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ into a linear Lie algebra for some vector space V .

We call V a \mathfrak{g} -module with action $g \cdot v = \phi(g)(v)$.

Example: The *adjoint representation*:

$$\begin{aligned} \text{ad} : \mathfrak{g} &\rightarrow \mathfrak{gl}(\mathfrak{g}) \\ x &\mapsto [x, \cdot]. \end{aligned}$$

Corollary 19. Any simple Lie algebra is isomorphic to a linear Lie algebra.

Proof: Since \mathfrak{g} is simple, the center $Z(\mathfrak{g}) = 0$. We can rewrite the center as

$$\begin{aligned} Z(\mathfrak{g}) &= \{x \in \mathfrak{g} \mid \text{ad } x(y) = 0 \quad \forall y \in \mathfrak{g}\} \\ &= \ker \text{ad } x. \end{aligned}$$

Using the first isomorphism theorem, we have $\mathfrak{g}/Z(\mathfrak{g}) \cong \text{im ad} \subseteq \mathfrak{gl}(\mathfrak{g})$. But $\mathfrak{g}/Z(\mathfrak{g}) = \mathfrak{g}$ here, so we are done.

3.2 Automorphisms

Definition: An automorphism of \mathfrak{g} is an isomorphism $\mathfrak{g} \rightarrow \mathfrak{g}$, and we define

$$\text{Aut}(\mathfrak{g}) = \{\phi : \mathfrak{g} \rightarrow \mathfrak{g} \mid \phi \text{ is an isomorphism}\}.$$

Proposition: If $\delta \in \text{Der}(\mathfrak{g})$ is nilpotent, then

$$\exp(\delta) := \sum \frac{\delta^n}{n!} \in \text{Aut}(\mathfrak{g}).$$

This is well-defined because δ is nilpotent, and a binomial formula holds:

$$\frac{\delta^n([x, y])}{n!} = \sum_{i=0}^n \left[\frac{\delta^i(x)}{i!}, \frac{\delta^{n-i}(y)}{(n-i)!} \right].$$

and for $n = 1$, $\delta([x, y]) = [x, \delta(y)] + [\delta(x), y]$.

Exercise: Show that

$$[(\exp \delta)(x), (\exp \delta)(y)] = \sum_{n=0}^{k-1} \frac{\delta^n([x, y])}{n!}.$$

Example: Let $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{F})$ and define

$$s = \exp(\text{ad } e) \exp(\text{ad } -f) \exp(\text{ad } e) \in \text{Aut } \mathfrak{g}.$$

where e, f are defined as (todo, see written notes).

Then define the Weyl group $W = \langle s \rangle$.

Exercise: Check that $s(e) = -f, s(f) = -e, s(h) = -h$, and so the order of s is 2 and $W = \{1, s\}$.

4 Lecture 4

4.1 Solvability

Idea: Define a semisimple Lie algebra

Definition: The derived series for \mathfrak{g} is given by

$$\begin{aligned} \mathfrak{g}^{(0)} &= \mathfrak{g} \\ \mathfrak{g}^{(1)} &= [\mathfrak{g}^{(0)}, \mathfrak{g}^{(0)}] \\ &\dots \\ \mathfrak{g}^{(i+1)} &= [\mathfrak{g}^{(i)}, \mathfrak{g}^{(i)}]. \end{aligned}$$

The Lie algebra \mathfrak{g} is *solvable* if there is some n for which $\mathfrak{g}^{(n)} = 0$.

Exercise (to turn in): Check that the Lie algebra of upper triangular matrices in $\mathfrak{gl}(n, \mathbb{F})$.

Example: Abelian Lie algebras are solvable

Example: Simple Lie algebras are *not* solvable.

Proposition: Let \mathfrak{g} be a Lie algebra, then

1. If \mathfrak{g} is solvable, then all subalgebras and all homomorphic images of \mathfrak{g} are also solvable.
2. If $I \trianglelefteq \mathfrak{g}$ and both I and \mathfrak{g}/I are solvable, then so is \mathfrak{g} .
3. If $I, J \trianglelefteq \mathfrak{g}$ are solvable, then so is $I + J$.

Corollary (of part 3 above): Any Lie algebra has a unique maximal solvable ideal, which we denote the *radical* $\text{Rad}(\mathfrak{g})$.

Definition: A Lie algebra is semisimple if $\text{Rad}(\mathfrak{g}) = 0$.

Example: Any simple Lie algebra is semisimple.

Example: Using part (2) above, we can deduce that we can construct a semisimple Lie algebra from *any* Lie algebra: for any \mathfrak{g} , the quotient $\mathfrak{g}/\text{Rad}(\mathfrak{g})$ is semisimple.

4.2 Nilpotency

$$\begin{aligned}\mathfrak{g}^0 &= \mathfrak{g} \\ \mathfrak{g}^1 &= [\mathfrak{g}^0, \mathfrak{g}^0] \\ &\quad \dots \\ \mathfrak{g}^{i+1} &= [\mathfrak{g}^i, \mathfrak{g}^i].\end{aligned}$$

Much like the previous case, we have

Example: Abelian Lie algebras are nilpotent.

Example: Nilpotent Lie algebras are solvable.

Example: The *strictly* upper triangular matrices (with zero on the diagonal) are nilpotent.

1. If \mathfrak{g} is nilpotent, then all subalgebras and all homomorphic images of \mathfrak{g} are also nilpotent.
2. If $\mathfrak{g}/Z(\mathfrak{g})$ is nilpotent, then so is \mathfrak{g} .
3. If $\mathfrak{g} \neq 0$ is nilpotent, then $Z(\mathfrak{g}) \neq 0$.

Claim: If \mathfrak{g} is nilpotent, then $\text{ad } x \in \text{End}(\mathfrak{g})$ is nilpotent for all $x \in \mathfrak{g}$.

Proof: This is because $\mathfrak{g}^n = 0 \iff [\mathfrak{g}, [\mathfrak{g}, [\mathfrak{g}, \dots]]] = 0$, and so for every $x_i, y \in \mathfrak{g}$ we have $[x_1, [x_2, \dots [x_n, y]]] = 0$, and so $\text{ad } x_1 \circ \text{ad } x_2 \circ \dots \circ \text{ad } x_n = 0$ which implies that $\text{ad } x^n = 0$ for all $x \in \mathfrak{g}$.

Theorem [Engel]: If $\text{ad } x$ is nilpotent for all $x \in \mathfrak{g}$, then \mathfrak{g} is nilpotent.

Remark: This can be confusing if \mathfrak{g} is a linear algebra, we can consider elements $x \in \mathfrak{g}$ and ask if it is the case x being nilpotent (as an endomorphism) iff $\mathfrak{g}x$ is nilpotent? False, a counterexample is $\mathfrak{g} = \mathfrak{gl}(2, \mathbb{C})$, where there exists an x which is *not* nilpotent while $\text{ad } x$ is nilpotent, which contradicts the above theorem.

Proof:

Lemma: Let $\mathfrak{g} \subseteq \mathfrak{gl}(V)$ be a Lie subalgebra for some finite dimensional vector space V . If x is nilpotent as an endomorphism on V for all $x \in \mathfrak{g}$, then there exists a nonzero vector $v \in V$ such that $\mathfrak{g}v = 0$, so $x \in \mathfrak{g} \implies x(v) = 0$.

Proof of lemma Use induction on $\dim \mathfrak{g}$, splitting into two separate base cases: - Case $\dim \mathfrak{g} = 0$, then $\mathfrak{g} = \{0\}$. - Case $\dim \mathfrak{g} = 1$, left as an exercise.

Inductive step: Let A be a maximal proper subalgebra and define $\phi : A \rightarrow \mathfrak{gl}(\mathfrak{g}/A)$ where $a \mapsto (x + A \mapsto [a, x] + A)$. We need to check that ϕ is a homomorphism, this just follows from using the Jacobi identity.

We also need to show that $\text{im } \phi \leq \mathfrak{gl}(\mathfrak{g}/A)$ is a Lie subalgebra, and $\dim \text{im } \phi < \dim \mathfrak{g}$. The claim is that $\phi(a) \in \text{End}(\mathfrak{g}/A)$ is nilpotent for all $a \in A$. By the inductive hypothesis, there is a nonzero coset $y + A \in \mathfrak{g}/A$ such that $(\text{im } \phi) \cdot (y + A) = A$. Since $y \notin A$, then $\phi(a)(y + A) = A$ for all $a \in A$, and so $[a, y] \in A$.

We want to show that A is a subalgebra of codimension 1, and $A \oplus F_y \leq \mathfrak{g}$ is a Lie subalgebra. This is because $[a_1 + c_1y, a_2 + c_2y] = [a_1, a_2] + c_2[a_1, y] - c_2[a_2, y] + c_1c_2[y, y]$. The last term is zero, the middle two terms are in A , and because A is closed under the bracket, the first term is in A as well.

But then $A \oplus F_y$ is a larger subalgebra than A , which was maximal, so it must be everything. So $A \oplus F_y = \mathfrak{g}$. So $A \leq \mathfrak{g}$ because $[a_1, a_2 + cy]$ is in A , $A \oplus F_y = \mathfrak{g}$ respectively, and this equals $[a_1, a_2] + c[a_1, y]$, where both terms are in A .

Proof to be continued on Friday!

5 Lecture 5

Last time: we had a theorem that said that if $\mathfrak{g} \in \mathfrak{gl}(V)$ and every $x \in \mathfrak{g}$ is nilpotent, then there exists a nonzero $v \in V$ such that $\mathfrak{g}v = 0$.

We proceeded by induction on the dimension of V , constructing $\text{im } \phi \subseteq \mathfrak{gl}(\mathfrak{g}/A)$, and showed that $\mathfrak{g} = A \oplus F_y$. Now consider

$$W = \{v \in V \mid Av = 0\},$$

which is \mathfrak{g} -invariant, so $\mathfrak{g}(W) \subseteq W$, or for all $a \in A, x \in \mathfrak{g}, v \in W$, we have $a \curvearrowright x(v) = 0$. This is true because $a \curvearrowright x = x \circ a + [a, x] \in \mathfrak{gl}(V)$. But V is killed by any element in A , and both of these terms are in A . In particular, the y appearing in F_y also satisfies $y \in W$. Consider $y|_W \in \text{End}(W)$, and we want to apply the inductive hypothesis to $F_y|_W \subseteq \mathfrak{gl}(W)$.

We need to check that $y|_W \in \text{End}(W)$, which is true exactly because y is nilpotent. So we can construct a nonzero $v \in W \subset V$ such that $y(v) = 0$, and so $\mathfrak{g}v = 0$.

Claim: $\phi(a) \in \text{End}(\mathfrak{g}/A)$ is nilpotent. Each $a \in A \subset \mathfrak{g}$ is nilpotent by assumption. Define the maps for left multiplication by a , $m_\ell : x \mapsto ax$, and the right multiplication $m_r : x \mapsto xa$. These are nilpotent, and since m_ℓ, m_r commute, the difference $m_\ell - m_r$ is nilpotent, and this is exactly $\text{ad } a$. But then $\phi(a)$ is nilpotent.

Good proof for using all of the definitions!

Now we can see what the consequences of having such a nonzero vector are. This theorem implies Engel's theorem, which says that if $\text{ad } x \in \text{End}(\mathfrak{g})$ is nilpotent for every $x \in \mathfrak{g}$, then \mathfrak{g} is nilpotent.

Proof: By induction on dimension. The base case is easy. For the inductive step, the previous theorem applies to $\text{ad } g \in \mathfrak{gl}(\mathfrak{g})$. So we can produce the nonzero $v \in \mathfrak{g}$ such that $\text{ad } v = 0$. Then $[x, v] = 0$ for all $x \in \mathfrak{g}$, so either $v \in Z(\mathfrak{g})$ or $Z(\mathfrak{g}) \neq 0$. In either case, $\mathfrak{g}/Z(\mathfrak{g})$ has smaller dimension. Since $\text{ad } x$ is nilpotent, so is $\text{ad } x + Z(\mathfrak{g})$, and so $\mathfrak{g}/Z(\mathfrak{g})$ is nilpotent. By an earlier proposition, since the quotient is nilpotent, so is the total space. \square

Let $\mathfrak{N}(F)$ be the subalgebra of $\mathfrak{gl}(F)$ consisting of strictly upper triangular matrices. We have a corollary: if $\mathfrak{g} \subset \mathfrak{gl}(n, F)$ is a Lie subalgebra such every $x \in \mathfrak{g}$ is nilpotent as an endomorphism of F , then the matrices of \mathfrak{g} with respect to some bases of in $\mathfrak{N}(n, F)$.

The proof is by induction on n , where the base case is easy. For the inductive step, we use the previous theorem to get a v_1 such that $x(v_1) = 0$ for all $x \in \mathfrak{g}$. Let $\bar{V} = F^n/Fv_1 \cong F^{n-1}$, and define $\phi : \mathfrak{g} \rightarrow \mathfrak{gl}(\bar{V})$ where $x \mapsto (\bar{y} \mapsto \overline{y(x)})$.

Then $\text{im } \phi \leq \mathfrak{gl}(n-1, F)$ as a subalgebra, and every $\phi(x) \in \text{End}(F^{n-1})$ is nilpotent, since x was nilpotent on the larger space. But (see notes) then x can be written as a strictly upper-triangular matrix.

5.1 Chapter 2: Semisimple Lie Algebras

We now assume $\text{char } F = 0$ and $\bar{F} = F$.

Theorem: If \mathfrak{g} is a solvable Lie subalgebra of $\mathfrak{gl}(V)$ for some finite dimensional V , then V contains a common eigenvector for a $x \in \mathfrak{g}$, i.e. a $\lambda : \mathfrak{g} \rightarrow F, x \mapsto \lambda(x)$ such that $x(v) = \lambda(x)v$ for all $x \in \mathfrak{g}$.

Proof: We will use induction on the dimension of \mathfrak{g} . For the inductive step:

Claim 1: There is an ideal $A \trianglelefteq \mathfrak{g}$ such that $\mathfrak{g} = A \oplus Fy$ for some $y \neq 0$, so A is a subalgebra of a solvable Lie algebra \mathfrak{g} and thus solvable itself. By hypothesis, we can produce a $w \in V \setminus \{0\}$, and thus a functional $\lambda : A \rightarrow F$ such that $aw = \lambda(a)w$ for all $a \in A$. So we define

$$V_\lambda = \{v \in V \mid av = \lambda(a)v \forall a \in A\}$$

where $w \in V_\lambda$.

Claim 2: $y(V_\lambda) \subseteq V_\lambda$, or $y|_{V_\lambda} \in \text{End}(V_\lambda)$.

Thus $F(y|_{V_\lambda}) \leq \mathfrak{gl}(V_\lambda)$ is a Lie algebra of dimension 1, and thus solvable. By the inductive hypothesis, we can find a $v \in V_\lambda$ and some $\mu \in F$ such that $y(v) = \mu v$. An arbitrary element $x \in \mathfrak{g}$ can be written as $x = a + cy$ for some $a \in A, c \in F$ and it acts by $x(v) = a(v) + cy(v) = \lambda(a)v + c\mu v = (\lambda(a) + c\mu)v \in V_\lambda$.

6 Lecture 6

Chapter 3: Theorems of Lie and Cartan

6.1 4.1: Lie's Theorem

Theorem: Let L be a solvable subalgebra of $\mathfrak{gl}(V)$, where V is finite-dimensional. If $V \neq 0$, then V contains a common eigenvector for all of the endomorphisms in L .

Proof: Use induction on $\dim L$. The case $\dim L = 0$ is trivial. We'll attempt to mimic the proof of Theorem 3.3. The idea is to

1. Locate an ideal of K of codimension 1,
2. Show by induction that common eigenvectors exist for k ,
3. Verify that L stabilizes a space consisting of such eigenvectors,
4. Find in that space an eigenvector for a single $z \in L$ satisfying $L = K + Fz$.

Step (1): Since L is solvable and of positive dimension, then $L \not\subseteq [L, L]$. Otherwise, if $L = [L, L]$, then $L^{(1)} = L \implies L^{(n)} = L$, which would contradict L being solvable.

Since $[L, L]$ is abelian, any subspace is automatically an ideal. So take a subspace of codimension one, then its inverse image $K \trianglelefteq L$ is an ideal satisfying $[L, L] \subseteq K$.

Step (2): Use induction to find a common eigenvector $v \in V$ for K . (K is solvable; if $K = 0$ then L is abelian of dimension 1 and any eigenvector for a basis vector of L finishes the proof.)

This means that $x \in K \implies x \curvearrowright v = \lambda(x)v$ for some $\lambda : K \rightarrow F$ a linear functional. Fix this λ , and let $W = \{w \in V \mid x \curvearrowright w = \lambda(x)w \forall x \in K\}$; note that $W \neq 0$.

Step (3): This will involve showing that L leaves W invariant. Assume for the moment that this is done, and proceed to step (4).

Step (4):

Write $L = K + Fz$. Since F is algebraically closed, we can find an eigenvector $v_0 \in W$ of z for some eigenvalue of z . Then v_0 is a common eigenvector for L , and λ can be extended to a linear function on L satisfying $x \curvearrowright v_0 = \lambda(x)v_0$ where $x \in L$.

It remains to show that L stabilizes W . Let $w \in W, x \in L$. To test whether or not $x \curvearrowright w \in W$, we take an arbitrary $y \in K$ and examine

$$yx \curvearrowright w = xy \curvearrowright w - [x, y] \curvearrowright w = \lambda(y)x \curvearrowright w - \lambda([x, y])w.$$

Note: the above equality is an important trick.

Thus we need to show that $\lambda([x, y]) = 0$. To this end, fix $w \in W, x \in L$. Let $n > 0$ be the smallest integer for which $w, x \curvearrowright w, \dots, x^n \curvearrowright w$ are all linearly *independent*. Let $W_i = \text{span}(\{w, x \curvearrowright w, \dots, x^{i-1} \curvearrowright w\})$ and set $W_0 = 0$. Then $\dim W_n = n$, and $W_{n+i} = W_n$ for all $i \geq 0$. Moreover, x maps W_n into itself. It is easy to check that each $y \in K$ is represented by an upper-triangular matrix with diagonal entries equal to $\lambda(y)$. This follows immediately from the congruence

$$yx^i \curvearrowright w = \lambda(y)x^i \curvearrowright w \pmod{W_i},$$

which can be proved by induction on i . The case $i = 0$ is trivial. For the inductive step, write

$$yx^i \curvearrowright w = yxx^{i-1} \curvearrowright w = xyx^{i-1} \curvearrowright w = [x, y]x^{i-1} \curvearrowright w$$

By induction,

$$yx^{i-1} = \lambda(y)x^{i-1} \curvearrowright w + w',$$

where $w' \in W_{i-1}$. Since x maps W_{i-1} into W_i by construction, the congruence holds for all i .

According to our description of the action of $y \in K$ on W_n , we have $\text{Tr}_{W_n}(y) = n\lambda(y)$. In particular, this is true for elements k of f of the special form $[x, y]$ where x is as it was above and y is in K .

But both x and y stabilize W_n , so $[x, y]$ acts on W_n as the commutator of two endomorphisms of W_n , and the trace is therefore zero.

We conclude that $n\lambda([x, y]) = 0$. Since $\text{char} F = 0$, this forces $\lambda([x, y]) = 0$ as required. \square


Corollary A (Lie's Theorem): Let $L \leq \mathfrak{gl}(V)$ be a solvable subalgebra where $\dim V = n < \infty$. Then L stabilizes some flag in V , i.e. the matrices of L relative to a suitable basis of V are upper triangular.

Proof: Use the above theorem, along with induction on $\dim V$. This is similar to the proof of corollary 3.3.

6.2 4.2: Jordan-Chevalley Decomposition

Fact 1:

The Jordan Canonical Form of a single endomorphism x over F algebraically closed is an expression



$$\begin{pmatrix} a & 1 & & \\ & a & \ddots & \\ & & \ddots & 1 \\ 0 & & & a \end{pmatrix}$$

of x in matrix form as a sum of blocks:

Fact 2:

Call $x \in \text{End} V$ *semisimple* if the roots of its minimal polynomial over F are all distinct. Equivalently, if F is algebraically closed, then x is semisimple iff x is diagonalizable.

Fact 3:

Two commuting semisimple endomorphisms can be simultaneously diagonalized. Therefore, their sum or difference is again semisimple.

Proposition: Let V be a finite dimensional vector space over F and $x \in \text{End} V$. Then

- There exist unique $x_s, x_n \in \text{End} V$ satisfying the conditions $x = x_s + x_n$, x_s is semisimple, x_n is nilpotent, and x_s, x_n commute.
- There exists polynomials $p(t), g(t)$ such that $x_s = p(x)$ and $x_n = g(x)$. In particular, x_s, x_n commute with any endomorphism commuting with x .
- If $A < B < V$ are subspaces and x maps B into A , then x_s, x_n also map B into A .

The decomposition $x = x_s + x_n$ is called the (additive) **Jordan-Chevalley decomposition** of x , or just the Jordan decomposition. x_s, x_n are respectively called the **semisimple part** and the **nilpotent part** of x .

Example:

$$x = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \implies x_s = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad x_n = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Note that $x_s x_n = x_n = x_n x_s$, $x_s = 2x - x^2$, and $x_n = x^2 - x$. We thus have $p(t) = 2t - t^2$ and $q(t) = t^2 - t$.

7 Lecture n+2

Definition (Jordan Decomposition)

Let $X \in \text{End}(V)$ for V finite dimensional. Then,

- (a) There exists a unique $X_s, X_n \in \text{End}(V)$ such that $X = X_s + X_n$ where X_s is semisimple, X_n is nilpotent, and $[X_s, X_n] = 0$.
- (b) There exists a $p(t), q(t) \in t\mathbb{F}[t]$ such that $X_s = p(X), X_n = q(X)$.

(Polynomials with no constant term.)

Proof of (a): Assume $X_s = X_s + X_n = X'_s + X'_n$, so both have bracket zero. Assuming that (b) holds, we have $X_s = p(X)$, and so

$$[X, X_s] = [X_s + X'_n, X'_s] = [X'_s, X'_s] + [X'_s, X'_n] = 0 \implies [p(X), X'_s] = 0 = [X_s, X'_s]$$

Using fact (c) from last time, then X_s, X'_s can be diagonalized simultaneously, and so $X_s - X'_s$ is semisimple.

On the other hand, if X'_n, X_n are nilpotent, and since these commute, $X_n - X'_n$ is nilpotent. But then this is a Jordan decomposition of the zero map, i.e.

$$0 = X - X = (X_s - X'_s) + (X_n + X'_n)$$

where the first term is semisimple and the second is nilpotent. Then each term is both semisimple *and* nilpotent, so they must be zero, which is what we wanted to show.

Proof of part (b): Let $m(t) = \prod_{i=1}^r (t - \lambda_i)^{m_i}$ be the minimal polynomial of X , where each $m_i \geq 1$ and the λ_i are distinct. Then the primary composition of V is given by

$$V = \bigoplus_{i=1}^r V_i, \quad V_i = \ker(X - \lambda_i I_V)^{m_i} \neq 0, \quad X(V_i) \subseteq V_i$$

Claim: There exists a polynomial $p \in \mathbb{F}[t]$ such that

$$p(x) \sim \left(\boxed{\lambda_1 I_{v_1}} \quad \boxed{\lambda_2 I_{v_2}} \quad \boxed{\lambda_r I_{v_r}} \quad \text{circles} \right)$$

Figure 1: ???

$$\begin{aligned} p &= \lambda \pmod{(t - \lambda_i)^{m_i}} \quad \forall i, \\ p &= 0 \pmod{t}. \end{aligned}$$

The existence follows from the Chinese Remainder Theorem.

What is $p(x) \curvearrowright V_i$? This acts by scalar multiplication by λ_i for all i . (Check). Because of the restrictive conditions, $p(x)$ has no constant term.

So $p(X) = X_s$ is the semisimple part we want. Now just set $q(t) = t - p(t)$, then $X_n := q(X) = X - X_s$ is nilpotent.

Example: The Jordan Decomposition is invariant under taking adjoints.

If we have $X = X_s + X_n$, then $\text{ad } X \in \text{End}(\text{End}(V))$. It can be shown that $(\text{ad } X)_s + (\text{ad } X)_n = \text{ad } (X_s) + \text{ad } (X_n)$.

Let e_{ii} be the elementary matrix with a 1 in the i, j position. You can write $\text{ad } X$ as a 4×4 matrix (see image).

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$X = X_S + X_n$$

$$= \begin{matrix} e_{11} \\ e_{12} \\ e_{21} \\ e_{22} \end{matrix} \begin{pmatrix} 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{pmatrix}$$

$$\rightarrow \text{JNF} \begin{pmatrix} 0 & | & & & \\ \hline & 0 & 1 & & \\ & & 0 & 1 & \\ & & & \ddots & \\ & & & & 0 \end{pmatrix}$$

$$= \begin{matrix} e_{11} \\ e_{12} \\ e_{21} \\ e_{22} \end{matrix} \begin{pmatrix} 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{pmatrix}$$

$$\rightarrow \text{JNF} \begin{pmatrix} 0 & & & \\ \hline & 0 & 1 & \\ & & 0 & 1 \\ & & & \ddots & 0 \end{pmatrix}$$

You can check that $(\text{ad } X)_S = 0$, $\text{ad } (X_S) = 0$, and $(\text{ad } X)_n$ is the Jordan form given above.

Lemma:

- (a) $x \in \text{End}(V) \implies \text{ad } (x)_S = \text{ad } (x_S)$ and $\text{ad } (x)_n = \text{ad } (x_n)$.
- (b) If A is a finite dimensional \mathbb{F} -algebra, then $\delta \in \text{Der}(A) \implies \delta_S, \delta_n \in \text{Der}(A)$ as well.

Proof of (a):

Check that $\text{ad } (x) = \text{ad } (x_S) + \text{ad } (x_n)$. Then for $y \in \text{End}(V)$, we have

$$\begin{aligned} (\text{ad } (x))(y) &= [x, y] \\ &= [x_S + x_n, y] \\ &= [x_S, y] + [x_n, y] \\ &= (\text{ad } (x_S))(y) + (\text{ad } (x_n))(y). \end{aligned}$$

Using theorem 3.3, x_n nilpotent $\implies \text{ad } (x_n)$ is also nilpotent. So write $x_S = \sum \lambda_i e_{ii}$ with the eigenvalues on the diagonal. Then $\text{ad } x_S(e_{ij}) = (\lambda_i - \lambda_j)e_{ij}$ for all i, j . But then $\text{ad } x_S$ is given

$$\begin{aligned}
 & (\delta - (\lambda + \mu)I)^n([x, y]) \\
 &= \sum_{i=0}^n \binom{n}{i} \left[(\delta - \lambda I)^i(x), (\delta - \mu I)^{n-i}(y) \right]
 \end{aligned}$$

Figure 2: Image

by a matrix with $\lambda_i - \lambda_j$ in the i, j position and zeros elsewhere. By the uniqueness of the Jordan decomposition, the statement follows.

Proof of (b):

Since $\delta \in \text{Der}(A)$, the primary decomposition with respect to δ is given by

$$A = \bigoplus_{\lambda \in F} A_\lambda \quad \text{where } A_\lambda = \left\{ a \in A \mid (\delta - \lambda I)^k a = 0 \text{ for some } k \gg 0 \right\}.$$

So $\delta_s \sim A_\lambda$ by scalar multiplication (by λ). Then for $\lambda, \mu \in F$, we have

So $[A_x, A_y] \subseteq A_{\lambda+\mu}$ for all $x, y \in A$. But then

and so $\delta_s \in \text{Der}(A)$, and $\delta_n = \delta - \delta_s \in \text{Der}(A)$ as well.

8 Lecture n+3

Todo

9 Lecture n+4

Review of bilinear forms: let $V = \mathbb{F}^n$.

Definition: A bilinear form $\beta : V^2 \rightarrow \mathbb{F}$ can be represented by a matrix B with respect to a basis $\{\mathbf{v}_i\}$ such that

$$\beta\left(\sum a_i \mathbf{v}_i, \sum b_i \mathbf{v}_i\right) = (a_1 \ a_2 \ \cdots) B (b_1 \ b_2 \ \cdots)$$

- β is *symmetric* iff $\beta(a, b) = \beta(b, a)$.
- β is *symplectic* iff $\beta(a, b) = -\beta(b, a)$.
- β is *isotropic* iff $\beta(a, a) = 0$.

$$S_S([x, y])$$

||

$$(\lambda + \mu)[x, y] = [\lambda x, y] + [x, \mu y]$$

||

$$[S_S(x), y] + [x, S_S(y)]$$

Figure 3: Image

For a subspace $U \leq V$, define

$$U^\perp := \{v \in V \mid \beta(u, v) = 0 \forall u \in U\}.$$

Note: in general, left/right orthogonality are distinguished, but these will be identical when β is symmetric/symplectic.

The form β is said to be *non-degenerate* iff $V^\perp = 0$ iff $\det B \neq 0$.

Assume F is an algebraically closed field, so $\bar{F} = F$, and $\text{char} F \neq 2$, then

- If β is non-degenerate and symmetric, then $B \sim I_n$
- If β is non-degenerate and symplectic, then $B \sim [0, I_{n/2}; I_{n/2}, 0]$.

Remark:

$\mathfrak{so}(n, \mathbb{F}) = \{x \in \mathfrak{gl}(n, F) \mid \beta(x(u), v) = -\beta(u, x(v))\}$, where B has the matrix $[0, I; I, 0]$ if n is odd, or this matrix with a 1 in the top-left corner if n is even.

Similarly, $\mathfrak{sp}(2m, \mathbb{F})$ can be described this way with the matrix $[0, -I_m; -I_m, 0]$.

Overview: The Killing form is defined as $\kappa : \mathfrak{g}^2 \rightarrow \mathbb{F}$ where $\kappa(x, y) = \text{tr}(\text{ad}_x \circ \text{ad}_y)$.

Then we have **Cartan's Criteria**:

- \mathfrak{g} solvable $\iff \kappa(x, y) = 0 \forall x \in [\mathfrak{g}, \mathfrak{g}], y \in \mathfrak{g}$.
- \mathfrak{g} semisimple $\iff \kappa$ is non-degenerate.

Note that if \mathfrak{g} is semisimple, then $\mathfrak{g} = \bigoplus_i I_i$ with each $I_i \trianglelefteq \mathfrak{g}$ and simple.

9.1 Cartan's Criteria

Some facts:

1. κ is symmetric
2. If \mathfrak{g} is finite dimensional, then κ is associative, i.e $\kappa([x, y], z) = \kappa(x, [y, z])$.

Exercise: Show that if $I \trianglelefteq \mathfrak{g}$, then $I^\perp \leq \mathfrak{g}$ is an ideal.

Proof of (2): In section 4.3, it was shown that $\text{tr}([a, b] \circ c) = \text{tr}(a \circ [b, c])$ for all $a, b, c \in \text{End}(V)$ (provided V is finite dimensional).

So

$$\begin{aligned} \kappa([x, y], z) &= \text{tr}(\text{ad}_{[x, y]} \circ \text{ad}_z) \\ &= \text{tr}([\text{ad}_x, \text{ad}_y] \circ \text{ad}_z) \\ &= \text{tr}(\text{ad}_x \circ [\text{ad}_y, \text{ad}_z]) \\ &= \text{tr}(\text{ad}_x \circ \text{ad}_{[y, z]}) \\ &= \text{tr}(x, [y, z]).. \end{aligned}$$

Theorem: \mathfrak{g} is semisimple iff κ is nondegenerate.

Proof: \implies : We want to show that $\mathfrak{g}^\perp = 0$. Note that $[\mathfrak{g}^\perp, \mathfrak{g}^\perp] \subseteq \mathfrak{g}$, and so for all $x \in [\mathfrak{g}^\perp, \mathfrak{g}^\perp]$ and for any $y \in \mathfrak{g}^\perp$, we have

$$\kappa(x, y) = \text{tr}(\text{ad}_x \circ \text{ad}_y) = 0$$

by the const(?) of \mathfrak{g}^\perp . This implies \mathfrak{g}^\perp is solvable.

Using fact (2), we have $\mathfrak{g}^\perp \trianglelefteq \mathfrak{g}$ and thus $\mathfrak{g}^\perp \subseteq \text{rad}(\mathfrak{g})$, which is 0 since because \mathfrak{g} is semisimple. So either $\mathfrak{g}^\perp = 0$ or κ is nondegenerate.

Used the fact that the radical was a maximal solvable ideal.

\impliedby : We want to show that for all $I \trianglelefteq \mathfrak{g}$ where $[I, I] = 0$, we have $I^\perp \subseteq \mathfrak{g}^\perp$.

For $x \in I, y \in \mathfrak{g}$, we have

$$(\text{ad}_x \circ \text{ad}_y)^2 = \mathfrak{g} \xrightarrow{\text{ad}_y} \mathfrak{g} \xrightarrow{\text{ad}_x} I \xrightarrow{\text{ad}_y} I \xrightarrow{\text{ad}_x} 0$$

And thus $\text{tr}(\text{ad}_x \circ \text{ad}_y) = 0$ and $I \subseteq \mathfrak{g}^\perp$.

Suppose that \mathfrak{g} is *not* semisimple. Then there exists a solvable ideal $J \neq 0$ such that the last term J^i in the derived series is an ideal $I \trianglelefteq \mathfrak{g}$ such that $[I, I] = 0$, forcing $J^i \subset \mathfrak{g}^\perp = 0$, which is a contradiction.

$$\kappa_{\mathfrak{g}} \sim I_i \begin{pmatrix} \kappa_{I_i} & \\ & \end{pmatrix}$$

Figure 4: Image

9.2 Section 5.2

Theorem: If \mathfrak{g} is semisimple, then

- There exist ideals $I_i \trianglelefteq \mathfrak{g}$ which are simple Lie algebras satisfying $\mathfrak{g} = \bigoplus I_i$. Note that $[I_i, I_j] \subseteq I_i \cap I_j = 0$, since direct summands intersect only trivially.
- Every simple $I \trianglelefteq \mathfrak{g}$ is one of these I_i .
- $\kappa_{I_i} = \kappa_{\mathfrak{g}}|_{I_i \times I_i}$, so

Remark: \mathfrak{g} is semisimple $\iff \mathfrak{g} = \bigoplus_i I_i$ for some simple Lie algebras I_i .

\Leftarrow : For all $i, S := \text{rad } \mathfrak{g}, I_i \trianglelefteq I_i$ is a solvable ideal. This implies that it is 0, since I_i is simple.

By definition, simple Lie algebras are not abelian.

Supposing that $S = I_i$, we would then have $[S, S] \neq 0$ since $[I_i, I_i] \neq 0$ by definition. But $[S, S] \neq S$ because S is solvable, which says that S is not simple (a contradiction).

Note that $[\text{rad } \mathfrak{g}, \mathfrak{g}] \subseteq \bigoplus [\text{rad } \mathfrak{g}, I_i] = 0$, which forces $\text{rad } \mathfrak{g} \subseteq Z(\mathfrak{g})$. Since I_i is simple, $Z(I_i) = 0$ for all i . But $Z(\mathfrak{g}) = \bigoplus Z(I_i) = 0$, and this forces $\text{rad } (\mathfrak{g}) \subseteq Z(\mathfrak{g}) \implies \text{rad } \mathfrak{g} = 0$. So \mathfrak{g} is semisimple.

Next time – starting the representation theory with $\mathfrak{sl}(2, \mathbb{F})$.

10 Lecture 10?

Recall the killing form:




Figure 5: Image

$$\begin{aligned} \kappa : \mathfrak{lieg}^2 &\rightarrow \mathbb{F} \\ (x, y) &\mapsto \text{tr}(\text{ad}_x \circ \text{ad}_y). \end{aligned}$$

and Cartan's criteria:

1. \mathfrak{g} is solvable $\iff \kappa(x, y) = 0 \ \forall x \in \mathfrak{g}, \mathfrak{g}], \ y \in \mathfrak{g}$.
2. \mathfrak{g} is semisimple $\iff \kappa$ is non-degenerate.

Theorem: If \mathfrak{g} is semisimple, then

- a. $\mathfrak{g} = \bigoplus_{i=1}^n I_i$ for some $I_i \trianglelefteq \mathfrak{g}$ which are all simple.
- b. Every simple ideal $I \trianglelefteq \mathfrak{g}$ is one of the I_i .
- c. $\kappa_{I_i} = \kappa_{\mathfrak{g}}|_{I_i \times I_i}$.

Proof of (a): Use induction on $\dim \mathfrak{g}$. If \mathfrak{g} has no nonzero proper ideals, then \mathfrak{g} is simple and we're done.

Otherwise, let I_1 be a minimal nonzero ideal of \mathfrak{g} . Then $I_1^\perp \trianglelefteq \mathfrak{g}$ is also an ideal, and thus $I := I_1 \cap I_1^\perp \trianglelefteq \mathfrak{g}$ is as well. Then for all $x \in [I, I]$, we must have $\kappa(x, y) = 0$ for any $y \in I \subseteq I_1^\perp$. So I is solvable, and thus $I = 0$. So $\mathfrak{g} = I_1 \oplus I_1^\perp$.

$$\begin{aligned}
 \text{ad } x &\sim \left(\begin{array}{c|c} A_x & B_x \\ \hline 0 & 0 \end{array} \right) \quad \kappa_{\mathfrak{g}}(x, y) = \text{tr} \left(\left(\begin{array}{c|c} A_x & B_x \\ \hline 0 & 0 \end{array} \right) \left(\begin{array}{c|c} A_y & B_y \\ \hline 0 & 0 \end{array} \right) \right) \\
 \text{ad } y &\sim \left(\begin{array}{c|c} A_y & B_y \\ \hline 0 & 0 \end{array} \right) \quad = \text{tr} \left(\begin{array}{c|c} A_x A_y & B_x B_y \\ \hline 0 & 0 \end{array} \right) \\
 &= \text{tr}(A_x A_y) \\
 &= \chi_{\mathcal{I}_i}(x, y)
 \end{aligned}$$

Figure 6: Image

Note that any ideal of I_1^\perp is also an ideal of \mathfrak{g} , which implies that $\text{rad}(I_1^\perp) \subseteq \text{rad}(\mathfrak{g})$, which is zero since \mathfrak{g} is semisimple, and thus I_1^\perp is semisimple as well.

By the inductive hypothesis, $I_1^\perp = I_2 \oplus \cdots \oplus I_n$ where each $I_j \trianglelefteq I_i^\perp$ is simple. Then $I_j \trianglelefteq \mathfrak{g} \implies [I_1, I_j] \subset I_1 \cap I_j$, since I_1 has no contribution. But this is a subset of $I_1 \cap I_1^\perp = 0$. \square

Proof of (b): If $I \trianglelefteq \mathfrak{g}$, then $[I, \mathfrak{g}] \trianglelefteq I$ because $[[I, \mathfrak{g}], I] \subseteq [I, I] \subseteq [I, \mathfrak{g}]$.

Since \mathfrak{g} is semisimple, $0 = \text{rad}(\mathfrak{g}) \supseteq Z(\mathfrak{g})$. So $[I, \mathfrak{g}] \neq 0$, and thus $[I, \mathfrak{g}] = I$ since I is simple. But then $[I, \mathfrak{g}] = \bigoplus [I, I_i]$ is simple as well. So only one direct summand can survive, since otherwise this would produce at least 2 nontrivial ideals, and $[I, \mathfrak{g}] = [I, I_i]$ for some i .

So for all $j \neq i$, we must have $I_j \cap I = I_j \cap [I, I_i] = 0$, and so $I \subseteq I_i$. But then $I = I_i$ since I_i itself is simple, and we're done.

Proof of (c):

(Without using the simplicity of I_i)

For $x, y \in I_i$, we have

10.1 Inner Derivations

Recall that $\text{ad } \mathfrak{g} \subseteq \text{Derg}$, and in fact (lemma) this is an ideal.

Theorem: If \mathfrak{g} is semisimple, then $\text{ad } \mathfrak{g} = \text{Derg}$.

Proof of lemma:

For all $\delta \in \text{Der } \mathfrak{g}$ and all $x, y \in \mathfrak{g}$, we have

$$\begin{aligned} [\delta, \text{ad } x](y) &= \delta([x, y]) - [x, \delta(y)] \\ &= [\delta(x), y] \\ &= [\text{ad } \delta(x)](y), \end{aligned}$$

and so $[\delta, \text{ad } x] \subseteq \text{ad } \mathfrak{g}$. \square

Proof of theorem:

If \mathfrak{g} is semisimple, then $0 = \text{rad } \mathfrak{g} \supseteq Z(\mathfrak{g}) = \ker \text{ad}$. Thus $\text{ad } \mathfrak{g} \cong \mathfrak{g} / \ker \text{ad} \cong \mathfrak{g}$ is also semisimple.

This means that $\kappa_{\text{ad } \mathfrak{g}}$ is non-degenerate, and thus $\text{ad } \mathfrak{g} \cap (\text{ad } \mathfrak{g})^\perp = 0$, where $(\text{ad } \mathfrak{g})^\perp \leq \text{Der}(\mathfrak{g})$.

(Note that the non-degeneracy of κ already forces $(\text{ad } \mathfrak{g})^\perp = 0$.)

Then $[(\text{ad } \mathfrak{g})^\perp, \text{ad } \mathfrak{g}] = 0$, and so for all $\delta \in (\text{ad } \mathfrak{g})^\perp$, we have $\delta(x) = [\delta, \text{ad } x]$ by the lemma, but we've shown that this is zero.

But then δ must be zero because ad is an isomorphism, and in particular it is injective. This means that $(\text{ad } \mathfrak{g})^\perp = 0$, and thus $\text{ad } \mathfrak{g} = \mathfrak{g}$. \square

We can use this to define an abstract Jordan decomposition by pulling back decompositions on adjoints:

11 Wednesday September 4th

11.1 4.3: Cartan's Criterion

Lemma: Let $A \subset B$ be two subspaces of $\mathfrak{gl}(V)$ where $\dim V < \infty$. Set $M = \{x \in \mathfrak{gl}(V) \mid [x, B] \subset A\}$. Suppose that $x \in M$ satisfies $\text{Tr}(xy) = 0$ for all $y \in M$. Then x is nilpotent.

Proof: Let $x = s + n$ (where $s = x_s$ and $n = x_n$) be the Jordan decomposition of x . Fix a basis $v_1 \cdots v_m$ of V relative to which s has matrix $\text{diag}(a_1 \cdots a_m)$. Let E be the vector subspace of F over the prime field Q spanned by the eigenvalues $a_1 \cdots a_m$. We have to show that $s = 0$, or equivalently that $E = 0$, since E has finite Q -dimension by construction. It will suffice to show that the dual space E^* is 0, i.e. that any linear functional $f : E \rightarrow Q$ is zero.

Given f , let y be the element of $\mathfrak{gl}(V)$ whose matrix is given by $\text{diag}(f(a_1), \dots, f(a_m))$. If $\{e_{ij}\}$ is a basis of $\mathfrak{gl}(V)$, then $\text{ad } s(e_{ij}) = (a_i - a_j)e_{ij}$ and $\text{ad } y(e_{ij}) = (f(a_i) - f(a_j))e_{ij}$.

Now let $r(t) \in F[t]$ be a polynomial with no constant term, satisfying $r(a_i - a_j) = f(a_i) - f(a_j)$ for all pairs i, j . The existence of such $r(t)$ follows from Lagrange interpolation, and the fact that if $a_i = a_j$ then $0 = r(a_j) - r(a_i) = r(a_i - a_j) = r(0)$, so r has no constant term. Thus there is no ambiguity in the assigned values, since $a_i - a_j = a_j - a_l$ would imply (by linearity of f) that $f(a_i) - f(a_j) = f(a_k) - f(a_l)$. Thus $\text{ad } y = r(\text{ad } s)$.

Note that Lagrange Interpolation is a special case of the Chinese Remainder Theorem for polynomials. If all x_i s are distinct, then $p_i(x) = x - x_i$ are all pairwise coprime. Then dividing $\frac{p(x)}{p_i(x)} = p(x_i)$. So letting $A_1 \cdots A_k$ be constants in k , there is a unique polynomial of degree less than k such that $p(x_i) = A_i$. Thus there is a polynomial $p(x)$ such that $p(x) \equiv A_i \pmod{p_i(x)}$, and $p(x_i) = A_i$.

$$\mathfrak{g} \subseteq \text{End}(V)$$

$$x \xrightarrow{\text{ad}} \text{ad } x$$

$$\parallel \text{JD}$$

$$\parallel \text{JD}$$

$$x_s \mapsto \text{ad } x_s = (\text{ad } x)_s$$

+

$$x_n \mapsto \text{ad } x_n = (\text{ad } x)_n$$



Can recover some x_s and x_n from the adjoints

Figure 7: Image

Now ad_s is the semisimple part of ad_x . By lemma A of 4.2, ad_s can be written as a polynomial in ad_x without a constant term. Therefore ad_y is also a polynomial in ad_x without constant term. By hypothesis, ad_x maps B into A , so we have $\text{ad}_y(B) \subset A$, and so $y \in M$. Using the hypothesis of the lemma, $\text{Tr}(xy) = 0$, and so $\sum a_i f(a_i) = 0$. The left side is a Q -linear combination of elements of E . Applying f , we obtain $\sum f(a_i)^2 = 0$. But the numbers $f(a_i)$ are rational, so this forces all of them to be zero. Finally, f must be identically 0 because the a_i span E . \square

Note that $\text{Tr}([x, y]z) = \text{Tr}(x[y, z])$. To verify this, write $[x, y]z = xyz - yxz$ and $x[y, z] = xyz - xzy$, then use the fact that $\text{Tr}(y(xz)) = \text{Tr}((xz)y)$.

Theorem (Cartan's Criterion): Let $L \leq \mathfrak{gl}(V)$ be a subalgebra with V finite dimensional. Suppose $\text{Tr}(xy) = 0$ for all $x \in [L, L]$ and $y \in L$. Then L is solvable.

Proof: It suffices to show that $[L, L]$ is nilpotent, or just that all $x \in [L, L]$ are nilpotent endomorphisms. We apply the above lemma, with V as given, $A = [L, L]$, and $B = L$, so $M = \{x \in \mathfrak{gl}(V) \mid [x, L] \subset [L, L]\}$. We have $L \subset M$. Our hypothesis is that $\text{Tr}(xy) = 0$ for all $x \in [L, L]$ and $y \in L$. To use the lemma to reach the desired conclusion, we need a stronger result: that $\text{Tr}(xy) = 0$ for $x \in [L, L]$ and $y \in M$.

If $[x, y]$ is a generator of $[L, L]$ and $z \in M$, then $\text{Tr}([x, y]z) = \text{Tr}(x[y, z]) = \text{Tr}([y, z]x)$. By definition of M , $[y, z] \in [L, L]$, so the right side is 0 by hypothesis.

Corollary: Let L be a Lie algebraic such that $\text{Tr}(\text{ad}_x \circ \text{ad}_y) = 0$ for all $x \in [L, L], y \in L$. Then L is solvable.

Proof: Apply the theorem to the adjoint representation of L . We then get $\text{ad } L$ is solvable. Since $\ker \text{ad} = Z(L)$ is also solvable, L itself is solvable.

11.2 Killing Form

11.2.1 Criterion for Semisimplicity

Let L be any lie algebra. If $x, y \in L$, then define $\kappa(x, y) = \text{Tr}(\text{ad}_x \circ \text{ad}_y)$. Then κ is a symmetric bilinear form on L , called the **killing form**.

Theorem: \mathfrak{g} is solvable $\iff \kappa(x, y) = 0$ for all $x \in [\mathfrak{g}, \mathfrak{g}], y \in \mathfrak{g}$.

Proof: \Leftarrow : By Cartan's Criterion.

\Rightarrow : Exercise.

Example: The killing form of $\mathfrak{sl}(2, F)$.

We have

$$\begin{aligned} x &= \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\ h &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\ y &= \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \end{aligned}$$

Then $\text{ad } h = \text{diag}(2, 0, -2)$, and

$$\begin{aligned}\text{ad } x &= \begin{pmatrix} 0 & -2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \\ \text{ad } y &= \begin{pmatrix} 0 & 0 & 0 \\ -1 & 0 & 0 \\ 0 & 2 & 0 \end{pmatrix}.\end{aligned}$$

and thus k has the matrix

$$\begin{pmatrix} 0 & 0 & 4 \\ 0 & 8 & 0 \\ 4 & 0 & 0 \end{pmatrix}.$$

where $k_{ij} = \kappa(x_i, x_j)$ where x_i is a basis of L .

12 Wednesday September 11th

Theorem: If L is semisimple and $x \in L$, there exists a unique x_s, x_n in L such that $x = x_s + x_n$, $[x_n, x_s] = 0$, $\text{ad } x_s$ is semisimple, and $\text{ad } x_n$ is nilpotent.

13 Friday September 13th

Todo

14 Monday September 16th

Let $S = \exp(\text{ad } e) \circ \exp(\text{ad } -f) \circ \exp(\text{ad } ei)$, which has the following matrix:

Where $\exp(\text{ad } e) = 1 + \text{ad } e + \frac{1}{2}(\text{ad } e)^2$, which would have the form

Theorem: If \mathfrak{g} is semisimple, then any finite dimensional \mathfrak{g} -module V is completely reducible, i.e. it splits into a direct sum of simple modules.

14.1 Proof of Weyl's(?) Theorem

If V itself is simple, then we're done, so suppose it is not.

Assume there exists a nonzero submodule $U \subsetneq V$. It suffices to show that $V = U \oplus U'$ for some U' .

$$\begin{pmatrix} & & -1 \\ & -1 & \\ -1 & & \end{pmatrix}$$

Figure 8: Image

$$\begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} + \begin{pmatrix} \cdot & 2 & \\ & \cdot & -1 \\ & & \cdot \end{pmatrix} + \begin{pmatrix} \cdot & \cdot & 1 \\ & \cdot & \cdot \\ & & \cdot \end{pmatrix}$$

Figure 9: Image

14.1.1 Step 1:

If $\dim V = 2$ and $\dim U = 1$.

Then $U, V/U$ are both trivial modules. So $g \curvearrowright u = 0$ for all $u \in U$. But then $g \curvearrowright (v + U) = U$ for all $v \in V$, since $g \curvearrowright v \in U$.

So for all $x, y \in \text{lieg}$ and all $v \in V$, we have $[x, y] \curvearrowright v = x \curvearrowright (y \curvearrowright v) - y \curvearrowright (x \curvearrowright v)$. But both of the terms in parenthesis are in U , and all elements in \mathfrak{g} kill elements in U , so this is zero. So $[\mathfrak{g}, \mathfrak{g}] \curvearrowright V$ trivially.

Exercise: If \mathfrak{g} is semisimple, then $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$.

So $\mathfrak{g} \curvearrowright V$ trivially. Thus any U' that is a complementary subspace of U will be a submodule of V .

14.1.2 Step 2:

Suppose U is simple and $\dim U > 1$, so $\dim V/U = 1$.

Let Ω be the Casimir element on U (faithful representation?). Then $\Omega u = c u$ for some $c \in \mathbb{F}$, and so $\Omega(U) \subseteq U$.

Since $\Omega : V \rightarrow V$ is a homomorphism, $\ker \Omega \subseteq V$ is a \mathfrak{g} -submodule. Then $\dim V/U = 1 \implies V/U$ is a trivial module. So $\mathfrak{g} \curvearrowright V/U = 0$, i.e. $\mathfrak{g} \curvearrowright V \subseteq U$.

Then $\Omega(v) = \sum_i x_i \curvearrowright (y_i \curvearrowright v) \in U$ for all $v \in V$. What is the matrix of Ω ?

In particular, $\text{Tr}(\Omega|_{V/U}) = 0$. So $\text{Tr}(\Omega) = \text{Tr}(\Omega|_U)$. From 6.2, we know that $\text{Tr}(\Omega) \neq 0 \implies c \neq 0$, where c is the scalar appearing above. So $\ker \Omega$ is 1-dimensional, and $\ker \Omega \cap U = \{0\}$.

So take $U' = \ker \Omega$.

14.1.3 Step 3:

Suppose U is *not* simple, but $\dim V/U = 1$.

We will induct on the dimension of U . Pick a proper nonzero submodule $\bar{U} \subsetneq U$, so that $\dim U/\bar{U} < \dim U$. Now $V/U \cong (V/\bar{U})/(U/\bar{U})$ by an isomorphism theorem. So U/\bar{U} is a submodule of V/\bar{U} of codimension 1. Applying the inductive hypothesis, we obtain $V/\bar{U} = U/\bar{U} \oplus \bar{V}/\bar{U}$ for some \bar{V} such that $U \subseteq \bar{V} \subseteq V$.

In particular, since $U \subseteq \bar{V}$ has codimension 1, $\dim \bar{U} < \dim U$. So apply the inductive hypothesis again: $\bar{V} = \bar{U} \oplus U'$ for some U' , and $V = U \oplus U'$.

14.1.4 Step 4: The general case

Recall that $\text{hom}(V, U)$ is a \mathfrak{g} -module where

$$(g \curvearrowright \phi)(v) = g \curvearrowright \phi(v) - \phi(g \curvearrowright v).$$

$$\begin{array}{c}
 u \qquad \qquad \qquad v/u \\
 \left(\begin{array}{c|c}
 u & * \\
 \hline
 v/u & 0 \dots 0
 \end{array} \right)
 \end{array}$$

Handwritten matrix representation on lined paper. The matrix is partitioned into four blocks by a vertical line. The top-left block is labeled u , the top-right block is labeled $*$, the bottom-left block is labeled v/u , and the bottom-right block contains a row of zeros $0 \dots 0$. The central part of the matrix is labeled cI .

Figure 10: Image

Define

$$S = \{\phi \in \text{hom}(V, U) \mid \phi|_U \in F1_U\}.$$

Then $S \leq \text{hom}(V, U)$ as a submodule. Define $T = \{\phi \in S \mid \phi|_U = 0\}$. Then $T \leq S$ as a submodule, and $\mathfrak{g}(S) \subseteq T$.

Now each $\phi \in S$ is determined (mod T) by the scalar $\phi|_U$. Note that $\dim(S/T) = 1$. By steps 1-3, we know that $S = T \oplus T'$ for some $T' \subseteq S$ of dimension 1. Then $T' = \text{span}_{\mathbb{F}}(f)$ for some nonzero map $f : V \rightarrow U$ such that $f(u) = cu$ for some $c \neq 0$.

Then $\mathfrak{g}(T \oplus T') = \mathfrak{g}(S) \subseteq T \implies \mathfrak{g}(T') = 0$. So for all $g \in \mathfrak{g}$, we have $0 = (g \curvearrowright f)(v) = f \curvearrowright f(v) - f(g \curvearrowright v)$. Then $f : V \rightarrow U$ is a lie algebra homomorphism, $\ker f = U'$, and thus $V = U \oplus U'$. \square

Some consequences of Weyl's theorem:

14.2 Preservation of Jordan Decomposition

Recall that when $\mathfrak{g} \in \mathfrak{gl}(V)$ is a linear lie algebra, then for $x \in \mathfrak{g}$ we have:

Jordan Decomposition: $x = x_s + x_n$ where $x_s, x_n \in \text{End}(V)$.

Abstract Jordan Decomposition:

$$\begin{aligned} \mathfrak{g} &\xrightarrow{\text{ad}} \text{ad}(\mathfrak{g}) \\ x &\mapsto \text{ad } x \\ x_s &\leftarrow (\text{ad } x)_s \\ x_n &\leftarrow (\text{ad } x)_n. \end{aligned}$$

and so $x = x'_s + x'_n$ for some x' . The theorem will be that these recover the usual Jordan decomposition.

Theorem: If $\mathfrak{g} \in \mathfrak{gl}(V)$ is semisimple and V is finite dimensional, then $x_s, x_n \in \mathfrak{g}$, and $x_s = x'_s, x'_n$.

Corollary: If \mathfrak{g} is semisimple and finite dimensional and $\phi : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ is a finite dimensional representation, then if $x = x_s + x_n$ is the abstract Jordan decomposition, then $\phi(x) = \phi(x_s) + \phi(x_n)$ is the Jordan decomposition in $\mathfrak{gl}(V)$.

Example: If $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{C})$ is semisimple and finite dimensional, and h is diagonal, then by JD $h = h + 0$, $\phi(h) = \phi(h) + 0$. Then $h \curvearrowright V$ semisimply, or $V = \bigoplus_{\lambda \in \mathbb{C}} V_\lambda$, where $V_\lambda = \{v \in V \mid h \curvearrowright v = \lambda v\}$ are the eigenspaces.

15 Wednesday Lecture

Last time: The abstract Jordan Decomposition coincides with the actual Jordan Decomposition.

$$h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Figure 11: Image

$$\begin{aligned} \phi : \mathfrak{g} &\rightarrow \mathfrak{gl}(V) \\ x &\mapsto \phi(x) = \phi(x)_s + \phi(x)_n = \phi(x_n) + \phi(x_s) \\ x_s + x_n &\mapsto \phi(x_s) + \phi(x_n). \end{aligned}$$

Therefore $x_s \curvearrowright V$ semisimply. The example we saw last time was $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{C})$, with a matrix $h = [1, 0; 0, -1]$ and $V = \bigoplus_{\lambda \in \mathbb{C}} V_\lambda$.

15.1 Finite Dimensional Representations of $\mathfrak{sl}(2, \mathbb{C})$

15.2 Weights and Maximal Vectors

Definition: If $V_\lambda \neq 0$, then V_λ is a *weight space* of V and $\lambda \in \mathbb{C}$ is a *weight* of h in V . We then define $W_t(V) = \{\text{weights in } V\}$.

Lemma: If $v \in V_\lambda$ then $e \curvearrowright v \in V_{\lambda+2}$ and $f \curvearrowright v \in V_{\lambda-2}$.

Proof:

$$\begin{aligned} h \curvearrowright (e \curvearrowright v) &= [h, e] \curvearrowright v + e \curvearrowright (h \curvearrowright v) \\ &= 2e \curvearrowright v + \lambda e \curvearrowright v \\ &= (\lambda + 2)e \curvearrowright v. \end{aligned}$$

and

$$\begin{aligned}
h \curvearrowright (f \curvearrowright v) &= [h, f] \curvearrowright v + f \curvearrowright (h \curvearrowright v) \\
&= -2f \curvearrowright v + \lambda f \curvearrowright v \\
&= (\lambda - 2)f \curvearrowright v.
\end{aligned}$$

So if V is a finite-dimensional \mathfrak{g} -module, then there exists a $V_\lambda \neq 0$ such that $V_{\lambda+2} = 0$. Any nonzero $v \in V_\lambda$ is called a *maximal vector*.

Note: in category \mathcal{O} , these always exist?

Some computations:

- $\mathfrak{g} = \mathfrak{gl}(2, \mathbb{C})$ Then $V = \mathbb{C}$ is the trivial module, and $g \curvearrowright V = 0$. So $W_t(V) = \{0\}$, and $V = V_0$.

If $V = \mathbb{C}^2$, then take the natural representation $\text{span}_{\mathbb{C}} \{v_1 = [1, 0], v_2 = [0, 1]\}$. Then $g \curvearrowright V$ by matrix multiplication, and if $h = [1, 0; 0, -1]$ then $h \curvearrowright v_1 = v_1$ and $h \curvearrowright v_2 = -v_2$ by just doing the matrix-vector multiplication. Then $\mathbb{C}([1, 0]) = V_1, \mathbb{C}([0, 1]) = V_{-1}$, so $W_t(V) = \{\pm 1\}$.

Taking $V = \mathbb{C}^3 = \text{ad } \mathfrak{g} = \text{span}_{\mathbb{C}} \{e, f, h\}$, then

$$\begin{aligned}
h \curvearrowright f &= [h, f] = -2f \\
h \curvearrowright h &= [h, h] = 0h \\
h \curvearrowright e &= [h, e] = 2e.
\end{aligned}$$

So $W_t(V) = \{2, 0, -2\}$ and $V_2 = \mathbb{C}e, V_0 = \mathbb{C}h, V_{-2} = \mathbb{C}f$.

Note the pattern: some largest value, then jumping by 2 to lower values, ending at negative the largest value. In some sense, the rest of the theory will reduce to the case of $\mathfrak{sl}(2, \mathbb{C})$.

Lemma: Let V be a finite dimensional simple $\mathfrak{sl}(2, \mathbb{C})$ -module, and $V_0 \in V_\lambda$ a maximal vector.

Set $V_{-1} = 0, V_i = f^{(i)} \curvearrowright v_0$ (where $f^{(i)} = \frac{f^i}{i!}$). Then for all $i \geq 0$, we have

- $h \curvearrowright v_i = (\lambda - 2i)v_i$
- $f \curvearrowright v_i = (i + 1)v_{i+1}$
- $e \curvearrowright v_i = (\lambda - i + 1)v_{i-1}$

Proof of (a): By lemma 7.1, we have $f \curvearrowright v_0 \in V_{\lambda-2}$, and so inductively $f^{(i)} \curvearrowright v_0 \in V_{\lambda-2i}$

Proof of (b): By definition.


Proof of (c):

$$\begin{aligned}
ie \curvearrowright v_i &= ie \curvearrowright \frac{f^i \curvearrowright v_0}{i!} \\
&= e \curvearrowright (f \curvearrowright v_{i-1}) \\
&= [e, f] \curvearrowright v_{i-1} + f \curvearrowright (e \curvearrowright v_{i-1}) \\
&= h \curvearrowright v_{i-1} + f \curvearrowright ((\lambda - i + 2)v_{i-2}) \text{ind} \\
&= (\lambda - 2i + 2)v_{i-2} + (\lambda + i - 2)(i - 1)v_{i-1} \\
&= i(\text{RHS}).
\end{aligned}$$

Theorem: If V is a finite dimensional and simple, then $V \cong L(m)$ for some $m \in \mathbb{Z}_{\geq 0}$ where $L(m) = \text{span}_{\mathbb{C}} \{v_0, v_1, \dots, v_m\}$ where each v_i is of weight $m - 2i$.

Thus $L(m) = L(m)_m \oplus L(m)_{m-2} \oplus \dots \oplus L(m)_{-m}$ where $\dim L(m)_{\mu} = 1$ for all μ and $\dim L(m) = m + 1$.)

Proof: Pick a maximal vector $v_0 \in V_{\lambda}$ for any weight λ . Define v_i as usual. Let $m = \min \{i \ni V_i \neq 0, V_{i+1} = 0\}$



Definition: A module V is a *highest weight module* of weight λ if $V = \mathfrak{g} \curvearrowright v_0$ for some maximal vector $v_0 \in V_{\lambda}$.

Then λ is referred to as the *highest weight*, and v_0 is the *highest weight vector*.

Corollary: If V is finite-dimensional, then

- $V = \bigoplus_{\lambda \in \mathbb{Z}} V_{\lambda}$
- The number of summands = $\dim V_0 + \dim V_1$.

Proof of (a): By Weyl's theorem, we know $V = \bigoplus W_i$ for some simple W_i . By theorem 7.2, this is equal to $\bigoplus_{m \in \mathbb{Z}_{\geq 0}} L(m)^{\mu_m}$

Proof of (b): $\dim V_0 = \# \{\text{summands where } m \text{ is even}\}$ $\dim V_1 = \# \{\text{summands where } m \text{ is odd}\}$

Remark: Let $V_d = \{f \in \mathbb{C}[x, y] \ni f \text{ is homogeneous of total degree } d\} = \text{span}_{\mathbb{C}} \{x^d, x^{d-1}y, \dots, y^d\}$.

Then $\mathfrak{sl}(2, \mathbb{C}) \curvearrowright V_d$ by

$$\begin{aligned}
e &\mapsto x \frac{\partial}{\partial y} \\
f &\mapsto y \frac{\partial}{\partial x} \\
h &\mapsto x \frac{\partial}{\partial x} - y \frac{\partial}{\partial y}.
\end{aligned}$$

Fact: For $L(m), \phi : \mathfrak{sl}(2, \mathbb{C}) \rightarrow \mathfrak{gl}(L(m))$, define

$$s = (\exp \phi(e)) \circ (\exp \phi(-f)) \circ (\exp \phi(e))$$

Then $s(v_i) = -v_{m-i}$.

16 Friday Lecture

Last time: Construction of simple finite-dimensional $\mathfrak{sl}(2, \mathbb{C})$ module.

Today: Root space decomposition for semisimple finite-dimensional \mathfrak{g} .

16.1 Root Space Decomposition

Let \mathfrak{g} be semisimple and finite dimensional, and let $\mathbb{F} = \mathbb{C}$.

16.1.1 Maximal Toral subalgebra and roots

Definition: A subalgebra $\mathfrak{h} \leq \mathfrak{g}$ is *toral* if $\mathfrak{h} \neq 0$ and it consists of only semisimple elements (i.e. $x_n = 0 \forall x \in \mathfrak{h}$)

Lemma:

- There exists a toral subalgebra of \mathfrak{g} , which is a nontrivial maximal toral subalgebra.
- Any toral subalgebra is abelian.

Proof of (a): Want to show that there exists an $x \in \mathfrak{g}$ such that $x_s \neq 0$, which will imply that $\mathfrak{h} = \mathbb{C}x_s$ is toral.

Suppose $x_s = 0$ for all $x \in \mathfrak{g}$, then $\text{ad } x = \text{ad } x_n$ is nilpotent. By Engel's theorem, this means \mathfrak{g} must be nilpotent. But this contradicts $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$ (since \mathfrak{g} is semisimple) so the derived series can never reach zero.

Proof of (b): Fix $x \in \mathfrak{h}$, want to show that $[x, h] = 0 \forall h \in \mathfrak{h}$. Then $x = x_s$, and so $\text{ad } x : \mathfrak{g} \rightarrow \mathfrak{g}$ is diagonalizable. It suffices to show that $\text{ad } x|_{\mathfrak{h}} = 0$ for all \mathfrak{h} .

Suppose that $[x, h] = ah$ for some vector h where $a \neq 0$. Decompose \mathfrak{h} into eigenspaces, so $\mathfrak{h} = \bigoplus_{\lambda} \mathfrak{h}_{\lambda}$ where $\mathfrak{h}_{\lambda} = \{y \in \mathfrak{h} \mid [h, y] = \lambda y\}$. But then $[h, x] \in \mathfrak{h}_0$, since $[h, [h, x]] = [h, -ah] = 0$.

So write $x = \sum_{\lambda} c_{\lambda} x_{\lambda}$, where $c_{\lambda} \in \mathbb{C}$ and $x_{\lambda} \in \mathfrak{h}_{\lambda}$. Then

$$\begin{aligned} [h, x] &= \sum_{\lambda} c_{\lambda} [h, x_{\lambda}] \\ &= \sum_{\lambda} c_{\lambda} \lambda x_{\lambda} \in \mathfrak{h}_0, \end{aligned}$$

so $\lambda c_{\lambda} = 0 \forall \lambda \neq 0$, which means $c_{\lambda} = 0 \forall \lambda \neq 0$, and thus $x \in \mathfrak{h}_0$ and $[h, x] = 0$. But this contradicts $[x, h] = ah$.

Now $\forall x, h \in \mathfrak{h}, g \in \mathfrak{g}$, we have $[h, [x, y]] = [x, [h, y]] + [y, [x, h]] = [x, [h, y]]$. Thus $\text{ad } h \circ \text{ad } x = \text{ad } x \circ \text{ad } h$ as elements of $\text{End}(\mathfrak{g})$.

So $\mathfrak{g} = \bigoplus_{\alpha \in \mathfrak{h}^*} \mathfrak{g}_{\alpha}$, where $\mathfrak{g}_{\alpha} = \{x \in \mathfrak{g} \mid [h, x] = \alpha(h)x \forall h \in \mathfrak{h}\}$.

Note that $\mathfrak{g}_0 = \{x \in \mathfrak{g} \mid [h, x] = 0 \forall h \in \mathfrak{h}\} = C_{\mathfrak{g}}(\mathfrak{h}) \supseteq \mathfrak{h}$, i.e. the centralizer of \mathfrak{h} in \mathfrak{g} .

Definition: Fix a toral subalgebra $\mathfrak{h} \subseteq \mathfrak{g}$, then a *root* is a nonzero $\alpha \in \mathfrak{h}^*$ such that $\mathfrak{g}_{\alpha} \neq 0$. \mathfrak{g}_{α} is referred to as the *root space*.

We write $\Phi = \{\text{roots}\}$ and $\mathfrak{g} = C_{\mathfrak{g}}(\mathfrak{h}) \oplus (\bigoplus_{\alpha \in \Phi} \mathfrak{g}_{\alpha})$.

Example: $\mathfrak{sl}(3, \mathbb{C})$.

TODO: Insert image from phone.

Then $\Phi = \{\alpha : \mathfrak{h} \rightarrow \mathbb{C}, h_1 \mapsto \alpha(h_1) \in \{\pm 1, \pm 2\}\}$. So

- $\mathfrak{g}_0 = \mathbb{C}h_1 \oplus \mathbb{C}h_2$
- $\mathfrak{g}_1 = \mathbb{C}f_2 \oplus \mathbb{C}e_3$
- $\mathfrak{g}_2 = \mathbb{C}e_1$
- $\mathfrak{g}_{-1} = \mathbb{C}f_3 \oplus \mathbb{C}e_2$
- $\mathfrak{g}_{-2} = \mathbb{C}f_1$.

TODO: Insert second and third image from phone

From these computations, we collect the eigenvalues as ordered pairs. If we choose a larger toral subalgebra, we get a finer decomposition. And if we take a maximal toral subalgebra, then $\mathfrak{h} = \mathfrak{g}_0$ and all $\dim \mathfrak{g}_{\alpha} = 1$.

Proposition (a): $[\mathfrak{g}_{\alpha}, \mathfrak{g}_{\beta}] \subseteq \mathfrak{g}_{\alpha+\beta}$ for all $\alpha, \beta \in \mathfrak{h}^*$.

Proposition (b): If $x \in \mathfrak{g}_{\alpha}$ and $\alpha \neq 0$ then $\text{ad } x$ is nilpotent.

Proposition (c): If $\alpha, \beta \in \mathfrak{h}^*$ and $\alpha + \beta = 0$, then $\kappa(x, y) = 0 \forall x \in \mathfrak{g}_{\alpha}, y \in \mathfrak{g}_{\beta}$.

Proof of (a): Easy exercise:

Proof of (b): For all $y \in \mathfrak{g}$, $y \in \mathfrak{g}_{\mu}$ for some $\mu \in \mathfrak{h}^*$. We have $\mathfrak{g}_u \xrightarrow{\text{ad } x} \mathfrak{g}_{\mu+\alpha} \xrightarrow{\text{ad } x} \mathfrak{g}_{\mu+2\alpha} \rightarrow \dots$ by $y \mapsto [x, y] \mapsto \dots$. Since \mathfrak{g} is finite dimensional, this must terminate, so $(\text{ad } x)^n(y) = 0$ for some n .

Proof of (c): If $\alpha + \beta = 0$, then there exists an $h \in \mathfrak{h}$ such that $\alpha(h) + \beta(h) \neq 0$. Since the killing form is associative, we have

Corollary: $\kappa|_{\mathfrak{g}_0}$ is nondegenerate.

Proof: We want to show $\kappa(h, y) = 0 \forall y \in \mathfrak{g}_0 \implies h = 0$ holds for any choice of $y \in \mathfrak{g}_{\alpha}$ with $\alpha \neq 0$.

$$\begin{array}{ccc}
 K([h, x], y) & = & \alpha(h) K(x, y) \\
 \parallel & \nearrow x \in \mathfrak{g}_\alpha & \\
 - K([x, h], y) & & \\
 \parallel & \nwarrow x \in \mathfrak{g}_\beta & \\
 - K(x, [h, y]) & = & -\beta(h) K(x, y)
 \end{array}$$

Figure 12: Image

By proposition (c), we have $\kappa(h, y) = 0$. Note that we have $\mathfrak{g} = \mathfrak{g}_0 \oplus (\bigoplus_{\alpha \neq 0} \mathfrak{g}_\alpha)$. This implies that $\kappa(h, y) = 0 \forall y \in \mathfrak{g}$. But then $h = 0$ because κ is nondegenerate and \mathfrak{g} is semisimple.

17 Monday Lecture

Last time: \mathfrak{h} is a *toral* subalgebra if it contains only semisimple elements, and implies that there is a *root space decomposition*

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \bigoplus_{\alpha \in \Phi} \mathfrak{g}_\alpha$$

where $\mathfrak{g}_\alpha = \{x \in \mathfrak{g} \mid [h, x] = \alpha(h)x \ \forall h \in \mathfrak{h}\}$ and $\Phi = \{\alpha : \mathfrak{h} \rightarrow \mathbb{C} \mid \mathfrak{g}_\alpha \neq 0, \alpha \neq 0\}$ and $\mathfrak{g}_0 = C_{\mathfrak{g}}(\mathfrak{h})$.

Take larger \mathfrak{h} yields finer decompositions, and a maximal \mathfrak{h} gives $\dim \mathfrak{g}_\alpha = 1 \forall \alpha \in \Phi$.

Corollary: $\kappa|_{\mathfrak{g}_0}$ is nondegenerate.

17.1 The Centralizer of \mathfrak{h}

If $x, y \in \text{End}(V)$ where V is finite dimensional, $xy = yx$, and y is nilpotent, then xy is nilpotent and $\text{Tr}(xy) = 0$.

Proposition: If $\mathfrak{h} \subseteq \mathfrak{g}$ is a maximal toral subalgebra, then $\mathfrak{h} = \mathfrak{g}_0$.

Proof:

Step 1: If $x \in \mathfrak{g}_0$, then $x_s, x_n \in \mathfrak{g}_0$.

If $x \in \mathfrak{g}_0$, then $\text{ad } x(\mathfrak{h}) \subseteq 0$. By proposition 4.2, $\text{ad } x_s(\mathfrak{h}) \subseteq 0, \text{ad } x_n(\mathfrak{h})$, and so $x_s, x_n \in \mathfrak{g}_0$.

Step 2: $\{x_s \ni x \in \mathfrak{g}_0\} \subseteq \mathfrak{h}$.

If $x \in \mathfrak{g}_0$, then by step 1 we have $x_s \in \mathfrak{g}_0$ and so $\mathfrak{h} + \mathbb{C}x_s$ is toral, and thus $x_s \in \mathfrak{h}$.

Step 3: $\kappa|_{\mathfrak{h}}$ is non-degenerate.

We want to show that $\kappa(h, x) = 0 \ \forall x \in \mathfrak{g} \implies h = 0$. By the corollary, it suffices to show that $\kappa(h, x) = 0 \ \forall x \in \mathfrak{g}_0$. By step 2, it suffices to check this only for $x \in \mathfrak{g}_0$ such that $x = x_n$.

If $x = x_n$, then $\text{ad } x_n$ is nilpotent and $\text{ad } h$ commutes with $\text{ad } x$ because $[h, x] = 0$ (since $x \in \mathfrak{g}_0$). By the lemma, $\text{Tr}(\text{ad } h \circ \text{ad } x) = 0$, since $\text{ad } h = \kappa(h, x)$.

Step 4: \mathfrak{g}_0 is nilpotent.

Pick $x \in \mathfrak{g}_0$. Then by step 2, $x_s \in \mathfrak{h}$, so $\text{ad } x_s : \mathfrak{g}_0 \odot$ is a zero map and thus nilpotent.

So $\text{ad } x_n$ is nilpotent, meaning that $\text{ad } x$ is nilpotent. By Engel's theorem, this implies that \mathfrak{g}_0 itself is nilpotent.

Step 5: \mathfrak{g}_0 is abelian.

Suppose that $I := [\mathfrak{g}_0, \mathfrak{g}_0] \neq 0$. We have $I \trianglelefteq \mathfrak{g}_0$, and I is not nilpotent whereas \mathfrak{g}_0 is.

By Lemma 3.3, we have $I \cap Z(\mathfrak{g}_0) \neq 0$, so pick x in the intersection. Note that $\kappa(\mathfrak{h}, I) = \kappa(\mathfrak{h}, [\mathfrak{g}_0, \mathfrak{g}_0])$, which by associativity equals $\kappa([\mathfrak{h}, \mathfrak{g}_0], \mathfrak{g}_0) = 0$.

By step 3, we have $\mathfrak{h} \cap I = 0$. By step 2, $x \neq x_s$, and thus $x_n \neq 0$. But we also have $x \in Z(\mathfrak{g}_0)$, so $[x, \mathfrak{g}_0] = 0$ and $\text{ad } x(\mathfrak{g}_0) \subseteq 0$. By Proposition 4.2, this holds for x_s, x_n as well, which are both in the center. So for all $y \in \mathfrak{g}_0$, $\text{ad } y$ commutes with $\text{ad } x_n$, which is nilpotent.

By the lemma, this implies that $0 = \text{Tr}(\text{ad } y \circ \text{ad } x_n) = \kappa(x_n, y)$ for all $y \in \mathfrak{g}_0$. So $x_n = 0$.

Step 6: Suppose $\mathfrak{g}_0 \not\subseteq \mathfrak{h}$. By step 2, there exists an $x \in \mathfrak{g}_0$ such that $x \notin \mathfrak{h}$, where $x_n \neq 0$. By step 5, $[x_n, y] = 0$ for all $y \in \mathfrak{g}_0$. Then $\text{ad } x$ (which is nilpotent) commutes with $\text{ad } y$. By the lemma, $0 = \kappa(x_n, y)$ for all $y \in \mathfrak{g}_0$, and thus $x_n = 0$. \square

Main idea: Choose a maximal toral subalgebra to get a nice root space decomposition, and so it coincides with \mathfrak{g}_0 .

Corollary: $\kappa|_{\mathfrak{g}}$ is nondegenerate.

Thus for all $\alpha \in \mathfrak{h}^*$, there exists a unique $t_\alpha \in \mathfrak{h}$ such that $\alpha = \kappa(t_\alpha, \cdot) : \mathfrak{h} \rightarrow \mathbb{C}$.

In other words, there is an identification

$$\begin{aligned} \mathfrak{h} &\xrightarrow{1-1} \mathfrak{h}^* \\ h &\mapsto \kappa(h, \cdot) \\ t_\alpha &\leftarrow \alpha. \end{aligned}$$

Definition: A subalgebra $\mathfrak{h} \subseteq \mathfrak{g}$ is a *Cartan subalgebra* if \mathfrak{h} is nilpotent and

$$\mathfrak{h} = N_{\mathfrak{g}}(\mathfrak{h}) = \{x \in \mathfrak{g} \ni [x, h] \subseteq \mathfrak{h}\}.$$

Note that $N_{\mathfrak{g}}(\mathfrak{h})$ is the largest subalgebra of \mathfrak{g} in which \mathfrak{h} is an ideal.

Remark: If \mathfrak{g} is semisimple and finite dimensional with $\text{char}(F) = 0$, we will have a correspondence:

$$\{\text{CSAs of } \mathfrak{g}\} \iff \{\text{maximal toral subalgebras of } \mathfrak{g}\}.$$

Maximal toral subalgebras advantages over Cartan subalgebra definition:

- Yields the finest root space decomposition
- $\mathfrak{h}^* = \mathfrak{h}$, Weyl group?
- Existence is easy compared to CSAs

On the other hand, CSA advantages:

- All CSAs are conjugate under G (some group to be defined)
- The dimensions of all CSAs are the same, giving a well-defined notion of dimension ($\text{rank } \mathfrak{g} = \dim \mathfrak{h}$).

17.2 8.3: Orthogonality Properties

From now on, \mathfrak{h} will be a maximal toral subalgebra.

Proposition: Let $\alpha \in \Phi$. Then

- Φ spans \mathfrak{h}^*
- $-\alpha \in \Phi$
- $\forall x \in \mathfrak{g}_\alpha, y \in \mathfrak{g}_{-\alpha}$, we have $[x, y] = \kappa(x, y)t_\alpha$
- $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}] = \mathbb{C}t_\alpha$ (let the nonzero scalar be λ)
- $\alpha(t_\alpha) = \kappa(t_\alpha, t_\alpha) \neq 0$.
- For any nonzero $e_\alpha \in \mathfrak{g}_\alpha$, there exists a unique $f_\alpha \in \mathfrak{g}_{-\alpha}$ such that $[e_\alpha, f_\alpha] = h_\alpha := \frac{\lambda}{\kappa(t_\alpha, t_\alpha)}t_\alpha$.
Moreover, $\langle e_\alpha, f_\alpha, h_\alpha \rangle = \mathfrak{sl}(2, \mathbb{C})$.

18 Lecture Wednesday

Today: Properties of the root space when the toral subalgebra is maximal.

Last time: We have $\mathfrak{g} = \mathfrak{g} \oplus (\bigoplus_{\alpha \in \Phi} \mathfrak{g}_\alpha)$ where $\kappa|_{\mathfrak{h}}$ is nondegenerate. We also have a correspondence

$$\begin{aligned} \mathfrak{h} &\iff \mathfrak{h}^* h \mapsto \kappa(\mathfrak{h}, \cdot) \\ t_\alpha &\leftarrow \alpha := \kappa(t_\alpha, \cdot). \end{aligned}$$

18.1 Orthogonality Properties

Proposition: Let $\alpha \in \Phi$. Then:

- Φ spans \mathfrak{h}^*
- $-\alpha \in \Phi$

- c. $[x, y] = \kappa(x, y)t_\alpha$ for all $x \in \mathfrak{g}_\alpha$ and $y \in \mathfrak{g}_{-\alpha}$
- d. $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}] = \mathbb{C}t_\alpha$
- e. $\alpha(t_\alpha) = \kappa(t_\alpha, t_\alpha) \neq 0$
- f. For each nonzero $e_\alpha \in \mathfrak{g}_\alpha$, there exists a unique $f_\alpha \in \mathfrak{g}_{-\alpha}$ such that $[e_\alpha, f_\alpha] = h_\alpha := \frac{2}{\kappa(t_\alpha, t_\alpha)}t_\alpha$.
Moreover, $\langle e_\alpha, t_\alpha, h_\alpha \rangle \cong \mathfrak{sl}(2, \mathbb{C})$.

Proof of (a): We want to show that $h \in \mathfrak{h}$ implies that if $\alpha(h) = 0$ for all $\alpha \in \Phi$, then $h = 0$.

Take $x \in \mathfrak{g}_\alpha$. Then $[h, x] = \alpha(h)x = 0$. So $[\mathfrak{h}, \mathfrak{g}] = 0$ because \mathfrak{h} is abelian. But then $[h, \mathfrak{g}] = 0$, or $h \in Z(\mathfrak{g}) = 0$ since \mathfrak{g} is semisimple.

Proof of (b): By Proposition 8.1c, we have $\kappa(\mathfrak{g}_\alpha, \mathfrak{g}_\beta) = 0$ for all $\beta \neq -\alpha$.

If $-\alpha \notin \Phi$, then $\mathfrak{g}_{-\alpha} = 0$. So $\kappa(\mathfrak{g}_\alpha, \mathfrak{g}) = 0$ by the non-degeneracy of κ .

Proof of (c): For all $h \in \mathfrak{h}$, we have

$$\begin{aligned}
 \kappa(h, [x, y]) &= \kappa([h, x], y) \\
 &= \kappa(\alpha(h)x, y) \\
 &= \kappa(t_\alpha, h)\kappa(x, y) \\
 &= \kappa(\kappa(x, y)t_\alpha, h) \\
 &= \kappa(h, \kappa(x, y)t_\alpha).
 \end{aligned}$$

which implies that $\kappa(h, [x, y]) - \kappa(x, y)t_\alpha = 0$, which forces the second argument to be zero by non-degeneracy.

Proof of (d): We will show that (d) implies (c), i.e. $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}] \subseteq \mathbb{C}t_\alpha$.

We want to show $\kappa(x, y)$ is not always zero.

Pick any nonzero $x \in \mathfrak{g}_\alpha$. Then $\kappa(x, \mathfrak{g}_\beta) = 0$ for all $\beta \neq -\alpha$. If $\kappa(x, \mathfrak{g}_{-\alpha}) = 0$, then $\kappa(x, \mathfrak{g}) = 0$. By non-degeneracy, this forces $x = 0$.

Proof of (e): We will skip this for now, and revisit with methods from later sections that make this proof simpler.

Proof of (f): Let $e_\alpha \neq 0$ in \mathfrak{g}_α . Then there exists a $y \in \mathfrak{g}_{-\alpha}$ such that $\kappa(e_\alpha, y) \neq 0$. Set $f_\alpha \in \mathfrak{g}_{-\alpha}$ such that $\kappa(e_\alpha, f_\alpha) = \frac{2}{\kappa(t_\alpha, t_\alpha)}$.

By (c), we have

$$\begin{aligned}
 [e_\alpha, f_\alpha] &= \kappa(e_\alpha, t_\alpha)t_\alpha \\
 &= \frac{2}{\kappa(t_\alpha, t_\alpha)}t_\alpha \\
 &= h_\alpha.
 \end{aligned}$$

and

$$\begin{aligned}
[h_\alpha, e_\alpha] &= \frac{2}{\kappa(t_\alpha, t_\alpha)} [t_\alpha, e_\alpha] \\
&= \frac{2}{\kappa(t_\alpha, t_\alpha)} \alpha(t_\alpha) e_\alpha \\
&= 2e_\alpha.
\end{aligned}$$

and similarly $[h_\alpha, f_\alpha] = -2f_\alpha$.

Definition:

Let $\mathfrak{sl}(2, \alpha) = \langle e_\alpha, f_\alpha, h_\alpha \rangle$ as in (f). A priori, this depends on a choice of $e_\alpha \neq 0$. We will show that this only depends on α .

18.2 Orthogonality/Integrality Properties

Proposition: Let $\alpha \in \Phi$. Then:

- a. $\dim \mathfrak{g}_\alpha = 1$. (Note that in general, $\dim \mathfrak{g}_0 = \dim \mathfrak{h} \geq 1$)
- b. $\mathbb{C}\alpha \cap \Phi = \{\pm\alpha\}$
- c. If $\beta \in \Phi$ such that $\alpha + \beta \in \Phi$, then $[\mathfrak{g}_\alpha, \mathfrak{g}_\beta] = \mathfrak{g}_{\alpha+\beta}$.
- d. If $\beta \neq -\alpha \in \Phi$, then let $p, q \in \mathbb{Z}$ be the largest such that $\beta - p\alpha$ and $\beta + q\alpha$ are both in Φ . Then $\beta + i\alpha \in \Phi$ for every $-p \leq i \leq q$, and

$$\beta(h_\alpha) = \kappa(t_\beta, h_\alpha) = \frac{2\kappa(t_\beta, t_\alpha)}{\kappa(t_\alpha, t_\alpha)} = p - q \in \mathbb{Z}.$$

Proof of (a) and (b):

Let $M = \mathfrak{h} \oplus \left(\bigoplus_{c \neq 0} \mathfrak{g}_{c\alpha} \right) \leq \mathfrak{g}$ as a subspace. By a routine check, M is an $\mathfrak{sl}(2, \alpha)$ submodule of \mathfrak{g} . Recall that $M = \bigoplus_{\lambda \in \mathbb{Z}} L(\lambda)$ as a direct sum of vector spaces. Applying Weyl's theorem, we also have $M = \bigoplus_{m \in \mathbb{Z}_{\geq 0}} L(m)^{\oplus \mu_m}$ as a direct sum of (irreducible?) modules.

For \mathfrak{h} , if we have $[h_\alpha, h] = 0$ for all $h \in \mathfrak{h}$, then $h \in M_0$. For $\mathfrak{g}_{c\alpha}$, $[h_\alpha, x] = c\alpha(h_\alpha)x$ for all $x \in \mathfrak{g}_{c\alpha}$. But this equals $2cx$. So this implies that $\mathfrak{g}_{c\alpha} \subseteq M_{2c}$.

Thus $2c \in \mathbb{Z}$, and thus $c \in \frac{1}{2}\mathbb{Z}$, and $M_0 = \mathfrak{h}$.

We then have $\dim M_0 = \sum_{m \in 2\mathbb{Z}} \mu_m$. So write $h = \mathbb{C}t_\alpha \oplus \ker \alpha$ as vector spaces. Consider the action $\mathfrak{sl}(2, \alpha) \curvearrowright \ker \alpha$, which is trivial since $h \in \ker \alpha$. We $[h_\alpha, h] = 0$, $[e_\alpha, h] = -\alpha(h)e_\alpha = 0$ since $h \in \ker \alpha$, and similarly $[f_\alpha, h] = 0$.

Thus $\ker \alpha = L(0)^{\oplus \dim \mathfrak{h} - 1}$. Moreover, $\mathfrak{sl}(2, \alpha) = L(2) = \text{span}(e_\alpha, t_\alpha, f_\alpha)^T$. But this forces the case that there is no other summand of the form $L(k)$ for k even in M .

Then $\mathfrak{g}_{2\alpha} \subseteq M_4$, which must be zero. So $2\alpha \notin \Phi$, so 2α is not a root. ("Twice a root is never a root")

So $\frac{1}{2}\alpha \notin \Phi$, otherwise we could apply this argument to conclude that α is not a root and reach a contradiction. Thus $M_1 = 0$, since $c \neq \frac{1}{2}$ implies that there is not summand of the form $L(k)$ for k odd in M . But this forces $M = \mathfrak{h} \oplus \mathfrak{sl}(2, \alpha)$.

Motto: reduce the complexity by using the $\mathfrak{sl}(2)$ module structure and its representation theory!

19 Lecture (Friday)

Last time, we saw $\Phi \subseteq \mathfrak{h}^* = \{\alpha : \mathfrak{h} \rightarrow \mathbb{C}\}$.

Suppose \mathfrak{g} is semisimple and \mathfrak{h} is a maximal toral subalgebra and take $F = \mathbb{C}$.

We have the following propositions:

- a. $\dim \mathfrak{g}_\alpha = 1 \ \forall \alpha \in \Phi$
- b. $\mathbb{C}\alpha \cap \Phi = \{\pm\alpha\}$, and $2\alpha \notin \Phi$ where $c\alpha : \mathfrak{h} \rightarrow \mathbb{C}, h \mapsto c \curvearrowright \alpha(h)$. Moreover, $M = \mathfrak{h} \oplus \left(\bigoplus_{c \neq 0} \mathfrak{g}_{c\alpha} \right)$
- c. If $\alpha, \beta \in \Phi$ and $\beta \neq -\alpha$ Let $p, q \in \mathbb{Z}$ be the largest such that $\beta - p\alpha$ and $\beta + q\alpha$ are in Φ . Moreover, $\beta(h_\alpha) = \kappa(t_\beta, t_\alpha) = p - q \in \mathbb{Z}$.

Proof of (c):

Set $M = \sum_{i \in \mathbb{Z}} \mathfrak{g}_{\beta+i\alpha}$, which is an $\mathfrak{sl}(2, \alpha)$ module. By (a), we have $\dim \mathfrak{g}_{\beta+i\alpha} = 1 \iff \beta+i\alpha \in \Phi$. But for all $x \in \mathfrak{g}_{\beta+i\alpha}$, we have $[h, x] = (\beta + i\alpha)(h)x$ for all $h \in \mathfrak{h}$. But then $[h_\alpha, x] = (\beta(h_\alpha) + i\alpha(h_\alpha))x = (\beta(h_\alpha) + 2i)x$

Then $\mathfrak{g}_{\beta+i\alpha} \subseteq M_{\beta(h_\alpha)+2i}$, so $\beta(h_\alpha) \in \mathbb{Z}$.

Moreover, $\text{Wt}(M) = 2\mathbb{Z}$ or $2\mathbb{Z} + 1$, and in particular $\dim M_0 + \dim M_1 = 1$.

Thus M is irreducible, and $M \cong L(m)$ for some $m \in \mathbb{Z}_{\geq 0}$. Moreover, $\text{Wt}(M) = \{m, m-2, \dots, -m\}$, and $\dim \mathfrak{g}_{\beta+i\alpha} = 1$ for all $i \in [-p, q]$. Thus $\beta + i\alpha \in \Phi$.

Proof of 8.3(e): $\alpha(t_\alpha) \neq 0$. The claim is that for all $\beta \in \Phi$, there exists an $r \in \mathbb{Q}$ such that $\beta(h) = r\alpha(h)$ for all $h \in [\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}]$.

There are two cases: if $\beta = -\alpha$, then we're done by the previous argument.

Otherwise, $\beta \neq -\alpha$. Take $M = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}_{\beta+i\alpha}$.

Then,

$$\begin{aligned} \text{Tr}_M(\text{ad } h) &= \sum_i \text{Tr}_M((\text{ad } e_i \circ \text{ad } f_i) - (\text{ad } f_i \circ \text{ad } e_i)) = \sum_i \text{Tr}_{\mathfrak{g}_{\beta+i\alpha}}(\text{ad } h) \\ &= \sum_i (\beta + i\alpha)(h) \dim \mathfrak{g}_{\beta+i\alpha} \\ &= \sum_i \dim \mathfrak{g}_{\beta+i\alpha} \beta(h) + \sum_i i \dim(\mathfrak{g}_{\beta+i\alpha}) \\ &\implies \beta(h) = \frac{-\sum_i \dim \mathfrak{g}_{\beta+i\alpha}}{\sum_i \dim \mathfrak{g}_{\beta+i\alpha}} \alpha(h). \end{aligned}$$