

Interesting Topological Spaces in Algebraic Geometry

D. Zack Garza

Tuesday 28th July, 2020

Contents

1	Ideas for Spaces	1
2	Analogies	2
2.1	Topological Category	3
2.2	Smooth Category	3
3	Kahlers	3
4	Calabi-Yaus	3

1 Ideas for Spaces

- Curves
 - Elliptic Curves
 - Higher genus
 - Hyperelliptic curves
 - The modular curve
- Surfaces
 - Compact Riemann surfaces
 - * Bolza Surface (Genus 2)
 - * Klein Quartic (Genus 3)
 - * Hurwitz Surfaces
 - Kummer surfaces
- Compact Complex Surfaces
 - Rational ruled
 - Enriques Surfaces
 - $K3$
 - * Kahler Manifolds
 - Kodaira
 - Toric
 - Hyperelliptic
 - Properly quasi-elliptic

-
- General type
 - Type VII
 - Fake projective planes
 - Conics
 - Calabi-Yau manifolds
 - Dimension 1: All elliptic curves (up to homeomorphism)
 - Dimension 2: $K3$ surfaces
 - Dimension 3 (threefolds): 500 million +, unknown if infinitely many
 - The bananafold
 - Hyperkähler
 - Hurwitz schemes
 - Topological galois groups, e.g. $G(\bar{F}/F)$ for $F = \mathbb{Q}, \mathbb{F}_p$.
 - $\text{Spec}(R)$ for R a DVR (a Sierpinski space)
 - Quiver Grassmannians
 - Rigid analytic spaces
 - Affine line with two origins
 - Moduli stack of elliptic curves $\mathcal{M}_{1,1}$.
 - Abelian Surface
 - Fano Varieties
 - Curves: isomorphic to \mathbb{P}^1
 - Surfaces: Del Pezzo surfaces
 - Weighted projective space
 - Toric Varieties
 - Grassmannian
 - Flag Varieties
 - Moduli Spaces

Due to Kunihiko Kodaira's classification of complex surfaces, we know that any compact hyperkähler 4-manifold is either a $K3$ surface or a compact torus T^4 . (Every Calabi-Yau manifold in 4 (real) dimensions is a hyperkähler manifold, because $SU(2)$ is isomorphic to $Sp(1)$.)

As was discovered by Beauville, the Hilbert scheme of k points on a compact hyperkähler 4-manifold is a hyperkähler manifold of dimension $4k$. This gives rise to two series of compact examples: Hilbert schemes of points on a $K3$ surface and generalized Kummer varieties.

2 Analogies

Notation: all dimensions are over \mathbb{R} .

Impossible goal: pick a category, understand all of the objects and all of the maps. Two main categories with a forgetful functor: **Diff** \longrightarrow **Top**. Question:

- What's in the "image" of this functor? (Manifolds that admit a differentiable structure.)
- What is the "fiber" above a given topological manifold? (Distinct differentiable structures)

Differentiable Manifolds: classified by geometric structure in low dimensions (≤ 4), algebraic data/methods in high dimensions

2.1 Topological Category

Identify objects up to homeomorphism

- Initial object: empty set
- Dimension 0: The point (terminal object)
- 1-manifolds: S^1, \mathbb{R}
- 2-manifolds: $\langle S, T, \mathbb{RP} \mid S = 0, 3\mathbb{RP} = \mathbb{RP} + T \rangle$. Classified by π_1 (orientability and “genus”). Riemann, Poincare, Klein.

2.2 Smooth Category

- 2-manifolds: Homeomorphic \iff diffeomorphic. Every surface admits a complex structure and a metric.
 - Uniformization: Conformally equivalent to a quotient of one of three spaces
 - * \mathbb{CP}^1 , positive curvature (spherical)
 - * \mathbb{D}° , zero curvature (flat)
 - * \mathbb{H} , negative curvature (hyperbolic)
- 3-manifolds: Thurston’s Geometrization
 - Oriented prime 3-manifolds can be decomposed into geometric “pieces” of 8 possible types
 - Geometric structure: a diffeo $M \cong \tilde{M}/\Gamma$ where Γ is a discrete Lie group acting freely/transitively on X
- 4-manifolds: classified in the topological category by surgery, but not in the smooth category
- n -manifolds, $n \geq 5$: classified by surgery

3 Kahlers

- For complex manifold, replace Riemannian metric with a Hermitian metric (positive definite sesquilinear inner product on tangent bundle)
- If skew-symmetric part is symplectic (closed and nondegenerate) then the metric is Kahler
- Includes smooth projective varieties, but not all complex manifolds
- Specialize to Calabi-Yaus: compact and Ricci-flat, or first Chern class vanishes

4 Calabi-Yaus

- As manifolds: Ricci-flat, i.e. Ricci curvature tensor vanishes (measures deviation of volumes of “geodesic balls” from Euclidean balls of the same radius).
- Applications: Physicists want to study G_2 manifolds (an exceptional Lie group, automorphisms of octonions), but no smooth or complex structures. Indirect approach: compactify an 11-dimension space, one small S^1 dimension \longrightarrow 10 dimensions, 4 spacetime and 6 “small” Calabi-Yau.