

Title

D. Zack Garza

Contents

1 Lecture 14: Monday, November 02

3

1 | Lecture 14: Monday, November 02

Recall that the *Hasse-Weil zeta function* of a one-variable function field K/\mathbb{F}_q over a finite ground field is defined in the following way: let $A_n = A_n(K)$ be the number of effective divisors of degree n . We have proved that A_n is finite, and for $n > 2g - 2$ we have a formula

$$Z(t) = \sum_{n=0}^{\infty} A_n t^n = \sum_{D \in \text{Div}^+(K)} t^{\deg(D)} \in \mathbb{Z}[[t]],$$

which is a formal power series with integer coefficients.

Remark 1.0.1 : Recall that we have proved that it is a rational function of t , and in particular when $g = 0, \delta = 1$ ¹ we get

$$Z(t) = \frac{1}{(1 - qt)(1 - t)}.$$

We got another expression which isn't fantastic: it involves this δ , which we'll work toward proving is equal to 1. When $g > 1$, we broke the zeta function into two pieces $Z(t) = F(t) + G(t)$. For divisors of sufficiently high degree, Riemann-Roch tells you what the dimension of the Riemann-Roch space is, and $G(t)$ explains the part coming from divisors of large degree. We obtained a formula previously for $F(t)$ and $G(t)$, and once we show $\delta = 1$ the formula for G will simplify. For $F(t)$, we specifically had

$$F(t) = \frac{1}{q-1} \sum_{0 \leq \deg(c) \leq 2g-2} q^{\ell(c)t^{\deg(c)}},$$

where the sum is over divisor classes and ℓ is the dimension of linear system corresponding to a divisor. But this isn't a great formula: what are these classes, how many are in each degree, and what is the dimension of the Riemann-Roch space?

Remark 1.0.2 : This is analogous to the Dedekind zeta function of a number field K , in which case

$$\zeta_K(s) = \sum_{T \in \ell(\mathbb{Z}_K)} |\mathbb{Z}_K/I|^{-s},$$

which will be covered in a separate lecture on *Serre zeta functions*.

Theorem 1.0.1 (F.K. Schmidt).

For all K/\mathbb{F}_q , we have $\delta = I(K) = 1$ where I is the index.

This will follow from the associated, but it much weaker. However, this is one of the facts we'd like to establish to use to *prove* the Riemann hypothesis.

¹The *index* of the function field, least positive degree of a divisor.

Remark 1.0.3 : Pete studied this in 2004 and found that every $I \in \mathbb{Z}^+$ arises as the index of a genus one function field K/\mathbb{Q} .

Notation: for $n \in \mathbb{Z}^+$, let μ_n denote the n th roots of unity in \mathbb{C} .

Lemma 1.1(?).

For $m, r \in \mathbb{Z}^+$, set $d := \gcd(m, r)$. Then

$$\left(1 - t^{mr/d}\right)^d = \prod_{\xi \in \mu_r} 1 - (\xi t)^m.$$

Proof (?).

In $\mathbb{C}[x]$, we have

$$(X^{r/1} - 1)^d = \prod_{\xi \in \mu_r} (X - \xi^m),$$

where both sides are monic polynomials whose roots include the (r/d) th roots of unity, each with multiplicity d . On the LHS, the distinct roots are the r/d th roots of unity, then raising to the d th power gives them multiplicity d . On the RHS, this is an exercise in cyclic groups: consider the n th power map on $\mathbb{Z}/r\mathbb{Z}$ and compute its image and kernel. As ξ ranges over r th roots of unity, ξ^m ranges over all r/d th roots of unity, each occurring with multiplicity d . Substituting $X = t^{-m}$ and multiplying both sides by t^r yields the original result.

Special case: set $m = r$, so $d = r$, then the RHS is r copies of 1. ■

Next up, we want to compare the zeta function $Z(t)$ for a function field over \mathbb{F}_q to the zeta function obtained when extending scalars to \mathbb{Q}^r .

Proposition 1.0.1(?).

Let K/\mathbb{F}_q be a function field, $r \in \mathbb{Z}^+$, and take the compositum K_r of K and \mathbb{F}_q^r viewed as a function field over \mathbb{F}_q^r . Let $Z(t)$ be the zeta function of K/\mathbb{F}_q and $Z_r(t)$ the zeta function of K_r/\mathbb{F}_q^r . Then

$$Z_r(t^r) = \prod_{q \in \mu_r} Z(qt).$$

Proof (?).

We have an Euler product formula

$$Z(t) = \prod_{p \in \Sigma(K/\mathbb{F}_q)} (1 - t^{\deg(p)})^{-1}.$$

where the sum is over places of the function field.

Proving this Euler product formula might show up in a separate lecture, but it is not any more difficult than proving it for the Riemann zeta function.

Exercise 1.0.1 (?): Why is this product expansion true? Write as a geometric series with ratio $t^{\deg(p)}$. Here just expand each summand to get

$$Z(t) = \prod_p \sum_{j=1}^{\infty} t^{j \deg(p)}.$$

Multiplying this out and collecting terms is in effect multiplying out the prime divisors to get effective divisors.

We now use the result about splitting that was stated (but not proved):

Claim: If $p \in \Sigma_m(K/\mathbb{F}_q)$ is a degree n place and $r \in \mathbb{Z}^+$, then there exist precisely $d := \gcd(m, r)$ places p^r of K_r lying over p . Moreover, each place p^r has degree m/d .

We now have

$$Z_r(t^r) = \prod_{p \in \Sigma(K/\mathbb{F}_q)} \prod.$$

■