Title

D. Zack Garza

November 26, 2019

Contents

1 Wednesday November 20

Last time:

$$\begin{aligned}
\mathbb{Z}\Lambda &\iff \{\mathfrak{h}^* \to \mathbb{Z}_{\geq 0} \mid \sim\} \\
&e(\mu) \mapsto e_\mu \\
e(\lambda)e(\mu) = e(\lambda+\mu) \mapsto f \star g(\lambda) &= \sum_{a+b=\lambda} f(a)g(b)
\end{aligned}$$

and $\mathrm{ch} L(\lambda) = \sum_{\mu \in \Lambda} \dim L(\lambda)_\mu e(\mu)$.

We have the Kostant function $p(\lambda) = \#\left\{(k_\alpha)_\alpha \mid -\lambda = \sum_{\alpha \in \Phi^+} k_\alpha \alpha\right\}$ and the Weyl function $q = e_\rho \star \prod_{\alpha\in\Phi^+}(1 - e_{-\alpha}) = \prod_{\alpha \in \Phi^+}(e_{\alpha/2} - e_{-\alpha/2})$.

Lemma: $p \star e_\lambda = \mathrm{ch} M(\lambda)$, so $q \star \mathrm{ch} M(\lambda) = e_{\lambda+\rho}$ and $q \star p = e_\rho$.

1.1 Weyl's Character Formula (24.2-3)

Definition: The *dot action* of W is given by $w \cdot \lambda = w(\lambda + \rho) - \rho$, i.e. a reflection for hyperplanes passing through $-\rho$.

E.g. for type $A2$, where $W(0) = 0$, we have:

Type A2

And for the dot action, we have

Image

where $W \cdot 0 = 0$ and $s(\alpha_1) = -\alpha_1$.

Theorem (Harish-Chandra): If $L(\mu)$ is a composition factor of $M(\lambda)$, then $\mu \in W \cdot \lambda$ for $\mu \leq \lambda$.

Proof: Postponed.

ch are characters, $L(\lambda)$ is a Verma module.

Remark: if we sum over $\mu \leq \lambda$, we obtain

$$\begin{aligned}
\mathrm{ch}M(\lambda) &= \sum_{\mu \in W\cdot\lambda} a_{\lambda\mu}\mathrm{ch}L(\mu) \\
\mathrm{ch}L(\lambda) &= \sum_{\mu \in W\cdot\lambda} b_{\lambda\mu}\mathrm{ch}M(\mu) &&= \sum_{W\cdot\lambda\in\Lambda} c_{\lambda W}\mathrm{ch}M(w\cdot\lambda).
\end{aligned}$$