

Notes on Lee's Manifolds

D. Zack Garza

Sunday 21st June, 2020

Contents

1	Point Set Review	1
1.1	Quotients	1
1.2	Subspaces	2
1.3	Products	2
1.4	Misc	3
2	Chapter 1	3
2.1	Recommended Problems	3
2.2	Notes	5

1 Point Set Review

1.1 Quotients

Definition 1.0.1 (Saturated).

A subset $A \subseteq X$ is *saturated* with respect to $p : X \rightarrow Y$ if whenever $p^{-1}(\{y\}) \cap A \neq \emptyset$, then $p^{-1}(\{y\}) \subseteq A$.

Equivalently, $A = p^{-1}(B)$ for some $B \subseteq Y$, i.e. it is a complete inverse image of some subset of Y , i.e. A is a union of fibers $p^{-1}(b)$.

Definition 1.0.2 (Quotient Map).

A continuous surjective map $p : X \rightarrow Y$ is a *quotient map* if $U \subseteq Y$ is open **iff** $p^{-1}(U) \subset X$ is open.

Note that \Rightarrow comes from the definition of continuity of p , but \Leftarrow is a stronger condition.

Equivalently, p maps saturated subsets of X to open subsets of Y .

Definition 1.0.3 (Universal Property of Quotients).

For $\pi : X \rightarrow Y$ a quotient map, if $g : X \rightarrow Z$ is a map that is constant on each $p^{-1}(\{y\})$, then there is a unique map f making the following diagram commute:

$$\begin{array}{ccc}
 X & & \\
 \downarrow \pi & \searrow g & \\
 Y & \xrightarrow{f} & Z
 \end{array}$$

Fact: an injective quotient map is a homeomorphism.

Fact: a product of quotient maps need not be a quotient map.

1.2 Subspaces

Definition 1.0.4 (The Subspace Topology).

$U \subset A$ is open iff $U = V \cap A$ for some open $V \subseteq X$.

Proposition 1.1 (*Universal Property of Subspaces*).

If X and $\iota_S : S \hookrightarrow Y$ is a subspace, then every continuous map $f : X \rightarrow S$ lifts to a continuous map $\tilde{f} : X \rightarrow Y$ where $\tilde{f} := \iota_S \circ f$:

$$\begin{array}{ccc}
 & & Y \\
 & \nearrow \exists! \tilde{f} & \uparrow \iota_S \\
 X & \xrightarrow{f} & S
 \end{array}$$

Note that we can view $\iota_S := \text{id}_Y|_S$. The subspace topology is the unique topology for which this property holds.

Some properties of subspace:

- The inclusion ι_S is a topological embedding.
- Restricting a continuous map to a subspace is still continuous.
- A basis for the subspace topology for $A \subset X$ can be obtained by intersecting basis elements of X with A .
- If X is Hausdorff/first/second-countable, then so is A .

1.3 Products

Definition 1.1.1 (The Product Topology).

The coarsest topology such that every projection map $p_\alpha : \prod_{\beta} X_\beta \rightarrow X_\alpha$ is continuous, i.e. for

every $U_\alpha \subseteq X_\alpha$ open, $p_\alpha^{-1}(U_\alpha) \in \prod_{\beta} X_\beta$ is open. For finite index sets, we can take the box topology: the collection of sets of the form $\prod_{i=1}^N U_i$ with each U_i open in X_i forms a basis for the

product topology on $\prod_{i=1}^N X_i$.

Why these differ: in \mathbb{R}^∞ , the set $S = \prod (-1, 1)$ is open in the box topology but not the product topology, since $\{0\}^\infty$ is not contained in any basic open neighborhood contained in S .

Some properties of products:

- Projections π_i are continuous by definition.
- A basis for the product topology can be obtained by taking the product of bases.
- A map $f : X \rightarrow \prod Y_i$ into a product is continuous iff each component function $F_i := \pi_i \circ f : X \rightarrow Y_i$ is continuous.
 - I.e. if we have continuous maps $f_i : X \rightarrow Y_i$ then the composite map $F = [f_1, f_2, \dots]$ is continuous.
- Separate continuity does not imply joint continuity: A map $f : \prod X_i \rightarrow Y$ out of a product need not be continuous even if (defining $\iota_j : X_j \hookrightarrow \prod X_i$) the map $f \circ \iota_j : X_j \rightarrow Y$ is continuous for all arbitrary inclusions ι_j .
- Any map of the form $f_{\mathbf{a}_j} : X_j \rightarrow \prod_{i=1}^n X_i$ where $x \mapsto (a_1, \dots, a_{j-1}, x, a_{j+1}, \dots, a_n)$ is a topological embedding.
- If X_i are Hausdorff/first/second-countable, then so is $\prod_{i=1}^n X_i$.

1.4 Misc

Definition 1.1.2 (Precompact).

A subset $A \subseteq X$ is *precompact* iff its closure $\text{cl}_X(A)$ is compact in X .

Definition 1.1.3 (Locally Compact).

A space X is *locally compact* iff every $x \in X$ has a neighborhood which is contained in some compact subset of X .

2 Chapter 1

Pages 1- 29.

2.1 Recommended Problems

Exercise (Problem 1.6) Show that if $M^n \neq \emptyset$ is a topological manifold of dimension $n \geq 1$ and M has a smooth structure, then it has uncountably many distinct ones.

Hint: show that for any $s > 0$ that $F_s(x) := |x|^{s-1}x$ defines a homeomorphism $F_x : \mathbb{D}^n \rightarrow \mathbb{D}^n$ which is a diffeomorphism iff $s = 1$.

Recommended problem

Exercise (Problem 1.7) Let $N := [0, \dots, 1] \in S^n$ and $S := [0, \dots, -1]$ and define the stereographic projection

$$\sigma : S^n \setminus N \rightarrow \mathbb{R}^n$$

$$[x^1, \dots, x^{n+1}] \mapsto \frac{1}{1 - x^{n+1}} [x^1, \dots, x^n]$$

and set $\tilde{\sigma}(x) = -\sigma(-x)$ for $x \in S^n \setminus S$ (projection from the South pole)

Recommended problem



Fig. 1.13 Stereographic projection

1. For any $x \in S^n \setminus N$ show that $\sigma(x) = \mathbf{u}$ where $(\mathbf{u}, 0)$ is the point where the line through N and x intersects the linear subspace $H_{n+1} := \{x^{n+1} = 0\}$.

Similarly show that $\tilde{\sigma}(x)$ is the point where the line through S and x intersects H_{n+1} .

2. Show that σ is bijective and

$$\sigma^{-1}(\mathbf{u}) = \sigma^{-1}\left([u^1, \dots, u^n]\right) = \frac{1}{\|\mathbf{u}\|^2 + 1} [2u^1, \dots, 2u^n, \|\mathbf{u}\|^2 - 1].$$

3. Compute the transition map $\tilde{\sigma} \circ \sigma^{-1}$ and verify that the atlas

$$\mathcal{A} := \{(S^n \setminus N, \sigma), (S^n \setminus S, \tilde{\sigma})\}$$

define a smooth structure on S^n .

4. Show that this smooth structure is equivalent to the standard smooth structure: Put graph coordinates on S^n as outlined in 2.2 to obtain $\{(U_i^\pm, \varphi_i^\pm)\}$.

For indices $i < j$, show that

$$\varphi_i^\pm \circ (\varphi_j^\pm)^{-1} [u^1, \dots, u^n] = [u^1, \dots, \hat{u}^i, \dots, \pm \sqrt{1 - \|\mathbf{u}\|^2}, \dots, u^n]$$

where the square root appears in the j th position. Find a similar formula for $i > j$. Show that if $i = j$, then

$$\varphi_i^\pm \circ (\varphi_j^\pm)^{-1} = \varphi_i^- \circ (\varphi_i^+)^{-1} = \text{id}_{\mathbb{D}^n}.$$

Show that these yield a smooth atlas.

Exercise (Problem 1.8) Define an *angle function* on $U \subset S^1$ as any continuous function $\theta : U \rightarrow \mathbb{R}$ such that $e^{i\theta(z)} = z$ for all $z \in U$.

Recommended problem

Show that U admits an angle function iff $U \neq S^1$, and for any such function θ , (U, θ) is a smooth coordinate chart for S^1 with its standard smooth structure.

Exercise (Problem 1.9) Show that \mathbb{CP}^n is a compact $2n$ -dimensional topological manifold, and show how to equip it with a smooth structure, using the correspondence

Recommended problem

$$\begin{aligned} \mathbb{R}^{2n} &\iff \mathbb{C}^n \\ [x^1, y^1, \dots, x^n, y^n] &\iff [x^1 + iy^1, \dots, x^n + iy^n]. \end{aligned}$$

2.2 Notes

Definition 2.0.1 (Topological Manifold).

A topological space M that satisfies

1. M is Hausdorff, i.e. points can be separated by open sets
2. M is second-countable, i.e. has a countable basis
3. M is locally Euclidean, i.e. every point has a neighborhood homeomorphic to an open subset \hat{U} of \mathbb{R}^n for some fixed n .

The last property says $p \in M \implies \exists U$ with $p \in U \subseteq M$, $\hat{U} \subseteq \mathbb{R}^n$, and a homeomorphism $\varphi : U \rightarrow \hat{U}$.

Note that second countability is primarily needed for existence of partitions of unity.

Exercise Show that in the last condition, \hat{U} can equivalently be required to be an open ball or \mathbb{R}^n itself.

Theorem 2.1 (*Topological Invariance of Dimension*).

Two nonempty topological manifolds of different dimensions can not be homeomorphic.

Exercise Show that in a Hausdorff space, finite subsets are closed and limits of convergent sequences are unique.

Exercise Show that subspaces and finite products of Hausdorff (resp. second countable) spaces are again Hausdorff (resp. second countable).

Thus any open subset of a topological manifold with the subspace topology is again a topological manifold.

Exercise Give an example of a connected, locally Euclidean Hausdorff space that is not second countable.

Definition 2.1.1 (Charts).

A chart on M is a pair (U, φ) where $U \subseteq M$ is open and $\varphi : U \rightarrow \hat{U}$ is a homeomorphism from U to $\hat{U} = \varphi(U) \subseteq \mathbb{R}^n$. If $p \in M$ and $\varphi(p) = 0 \in \hat{U}$, then the chart is said to be *centered* at p . Note that any chart about p can be modified to a chart (φ_1, \hat{U}_1) that is centered at p by defining $\varphi_1(x) = x - \varphi(p)$.



Fig. 1.2 A coordinate chart

U is the *coordinate domain* and φ is the *coordinate map*.

Note that we can write φ in components as $\varphi(p) = [x^1(p), \dots, x^n(p)]$ where each x^i is a map $x^i : U \rightarrow \mathbb{R}$. The component functions x^i are the *local coordinates* on U .

Shorthand notation: $[x^i] := [x^1, \dots, x^n]$.

Example 2.1 (Graphs of Continuous Functions).

Define

$$\Gamma(f) = \left\{ (x, y) \in \mathbb{R}^n \times \mathbb{R}^k \mid x \in U, y = f(x) \in \hat{U} \right\}.$$

This is a topological manifold since we can take $\varphi : \Gamma(f) \rightarrow U$ by restricting $\pi_1 : \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R}^n$ to the subspace $\Gamma(f)$. Projections are continuous, restrictions of continuous functions are continuous.

This is a homeomorphism because the map $g : x \mapsto (x, f(x))$ is continuous and $g \circ \pi_1 = \text{id}_{\mathbb{R}^n}$ is continuous with $\pi_1 \circ g = \text{id}_{\Gamma(f)}$. Note that $U \cong \Gamma(f)$, and thus $(U, \varphi) = (\Gamma(f), \varphi)$ is a single *global* coordinate chart, called the *graph coordinates* of f .

Thus graphs of continuous functions $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ are locally Euclidean?

Note that this works in greater generality:: “The same observation applies to any subset of \mathbb{R}^{n+k} by setting *any* k of the coordinates equal to some continuous function of the other n .”

Coordinates as numbers vs functions?

Example 2.2 (Spheres).

S^n is a subspace of \mathbb{R}^{n+1} and is thus Hausdorff and second-countable by exercise 2.2.




Fig. 1.3 Charts for \mathbb{S}^n

To see that it's locally Euclidean, take

$$U_i^+ := \left\{ [x^1, \dots, x^n] \in \mathbb{R}^{n+1} \mid x^i > 0 \right\} \quad \text{for } 1 \leq i \leq n+1$$

$$U_i^- := \left\{ [x^1, \dots, x^n] \in \mathbb{R}^{n+1} \mid x^i < 0 \right\} \quad \text{for } 1 \leq i \leq n+1.$$

Define

$$f : \mathbb{R}^n \longrightarrow \mathbb{R}^{\geq 0}$$

$$\mathbf{x} \mapsto \sqrt{1 - \|\mathbf{x}\|^2}.$$

Note that we immediately need to restrict the domain to $\mathbb{D}^n \subset \mathbb{R}^n$, where $\|x\|^2 \leq 1 \implies 1 - \|x\|^2 \geq 0$, to have a well-defined real function $f : \mathbb{D}^n \longrightarrow \mathbb{R}^{\geq 0}$.

Then (claim)

$$\begin{aligned} U_i^+ \cap S^n & \text{ is the graph of } x^i = f(x^1, \dots, \widehat{x^i}, \dots, x^{n+1}) \\ U_i^- \cap S^n & \text{ is the graph of } x^i = -f(x^1, \dots, \widehat{x^i}, \dots, x^{n+1}). \end{aligned}$$

This is because

$$\begin{aligned} \Gamma(x^i) &:= \{(\mathbf{x}, f(\mathbf{x})) \subseteq \mathbb{R}^n \times \mathbb{R}\} \\ &= \left\{ [x_1, \dots, \widehat{x^i}, \dots, x^{n+1}], f([x_1, \dots, \widehat{x^i}, \dots, x^{n+1}]) \subseteq \mathbb{R}^n \times \mathbb{R} \right\} \\ &= \left\{ [x_1, \dots, \widehat{x^i}, \dots, x^{n+1}], \left(1 - \sum_{\substack{j=1 \\ j \neq i}}^{n+1} (x^j)^2 \right)^{\frac{1}{2}} \subseteq \mathbb{R}^n \times \mathbb{R} \right\} \end{aligned}$$

and any vector in this set has norm satisfying

$$\|(\mathbf{x}, y)\|^2 = \sum_{\substack{j=1 \\ j \neq i}}^{n+1} (x^j)^2 + \left(1 - \sum_{\substack{j=1 \\ j \neq i}}^{n+1} (x^j)^2 \right) = 1$$

and is thus in S^n .

To see that any such point also has positive i coordinate and is thus in U_i^+ , we can rearrange (?) coordinates to put the value of f in the i th coordinate to obtain

$$\Gamma(x_i) = \left\{ [x^1, \dots, f(x^1, \dots, \widehat{x^i}, \dots, x^n), \dots, x^n] \right\}$$

Seems like f is always the *last* coordinate in the graph

and note that the square root only takes on positive values.

Thus each $U_i^\pm \cap S^n$ is the graph of a continuous function and thus locally Euclidean, and we can define chart maps

$$\begin{aligned} \varphi_i^\pm : U_i^\pm \cap S^n &\longrightarrow \mathbb{D}^n \\ [x^1, \dots, x^n] &\mapsto [x^1, \dots, \widehat{x^i}, \dots, x^{n+1}] \end{aligned}$$

yield $2(n+1)$ charts that are graph coordinates for S^n .

Example 2.3 (Projective Space).

Define \mathbb{RP}^n as the space of 1-dimensional subspaces of \mathbb{R}^{n+1} with the quotient topology determined by the map

$$\begin{aligned} \pi : \mathbb{R}^{n+1} \setminus \{0\} &\longrightarrow \mathbb{RP}^n \\ \mathbf{x} &\mapsto \text{span}_{\mathbb{R}} \{\mathbf{x}\}. \end{aligned}$$

How is this map a quotient map?

Notation: for $\mathbf{x} \in \mathbb{R}^{n+1} \setminus \{0\}$ write $[\mathbf{x}] := \pi(\mathbf{x})$, the line spanned by \mathbf{x} .

Define charts:

$$\tilde{U}_i := \left\{ \mathbf{x} \in \mathbb{R}^{n+1} \setminus \{0\} \mid x^i \neq 0 \right\}, \quad U_i = \pi(\tilde{U}_i) \subseteq \mathbb{RP}^n$$

and chart maps

$$\begin{aligned} \tilde{\varphi}_i : \tilde{U}_i &\longrightarrow \mathbb{R}^n \\ [x^1, \dots, x^{n+1}] &\mapsto \left[\frac{x^1}{x^i}, \dots, \widehat{x^i}, \dots, \frac{x^{n+1}}{x^i} \right]. \end{aligned}$$

Then (claim) this descends to a continuous map $\varphi_i : U_i \longrightarrow \mathbb{R}^n$ by the universal property of the quotient:

$$\begin{array}{ccc} \tilde{U}_i & & \\ \pi_U \downarrow & \searrow \tilde{\varphi}_i & \\ U_i & \xrightarrow{\varphi_i} & \mathbb{R}^n \end{array}$$

- The restriction $\pi_U : \tilde{U}_i \longrightarrow U_i$ of π is still a quotient map because $\tilde{U}_i = \pi_U^{-1}(U_i)$ where $U_i \subseteq \mathbb{RP}^n$ is open in the quotient topology and thus \tilde{U}_i is saturated.

Thus π_U sends saturated sets to open sets and is thus a quotient map.

- $\tilde{\varphi}_i$ is constant on preimages under π_U : fix $y \in U_i$, then $\pi_U^{-1}(\{y\}) = \left\{ \lambda \mathbf{y} \mid \lambda \in \mathbb{R} \setminus \{0\} \right\}$, i.e. the point $y \in \mathbb{RP}^n$ pulls back to every nonzero point on the line spanned by $\mathbf{y} \in \mathbb{R}^n$.

But

$$\begin{aligned} \tilde{\varphi}_i(\lambda \mathbf{y}) &= \varphi_i\left(\left[\lambda y^1, \dots, \lambda y^i, \dots, \lambda y^n\right]\right) \\ &= \left[\frac{\lambda y^1}{\lambda y^i}, \dots, \widehat{\lambda y^i}, \dots, \frac{\lambda y^{n+1}}{\lambda y^i}\right] \\ &= \left[\frac{y^1}{y^i}, \dots, \widehat{y^i}, \dots, \frac{y^{n+1}}{y^i}\right] \\ &= \tilde{\varphi}_i(\mathbf{y}). \end{aligned}$$

So this yields a continuous map

$$\varphi_i : U_i \longrightarrow \mathbb{R}^n.$$

We can now verify that φ is a homeomorphism since it has a continuous inverse given by

$$\begin{aligned} \varphi_i^{-1} : \mathbb{R}^n &\longrightarrow U_i \subseteq \mathbb{RP}^n \\ \mathbf{u} := [u^1, \dots, u^n] &\mapsto [u^1, \dots, u^{i-1}, \mathbf{1}, u^{i+1}, \dots, u^n]. \end{aligned}$$

It remains to check:

Exercise

1. The $n + 1$ sets U_1, \dots, U_{n+1} cover \mathbb{RP}^n .
2. \mathbb{RP}^n is Hausdorff
3. \mathbb{RP}^n is second-countable.

Exercise (1.6) Show that \mathbb{RP}^n is Hausdorff and second countable.

Exercise (1.7) Show that \mathbb{RP}^n is compact. (Hint: show that π restricted to S^n is surjective.)

Definition 2.1.2 (Topological Embedding).

A continuous map $f : X \rightarrow Y$ is a *topological embedding* iff it is injective and $\tilde{f} : X \rightarrow f(X)$ is a homeomorphism.

Example 2.4 (Product Manifolds).

Let $M := M_1 \times \dots \times M_k$ be a product of manifolds of dimensions n_1, \dots, n_k respectively. A product of Hausdorff/second-countable spaces is still Hausdorff/second-countable, so just need to check that it's locally Euclidean.

- Let $\mathbf{p} \in \prod_{i=1}^N M_i$, so $p_i \in M_i$
- Choose a chart (U_i, φ_i) with $p_i \in U_i$ and assemble a product map:

$$\Phi := \prod \varphi_i : \prod U_i \rightarrow \prod \mathbb{R}^{n_i} \cong \mathbb{R}^{\sum n_i} := \mathbb{R}^N.$$

- Claim: Φ is a homeomorphism onto its image in \mathbb{R}^N .
 - Each φ_i is a homeomorphism onto $\varphi_i(U_i)$ (by the definition of a chart on M_i)
 - It suffices to show that Φ^{-1} exists and is continuous, where

$$\Phi^{-1}(V) := \left(\prod \varphi_i \right)^{-1}(\prod V_i).$$

- Φ is a product of continuous functions and thus continuous.
- $\Phi^{-1} := \left(\prod \varphi_i \right)^{-1} = \prod \varphi_i^{-1}$, which are all assumed continuous since φ_i were homeomorphisms.

Example 2.5 (Torii).

$T^n := \prod_{i=1}^n S^1$ is a topological n -manifold.

Proposition 2.2.

Every topological manifold has a countable basis of precompact coordinate balls.

Todo list

Recommended problem	3
Recommended problem	3
Recommended problem	5
Recommended problem	5
Thus graphs of continuous functions $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$ are locally Euclidean?	6
Coordinates as numbers vs functions?	6
Seems like f is always the *last* coordinate in the graph	8
How is this map a quotient map?	8
Exercise	10