

Elliptic Curves

D. Zack Garza

March 14, 2020

Contents

1	Wednesday January 8	1
2	Mordell-Weil Groups	2
3	Monday January 13th	4
3.1	Every Abelian Group is a Class Group	4
3.2	Proof Sketch	4
3.2.1	Step 1	4
3.2.2	Step 2	4
3.2.3	Step 3	5
3.3	Mordell-Weil	6
4	Wednesday January 15th	6

List of Definitions

3.1	Definition – Replete	5
3.2	Definition – Weakly Replete	5
4.1	Definition	8

List of Theorems

3.1	Theorem – Claborn - Leedham - Green - Clark	4
3.2	Theorem – Mordell-Weil	6
4.1	Theorem – Height Descent	8

1 Wednesday January 8

Summary:

1. Mordell-Weil theorem
 - For elliptic curves over global fields (number fields, function fields, finite fields, etc)

- Proof uses Galois cohomology and height functions, essentially one proof!
 - Holds for abelian varieties, but more difficult (need an analog of height functions, i.e. an x -coordinate)
2. Height functions (possibly)
 3. Elliptic curves over \mathbb{Q}_p or complete discrete valuation fields (see Silverman for basics, possibly Chapter 5), particularly Tate curves
 4. Weil-Chatelet groups E/k related to $H^1(k; E)$ with coefficients in the elliptic curve
 5. Galois representation of E/k for $\text{char } k = 0$, for $\rho_n g_k \rightarrow \text{GL}(2, \mathbb{Z}/n\mathbb{Z})$ which leads to $\hat{\rho} : g_k \rightarrow \text{GL}(\hat{\mathbb{Z}})$.

2 Mordell-Weil Groups

Let E/k be an elliptic curve over a field k , i.e. a smooth, projective, geometrically integral curve of genus 1 with a k -rational point O .

Note: Silverman good for foundations, but assumes k is perfect! Here we'll assume k is arbitrary.

Remark: If k is not algebraically closed, such a point O may not exist.

By Riemann-Roch (easy computation) E embeds (non-canonically) into \mathbb{P}^2/k as a Weierstrass cubic

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6 \quad \Delta \neq 0.$$

This is a smoothness condition, and this equation has a k -rational point at infinity $[0 : 1 : 0]$. The line at infinity is a flex line (?), and so only intersects this curve at one point.

If $\text{char } k \neq 2, 3$ then $y^2 = x^3 + Ax + B$.

Every elliptic curve is given by a Weierstrass equation, although not in a unique way.

An amazing fact: The k -rational points $E(k)$ forms an abelian group with zero as the identity.

Proof:

1. Given any plane cubic C/k and an origin $O \in C(k)$, the chord and tangent process yields a group structure. Note that there is a symmetry in connecting rational points a, b with a line intersecting at another rational point c which is not present in most groups, so an additional inversion about O is needed to actually make this into a group. Proving associativity: difficult!
2. Look at $\text{Pic}^0 E$, the degree 0 divisors on E mod birational equivalence (?), which is equal to the degree 0 line bundles on E mod bundle isomorphism.

Exercise: Show there is a map $C(k) \rightarrow \text{Pic}^1 C$ given by sending p to its equivalence class; this is a bijection by Riemann-Roch (straightforward exercise).

We can then compose this with a map $\text{Pic}^1 \rightarrow \text{Pic}^0 C$ given by $D \mapsto D - [O]$, which decreases the degree by 1. This gives a map $\Phi : C(k) \rightarrow \text{Pic}^0 C$, just need to check that $\Phi(P \oplus Q) = \Phi(P) + \Phi(Q)$.

Check that the groups are independent of the k -rational point chosen, i.e. changing rational points yields isomorphic groups. So the group law itself does actually depend on the rational point, although the structure doesn't.

Exercise: Let $(E, O)/k$ be an elliptic curve and define $E^0 = E \setminus \{O\}$ the (nonsingular, integral) affine curve given by removing the point at infinity. Then the affine coordinate ring $k[E^0]$ is defined as $k[x, y]/(y^2 - x^3 - Ax - B)$, which is a Dedekind ring.

The interesting thing about Dedekind domains: the ideal class group! (i.e. the Picard group)

This has ideal class group $\text{Pic}[E^0]$, and one can show that

$$\begin{aligned} \text{Pic}^0 E &\longrightarrow \text{Pic}[E^0] \\ \sum_p n_p \deg(p)[p] &\mapsto \sum_{p \neq 0} n_p [p] = \prod_p p^{n_p} \end{aligned}$$

with the sum ranging over all closed points is an isomorphism.

Just note that the RHS can't have a point at infinity, so we just forget it. The isomorphism follows from some exact sequence with correction terms that vanish.

So the Mordell-Weil group of $E(k)$ is isomorphic to $\text{Pic}[E^0]$, the class group of a dedekind domain (?).

Definitions: Let G be a commutative group.

- G is a *class group* iff there exists a dedekind domain R such that $G \cong \text{Pic} R$.
- G is an (*elliptic*) *Mordell-Weil group* iff there exists a field k and an elliptic curve E/k such that $G \cong E(k)$.

Questions:

1. Which G are class groups?
2. Which G are Mordell-Weil groups?

An answer to question 1:

Theorem (Clayborn, 1966): Every commutative G is a class group.

Subsequent proofs: Leetham-Green (1972) and Clark (2008) following Rosen, and uses elliptic curves. See the end of Pete's Commutative Algebra notes!

An answer to question 2:

Consider E/\mathbb{C} , then $E(\mathbb{C}) \cong S^1 \times S^1$, so the torsion subgroup is $T(1) := (\mathbb{Q}/\mathbb{Z})^2 = \bigoplus_{\ell} (\mathbb{Q}_{\ell}/\mathbb{Z}_{\ell})^2$.

This in fact holds for any algebraically closed field of characteristic zero.

Fact: For any E/k , the Mordell-Weil group $E(k)$ is “ $T(1)$ -constrained”, i.e. $E(k)[\text{tors}] \hookrightarrow T(1)$.

Theorem (Clark, 2012): G is a Mordell-Weil group $\iff G$ is $T(1)$ -constrained.

Note: the analogous statement for abelian varieties, i.e being $T(g)$ constrained for some other genus $g \neq 1$, is open. Fixing $k = \mathbb{Q}$ still yields very interesting problems. Computing the rank and torsion subgroups is currently open, and the subject of modern research.

3 Monday January 13th

3.1 Every Abelian Group is a Class Group

Theorem 3.1 (*Claborn - Leedham - Green - Clark*).

Any commutative group is the class group of some Dedekind domain.

Also see: partial re-proof by Rosen that uses elliptic curves. This theorem: mostly a proof in commutative algebra, see end of Pete's commutative algebra notes.

3.2 Proof Sketch

Let E/k be an elliptic curve over a field.

3.2.1 Step 1

Note that $\text{End}_k(E) \cong_{\mathbb{Z}} \mathbb{Z}^{a(E)}$ where $a(E) \in \{1, 2, 4\}$.

Could be \mathbb{Z} as a \mathbb{Z} -module, could be an order in the imaginary quadratic field (e.g. a quaternion algebra)

There is a short exact sequence

$$0 \longrightarrow E(k) \longrightarrow E(k(E)) \longrightarrow \text{End}_K(E) \longrightarrow 0.$$

This splits because (as seen above), the RHS term is free and thus projective. So

$$E/k(E) \cong E(k) \oplus \mathbb{Z}^{a(E)}.$$

Note that $k(E)$ is an extension of E_k to $E_{k(E)}$ the field of rational functions over k ? (function field). To simplify, take $a(E) = 1$ and $E(k) = \{0\}$.

Taking $k = \mathbb{Q}$, this happens (probably, asymptotically) half of the time. It's easy to write down an elliptic curve that satisfies these conditions

Then $E/k(E) \cong \mathbb{Z}$.

Now pass to the field of rational functions over this field, taking $E(k(E)(E/k(E)))$. Then $k^2(E) := k(E)(E/k(E))$, and inductively define $k^n(E)$ by passing to function fields. So $E(k^n(E)) \cong \mathbb{Z}^n$.

So we can construct elliptic curves that have any free commutative group as their Mordell-Weil group.

3.2.2 Step 2

Loosely speaking, we'll iterate this process transfinitely. Then for any set S , there exists a field k and an elliptic curve E/k such that $E(k) \cong \bigoplus_S \mathbb{Z}$. We now want to introduce a process that allows passing to quotients. And $R := k[E^0]$ is the affine coordinate ring of E , remove the point at infinity (?).

3.2.3 Step 3

Let R be a Dedekind domain. Note it has a fraction field with a certain ideal class group. Let $W \subset \max\text{Spec } (R)$, then

$$R^W := \bigcap_{\mathfrak{p} \in \max\text{Spec } R \setminus W} R_{\mathfrak{p}}.$$

Then R^W is Dedekind (and every overring of a Dedekind domain is of this form) and $\max\text{Spec } (R^W) = \max\text{Spec } (R \setminus W)$.

Then

$$\text{Pic } R^W = \text{Pic } R / \langle [\mathfrak{p}] \mid \mathfrak{p} \in W \rangle.$$

Note that if $(A, +)$ is a commutative group, writing $A = \bigoplus_S \mathbb{Z}/H$, we have a Dedekind domain $R = k[E^0]$ such that $\text{Pic } R = \bigoplus_S \mathbb{Z}$.

Note: $\text{Pic } R$ is the class group.

Definition 3.1 (Replete).

A Dedekind domain R is **replete** iff every element of the class group $\text{Pic } R$ is the class group $[\mathfrak{p}]$ of some ideal $\mathfrak{p} \in \max\text{Spec } (R)$.

Is every ideal class the class of a prime ideal? For k a field, $R = \mathbb{Z}_k$. This follows from Chebotom (?) Density (most important theorem for arithmetic geometers!)

Definition 3.2 (Weakly Replete).

A Dedekind domain R is **weakly replete** iff every subgroup $H \subset \text{Pic } R$ is generated by classes of prime ideals.

Exercise (Easy) $K[E^0]$ is weakly replete, and an easy application of Riemann-Roch shows that if $0 \neq p \in E(k) = \text{Pic } k[E^0]$, then $[p] \in \text{Pic } k[E^0]$ is generated by a prime ideal.

Note: most applications of Riemann-Roch to elliptic curves are easy! In this case, it gives you an identification $E \cong \text{Pic } {}^1(E)$.

So there exists a subset $W \subset \max\text{Spec } k[E^0]$ such that $\langle [p] \mid p \in W \rangle = H$. Then

$$\text{Pic } k[E^0]^W \cong \bigoplus_S \mathbb{Z}/H \cong A.$$

■

Note that Dedekind domains don't have to be replete or even weakly replete. The class group of a Dedekind domain could be \mathbb{Z} , and the class of every prime ideal could be $1 \in \mathbb{Z}$

Proof (Claborn).

Start with an arbitrary Dedekind domain R and attach one that's replete.

Can ask for a similar result for abelian varieties, there are conjectures here, few clear results.

Need to get $\mathbb{Z}/(m) \times \mathbb{Z}/(n)$, since these occur as Mordell-Weil groups. Take a modular curve and a generic point. Look at universal elliptic curves over elliptic curves and take their Mordell-Weil groups (?)

If k is algebraically closed and $\text{char } k = p$, can't have $\mathbb{Z}(p) \times \mathbb{Z}/(p)$. Consider the p -primary torsion $E_k[p^\infty]$. It is zero iff E is supersingular (no points of order p). It is $\mathbb{Q}_p/\mathbb{Z}_p = \varinjlim_n \mathbb{Z}/(p^n)$ iff E is ordinary.

Can sometimes reduce to cases where $k = \mathbb{C}$ and do things analytically. ■

3.3 Mordell-Weil

Theorem 3.2 (Mordell-Weil).

Let k be a global field (extension of \mathbb{Q} or function field over \mathbb{F}_p) and E/k an elliptic curve. Then $E(k) \cong \mathbb{Z}^r \oplus T$ (by classification of abelian groups) where T is finite, and $T \cong \mathbb{Z}/(m) \oplus \mathbb{Z}/(n)$ for $m \mid n$. So T is generated by at most two elements.

Proof (3 steps).

Step 1: Weak Mordell-Weil theorem.

Take any $n \geq 2$ and $\text{char } k$ not dividing n . Show that $E(k)/nE(k)$ is finite.

Step 2: Define a height function $h : E(k) \rightarrow \mathbb{R}$ satisfying 3 properties (see next time). This is approximately a quadratic form.

Decompose at places of a number field, see Number Theory II.

Step 3: For any commutative group A , there is a notion of a height function

$$h : A \rightarrow \mathbb{R}.$$

Show the Height Descent Theorem: if A admits a height function and A/nA is finite for some $n \geq 2$, then A is finitely generated.

Also how you'd prove this theorem for abelian varieties, more difficulty defining h . ■

4 Wednesday January 15th

Recall that we're trying to prove the Mordell-Weil theorem. Let K be a global field, so it's the field of functions over some nice curve. Then the Mordell-Weil group $E(K)$ is finitely generated.

Step 1: The weak Mordell-Weil theorem for all $n \geq 2$ with $\text{char } k$ not dividing n , $E(k)/nE(k)$ is finite.

Step 2: Construction of a height function $h : E(K) \rightarrow \mathbb{R}$ that is "trying" to be a quadratic form.

Step 3 (Today): The Height Descent Theorem, i.e. if $(A, +)$ is a commutative group such that A/nA is finite for some $n \geq 2$ and it admits a height function $h : A \rightarrow \mathbb{R}$, then A is finitely generated.

Question: What does the weak Mordell-Weil group $E(K)/nE(K)$ tell us about $E(K)$?

Note that we'll inject this into a larger group, which we'll show is finite, but this isn't great for learning about the size.

Example 4.1.

Consider E/\mathbb{C} , then $E(\mathbb{C}) = S^1 \times S^1$ and $E(\mathbb{C})/nE(\mathbb{C}) = 0$, so the map $x \rightarrow nx$ is a surjective map and $E(K)$ is n -divisible here. In general, whenever $K = \overline{K}$ is algebraically closed, then $x \mapsto nx$ is again surjective and the weak Mordell-Weil group is trivial. So knowing this is small doesn't tell us much about $E(K)$ at all.

Example 4.2.

For E/\mathbb{R} , $E(\mathbb{R})$ is either S^1 (cubic with one real root, $\Delta = 0$) or $S^1 \times \mathbb{Z}/(2)$ (cubic with three real roots, $\Delta > 0$) are the two possible group structure.

Then

$$\begin{cases} 0 & n \text{ odd} \\ 0 & n \text{ even and } \Delta < 0. \\ \mathbb{Z}/(2) & n \text{ even and } \Delta > 0 \end{cases}.$$

Example 4.3.

Consider E/\mathbb{Q}_p , then for all $\ell \gg 0$ $E(\mathbb{Q}_p) \xrightarrow{[\ell]} E(\mathbb{Q}_p)$ with $E(\mathbb{Q}_p)/\ell E(\mathbb{Q}_p) = 0$ while $E(\mathbb{Q}_p)/pE(\mathbb{Q}_p)$ is not zero.

Note: here is an example of a Boolean space, that ends up being homeomorphic to a Cantor set.

Suppose $E(K)$ is finitely generated, so $E(K) \cong \mathbb{Z}^r \oplus T$ with T finite. Then knowing $E(K)/nE(K)$ gives an upper bound on r .

Example 4.4.

Take $n = 2$, then $E(K)/nE(K) \cong (\mathbb{Z}/(2))^s$ for some $s \in \mathbb{N}$. Then $(\mathbb{Z}^r \oplus T)/2(\mathbb{Z}^r \oplus T) \cong (\mathbb{Z}/(2))^r \oplus T/2T$ for $r \leq s$. Then either

- $r = 2$ and $E(K[2]) = (0)$.
- $r = 1$ and $E(K[2]) \cong \mathbb{Z}/(2)$,
- $r = 0$ and $E(K[2]) \cong (\mathbb{Z}/(2))^2$.

Note that we don't need the Mordell-Weil theorem to compute the torsion subgroups of $E(k)$. It is often easier to compute these directly. For all non-archimedean places v of K , $E(K_v)[\text{tors}]$ is finite (see Silverman?) and embeds into a number of finite things.

To compute $E(K_v)[\text{tors}]$,

1. Find $N \in \mathbb{Z}^+$ such that $E(k)[\text{tors}] \subset E[N]$.
 - Choose 2 different places v_0, v_1 of good reduction (from Weierstrass equation) with different residue characteristics $\ell_1 \neq \ell_2$

- Consider the map $E(K_{v_i})[\text{tors}] \rightarrow E(\mathbb{F}_{v_i})$
 - The kernel is a finite p_i -primary group.
 - Comes down to torsion and formal groups, see first course.
2. Compute $E[N](K)$ (several algorithms, just checking for rational points on a zero-dimensional variety?)

See division polynomials, can check for roots of polynomials over any global field. Easy to check for rational points on finite fields.

Suppose $E(K) \cong \mathbb{Z}^r \oplus T$ is finitely generated and we know $E(K)/nE(K)$ for some n and we know T . Then we explicitly know r .

See Tate Shafarevich group – important! But difficult, a piece of information that helps compute the rank (?).

Definition 4.1.

Fix $n \geq 2$. An n -height function on $(A, +)$ is a map $h : A \rightarrow \mathbb{R}$ satisfying

1. For all $R \geq 0$, the set $h^{-1}(-\infty, R)$ is finite.
2. For all $Q \in A$, there exists a $C_2 = C_2(A, Q)$ such that for all $P \in A$, $h(P + Q) \leq 2h(P) + C_2$.
(?)
3. There exists a $C_3 = C_3(A, n)$ such that for all $P \in A$, $h(nP) \geq n^2h(P) - C_3$

Note: (3) would be an equality for an honest quadratic function, so this deviates in a controlled way.

Theorem 4.1 (Height Descent).

Let $(A, +)$ be a commutative group with an n -height function $h : (A, +) \rightarrow \mathbb{R}$. If A/nA is finite, then A is finitely generated.

Proof: Let r be the size of A/nA . Choose coset representatives Q_1, \dots, Q_r of nA in A . Let $p \in A$ and define a sequence $\{P_k\}_{k=0}^\infty$ in A by $P_0 = P$ and for $k \geq 1$, choose P_k such that $P_{k-1} = nP_k + Q_{i_k}$.

Then for all $k \in \mathbb{Z}^+$, it's true that $P = n^k P_k + \sum_{j=1}^k n^{j-1} Q_{i_j}$.

Claim 1.

There exists a constant $c > 0$ depending only on A, n such that for all $P \in A$, there exists a $K = K(P)$ such that for all $k \geq K$, we have $h(P_k) \leq 0$.

Note that this is sufficient – if so, A is generated by $\{Q_1, \dots, Q_r\} \cup h^{-1}((-\infty, C])$, which are both finite.

Next time: proof of claim.

Note: similar setup goes through for abelian varieties, see Néron–Tate height canonical height, which yields an honest “quadratic form”.