# **Algebra**

### D. Zack Garza

## August 15, 2019

#### **Contents**

1		Lecture 1 (Thu 15 Aug 2019)		
	1.1	Cyclic	Groups	2
	1.2	Homo	morphisms	3
	1.3	Direct	Products	3
	1.4	Finite	ly Generated Abelian Groups	3
	1.5	Funda	mental Homomorphism Theorem	4
		1.5.1	The First Homomorphism Theorem	4
		1.5.2	The Second Theorem	4
2	Lect	cure 2		4

# 1 Lecture 1 (Thu 15 Aug 2019)

**Definition 1.** >  $\{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$  A **group** is an ordered pair  $(G, \cdot : G \times G \to G)$  where G is a set and  $\cdot$  is a binary operation, which satisfies the following axioms:

- Associativity:  $(g_1g_2)g_3 = g_1(g_2g_3)$ ,
- Identity:  $\exists e \in G \ni ge = eg = g$ ,
- Inverses:  $g \in G \implies \exists h \in G \ni gh = gh = e$ .

> {{{}}}{{{}}}}{{{{}}}}

Example 1. > {{{}}}{{{}}}{{{}}}{{{}}}{{{}}}{{{}}}}   
 (GL(n, 
$$\mathbb{R}), \times) = {A \in Mat_n \ni det(A) \neq 0} - (S_n, \circ)$$

> {{{}}}}{{{{}}}}

**Definition 2.**  $> \{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}$  A subset  $S \subseteq G$  is a **subgroup** of G iff

- $1. \ s_1, s_2 \in S \implies s_1 s_2 \in S$
- $2. e \in S$
- $3. \ s \in S \implies s^{-1} \in S$
- > {{{}}}}{{{{}}}}

We denote such a subgroup  $S \leq G$ .

Examples of subgroups:

- $(\mathbb{Z},+) \leq (\mathbb{Q},+)$
- $SL(n,\mathbb{R}) \leq GL(n,\mathbb{R})$ , where  $SL(n,\mathbb{R}) = \{A \in GL(n,\mathbb{R}) \ni \det(A) = 1\}$

#### 1.1 Cyclic Groups

**Definition 3.** >  $\{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}$  A group G is **cyclic** iff G is generated by a single element.

> {{{}}}}{{{}}}}

**Exercise 1.**  $> \{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$  Show  $\langle g \rangle = \{g^n \ni n \in \mathbb{Z}\} \cong \bigcap \{H \leq G \ni g \in H\}.$ 

> {{{}}}}{{{{}}}}

**Theorem 1.** > {{{}}}{{{}}}{{{}}}{{{}}}}{{{}}}. So G(g).

- If  $|G| = \infty$ , then  $G \cong \mathbb{Z}$ .
- If  $|G| = n < \infty$ , then  $G \cong \mathbb{Z}_n$ .
- > {{{}}}}{{{{}}}}

**Definition 4.** >  $\{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$  Let  $H \leq G$ , and define a **right coset of** G by  $aH = \{ah \ni H \in H\}$ . A similar definition can be made for **left cosets**.

Then  $aH = bH \iff b^{-1}a \in G \text{ and } Ha = Hb \iff ab^{-1} \in H.$ 

Some facts:

- Cosets partition H, i.e.  $b \notin H \implies aH \cap bH = \{e\}$ .
- |H| = |aH| = |Ha| for all  $a \in G$ .

**Theorem 2** (Lagrange). >  $\{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$  If G is a finite group and  $H \leq G$ , then  $|H| \mid |G|$ .

**Definition 5.** >  $\{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$  A subgroup  $N \leq G$  is **normal** iff gN = Ng for all  $g \in G$ , or equivalently  $gNg^{-1} \subseteq N$ . I denote this  $N \subseteq G$ .

When  $N \leq G$ , the set of left/right cosets of N themselves have a group structure. So we define

$$G/N = \{gN \ni g \in G\}$$
 where  $(g_1N)(g_2N) = (g_1g_2)N$ .

Given  $H, K \leq G$ , define  $HK = \{hk \ni h \in H, k \in K\}$ . We have a general formula,

$$|HK| = \frac{|H||K|}{|H \cap K|}.$$

#### 1.2 Homomorphisms

**Definition 6.** >  $\{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$  Let G, G' be groups, then  $\varphi: G \to G'$  is a **homomorphism** if  $\varphi(ab) = \varphi(a)\varphi(b)$ .

Example 2.  $> \{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$ 

- $\exp: (\mathbb{R}, +) \to (\mathbb{R}^{>0}, \cdot)$  where  $\exp(a+b) = e^{a+b} = e^a e^b = \exp(a) \exp(b)$ .
- det:  $(GL(n,\mathbb{R}),\times) \to (\mathbb{R}^{\times},\times)$  where  $\det(AB) = \det(A)\det(B)$ .
- Let  $N \subseteq G$  and  $\varphi G \to G/N$  given by  $\varphi(g) = gN$ .
- Let  $\varphi: \mathbb{Z} \to \mathbb{Z}_n$  where  $\phi(g) = [g] = g \mod n$  where  $\mathbb{Z}_n \cong \mathbb{Z}/n\mathbb{Z}$

**Definition 7.** >  $\{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$  Let  $\varphi: G \to G'$ . Then  $\varphi$  is a **monomorphism** iff it is injective, an **epimorphism** iff it is surjective, and an **isomorphism** iff it is bijective.

#### 1.3 Direct Products

Let  $G_1, G_2$  be groups, then define

$$G_1 \times G_2 = \{(g_1, g_2) \ni g_1 \in G, g_2 \in G_2\}$$
 where  $(g_1, g_2)(h_1, h_2) = (g_1h_1, g_2, h_2)$ .

We have the formula  $|G_1 \times G_2| = |G_1||G_2|$ .

#### 1.4 Finitely Generated Abelian Groups

**Definition 8.** > {{{}}}{{{}}}{{{}}}{{{}}}}{{{}}} We say a group is **abelian** if G is commutative, i.e.  $g_1, g_2 \in G \implies g_1g_2 = g_2g_1$ .

This generalizes the notion of a cyclic group, where we can simply intersect all of the subgroups that contain the  $g_i$  to define it.

We know what cyclic groups look like – they are all isomorphic to  $\mathbb{Z}$  or  $\mathbb{Z}_n$ . So now we'd like a structure theorem for abelian finitely generated groups.

**Theorem 3.**  $> \{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}$  Let G be a finitely generated abelian group. Then

$$G \cong \mathbb{Z}^r \times \prod_{i=1}^s \mathbb{Z}_{p_i^{\alpha_i}}$$

for some finite  $r, s \in \mathbb{N}$  and  $p_i$  are (not necessarily distinct) primes.


**Example 3.**  $> \{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$  Let G be a finite abelian group of order 4. Then  $G \cong \mathbb{Z}_4$  or  $\mathbb{Z}_2^2$ , which are not isomorphic because every element in  $\mathbb{Z}_2^2$  has order 2 where  $\mathbb{Z}_4$  contains an element of order 4.

#### 1.5 Fundamental Homomorphism Theorem

Let  $\varphi: G \to G'$  be a group homomorphism and define  $\ker \varphi = \{g \in G \ni \varphi(g) = e'\}$ .

#### 1.5.1 The First Homomorphism Theorem

**Theorem 4.** > {{{}}}{{{}}}{{{}}}{{{}}}}{{{}}}} {{{}}}} There exists a map  $\varphi': G/\ker \varphi \to G'$  such that the following diagram commutes:



That is,  $\varphi = \varphi' \circ \eta$ , and  $\varphi'$  is an isomorphism onto its image, so  $G/\ker \varphi = \operatorname{im} \varphi$ . This map is give by  $\varphi'(g(\ker \varphi)) = \varphi(g)$ .

**Exercise 2.**  $> \{\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}$  Check that  $\varphi$  is well-defined.

#### 1.5.2 The Second Theorem

**Theorem 5.**  $> \{:\{\{\}\}\}\{\{\{\}\}\}\{\{\{\}\}\}\}\{\{\{\}\}\}\}$  Let  $K, N \leq G$  where  $N \subseteq G$ . Then

$$\frac{K}{N \cap K} \cong \frac{NK}{N}$$

#### 2 Lecture 2