

Problem Set 10

D. Zack Garza

December 1, 2019

Contents

1 Problem 1	1
2 Problem 2	1
3 Problem 3	2
4 Problem 4	2
5 Problem 5	2
5.1 Part 1	2

1 Problem 1

Let ϕ be an n -form. It suffices to show these statements for $n = 2$.

\implies : Suppose ϕ is alternating, then $\phi(b, b) = 0$ for all $b \in B$.

Letting $a, b \in B$ be arbitrary, we then have

$$\begin{aligned}\phi(a + b, a + b) &= \phi(a, a + b) + \phi(b, a + b) \\ &= \phi(a, a) + \phi(a, b) + \phi(b, a) + \phi(b, b) \\ &= \phi(a, b) + \phi(b, a) \\ &\implies \phi(a, b) = -\phi(b, a),\end{aligned}$$

which shows that ϕ is skew-symmetric.

\Leftarrow Suppose ϕ is skew-symmetric, so $\phi(a, b) = -\phi(b, a)$ for all $a, b \in B$. Then $\phi(b, b) = -\phi(b, b)$ by transposing the terms, which says that $\phi(b, b) = 0$ for all $b \in B$ and thus ϕ is alternating.

2 Problem 2

Let $f(x) = \det(P + xQ) \in R[x]$, then f is a polynomial in x which is not identically zero.

To see that $f \neq 0$, we can use that fact that P is invertible to evaluate $f(0) = \det(P) \neq 0$.

We can now note that f has finite degree, and thus finitely many zeroes in R .

3 Problem 3

Letting $k[x] \curvearrowright_\phi E$ to yield a $k[x]$ -module structure on E and take an invariant factor decomposition,

$$E = E_1 \oplus E_2 \oplus \cdots \oplus E_t, \quad E_i = \frac{k[x]}{(q_i)}, \quad q_1 \mid q_2 \mid \cdots \mid q_t$$

where $E_i = k[x]/(q_i)$. Then $q_t = q$, the minimal polynomial of E .

In particular, E_t is a ϕ -invariant subspace of E , and if $\deg q_t = m$, then E_t is in fact an m -dimensional cyclic module with basis $\{\mathbf{v}, \phi(\mathbf{v}), \phi^2(\mathbf{v}), \dots, \phi^{m-1}(\mathbf{v})\}$ for some $\mathbf{v} \in E_t$.

But since $E_t \leq E$ is a subspace, we have

$$m = \deg q(x) = \deg q_t(x) = \dim E_t \leq \dim E.$$

4 Problem 4

\Rightarrow : Suppose $A \sim D$ where D is diagonal. Then $JCF(A) = JCF(D) = D$, which means that every Jordan block of A has size exactly 1.

Since the elementary divisors of A are precisely the minimal polynomials of the Jordan blocks of A , and the minimal polynomial of any 1×1 matrix $[a_{ij}]$ is given by the linear polynomial $x - a_{ij}$, every elementary divisor of A must be linear.

\Leftarrow : Suppose all of the elementary divisors of A are linear. Every elementary divisor is the minimal polynomial of a Jordan block of A , and so if we write $JCF(A) = \bigoplus M_i$, then the minimal polynomial of each M_i is linear.

Supposing that M_i has minimal polynomial $p_i(x) = x - c$ for some scalar c , we have

$$p_i(M_i) = 0 \implies M_i - cI_n = 0 \implies M_i = cI_n,$$

which shows that M_i is a diagonal matrix with only c on its diagonal.

But if every Jordan block of A is diagonal, then $JCF(A) = D$ is diagonal and $A \sim D$.

5 Problem 5

5.1 Part 1

We'll use the fact that the minimal polynomial q is the invariant factor of highest degree, and so every other invariant factor must divide q .

Moreover, $RCF(A) = C_1 \oplus C_2 \oplus \cdots \oplus C_k$ where each C_i is the companion matrix of the i th invariant factor if we write $V \cong \bigoplus_{i=1}^k k[x]/(a_i)$. So it suffices to determine all of the possible distinct combinations of invariant factors.

We can restrict this list by noting that the characteristic polynomial satisfies $\chi_A(x) = \prod a_i$, and in particular, $\deg \chi_A(x) = 6$. Noting that $\deg q(x) = 3$, the degrees of the remaining invariant factors must sum to 3.

These are:

- $a_1 = (x - 2), a_2 = q(x)$
- $a_1 = (x - 2), a_2 = (x - 2)^2, a_3 = q(x)$
- $a_1 = (x - 2), a_2 = (x - 2)(x - 3), a_3 = q(x)$
- $a_1 = (x - 3), a_2 = q(x)$
- $a_1 = (x - 3), a_2 = (x - 2)(x - 3), a_3 = q(x)$