

# Moduli Spaces

D. Zack Garza

March 18, 2020

## Contents

<b>1</b>	<b>Preface</b>	<b>2</b>
<b>2</b>	<b>Thursday January 9th</b>	<b>2</b>
2.1	Representability . . . . .	2
2.2	Projective Space . . . . .	6
<b>3</b>	<b>Tuesday January 14th</b>	<b>9</b>
<b>4</b>	<b>Thursday January 16th</b>	<b>15</b>
4.1	Subfunctors . . . . .	15
4.2	Actual Geometry: Hilbert Schemes . . . . .	16
4.2.1	Proof . . . . .	18
<b>5</b>	<b>Thursday January 23</b>	<b>19</b>
5.0.1	Hypersurfaces . . . . .	20
5.0.2	Example: Twisted Cubics . . . . .	22
<b>6</b>	<b>Tuesday January 28th</b>	<b>24</b>
<b>7</b>	<b>Thursday January 30th</b>	<b>31</b>

## List of Definitions

2.2.1	Definition – Moduli Functor . . . . .	3
2.2.2	Definition – Moduli Space . . . . .	3
3.0.1	Definition – Equalizer . . . . .	9
3.0.2	Definition – Coequalizer . . . . .	10
3.0.3	Definition – Zariski Sheaf . . . . .	11
3.0.4	Definition – Subfunctors, Open/Closed Functors . . . . .	12
3.0.5	Definition – Open Covers . . . . .	12
4.0.1	Definition – Hilbert Functor . . . . .	16
4.0.2	Definition – Flatness . . . . .	16
4.3.1	Definition – Hilbert Polynomial Subfunctor . . . . .	19
7.0.1	Definition . . . . .	31

## List of Theorems

2.1	Theorem – Yoneda . . . . .	3
2.3	Proposition . . . . .	7
3.1	Proposition . . . . .	13
4.1	Proposition . . . . .	17
4.2	Proposition – Modified Characterization of Flatness for Sheaves . . . . .	18
4.4	Theorem – Grothendieck . . . . .	19

## 1 Preface

These are notes live-tex'd from a course in Moduli Spaces taught by Ben Bakker at the University of Georgia in Spring 2020. Any errors or inaccuracies are almost certainly my own.

D. Zack Garza, March 18, 2020

## 2 Thursday January 9th

Some references:

- Course Notes
- Hilbert schemes/functors of points: Notes by Stromme
  - Slightly more detailed: Nitsure, . . . Hilbert schemes, Fundamentals of Algebraic Geometry
  - Mumford, Curves on Surfaces
- Harris-Harrison, Moduli of Curves (chatty and less rigorous)

### 2.1 Representability

Last time: Fix an  $S$ -scheme, i.e. a scheme over  $S$ .

Then there is a map

$$\begin{aligned}\mathrm{Sch}/S &\longrightarrow \mathrm{Fun}(\mathrm{Sch}/S^{\mathrm{op}}, \mathrm{Set}) \\ x &\mapsto h_x(T) = \mathrm{hom}_{\mathrm{Sch}/S}(T, x).\end{aligned}$$

where  $T' \xrightarrow{f} T$  is given by

$$\begin{aligned}h_x(f) : h_x(T) &\longrightarrow h_x(T') \\ (T \mapsto x) &\mapsto \text{triangles of the form}\end{aligned}$$

$$\begin{array}{ccc} T' & \xrightarrow{\quad} & X \\ & \searrow & \nearrow \\ & T & \end{array} .$$

**Theorem 2.1 (Yoneda).**

$$\mathrm{hom}_{\mathrm{Fun}}(h_x, F) = F(x).$$

**Corollary 2.2.**

$$\mathrm{hom}_{\mathrm{Sch}/S}(x, y) \cong \mathrm{hom}_{\mathrm{Fun}}(h_x, h_y).$$

**Definition 2.2.1** (Moduli Functor).

A **moduli functor** is a map

$$\begin{aligned} F : (\mathrm{Sch}/S)^{\mathrm{op}} &\longrightarrow \mathrm{Set} \\ F(x) &= \text{"Families of something over } x\text{"} \\ F(f) &= \text{"Pullback"}. \end{aligned}$$

**Definition 2.2.2** (Moduli Space).

A **moduli space** for that “something” appearing above is an  $M \in \mathrm{Obj}(\mathrm{Sch}/S)$  such that  $F \cong h_M$ .

Now fix  $S = \mathrm{Spec}(k)$ .

$h_m$  is the functor of points over  $M$ .

**Remark (1)**  $h_m(\mathrm{Spec}(k)) = M(\mathrm{Spec}(k)) \cong \text{“families over } \mathrm{Spec}(k)\text{”} = F(\mathrm{Spec}(k))$ .

**Remark (2)**  $h_M(M) \cong F(M)$  are families over  $M$ , and  $\mathrm{id}_M \in \mathrm{Mor}_{\mathrm{Sch}/S}(M, M) = \xi_{\mathrm{Univ}}$  is the universal family.

Every family is uniquely the pullback of  $\xi_{\mathrm{Univ}}$ . This makes it much like a classifying space.

For  $T \in \mathrm{Sch}/S$ ,

$$\begin{aligned} h_M &\xrightarrow{\cong} F \\ f \in h_M(T) &\xrightarrow{\cong} F(T) \ni \xi = F(f)(\xi_{\mathrm{Univ}}). \end{aligned}$$

where  $T \xrightarrow{f} M$  and  $f = h_M(f)(\mathrm{id}_M)$ .

**Remark (3)** If  $M$  and  $M'$  both represent  $F$  then  $M \cong M'$  up to unique isomorphism.

$$\xi_M \qquad \qquad \xi_{M'}$$

$$M \xrightarrow{f} M'$$

$$M' \xrightarrow{g} M$$

$$\xi_{M'} \qquad \qquad \xi_M$$

which shows that  $f, g$  must be mutually inverse by using universal properties.

**Example 2.1.**

A length 2 subscheme of  $\mathbb{A}_k^1$  (??) then

$$F(S) = \{V(x^2 + bx + c)\} \subset \mathbb{A}'_5$$

where  $b, c \in \mathcal{O}_s(s)$ , which is functorially bijective with  $\{b, c \in \mathcal{O}_s(s)\}$  and  $F(f)$  is pullback.


Then  $F$  is representable by  $\mathbb{A}_k^2(b, c)$  and the universal object is given by

$$V(x^2 + bx + c) \subset \mathbb{A}^1(?) \times \mathbb{A}^2(b, c)$$

where  $b, c \in k[b, c]$ .

Moreover,  $F'(S)$  is the set of effective Cartier divisors in  $\mathbb{A}'_5$  which are length 2 for every geometric fiber.  $F''(S)$  is the set of subschemes of  $\mathbb{A}'_5$  which are length 2 on all geometric fibers. In both cases,  $F(f)$  is always given by pullback.

Problem:  $F''$  is not a good moduli functor, as it is not representable. Consider  $\text{Spec } k[\varepsilon]$ .



$F$	✓	x	x
$F'$	✓	x	x
$F''$	✓	✓	✓

$$\begin{array}{ccccc}
 \mathrm{Spec} k & \xrightarrow{i} & \mathrm{Spec} k[\varepsilon] & & \\
 & & & \nearrow & \\
 & & & & = F'(\mathrm{Spec} k) \\
 F(\mathrm{Spec} k[\varepsilon]) & \xrightarrow{F(i)} & F(\mathrm{Spec} k) & & \searrow \\
 & & & & = F''(\mathrm{Spec} k) \\
 \uparrow \subset & & \uparrow \in & & \\
 T_p F','' & & P = V(x(x-1)) & & 
 \end{array}$$

We think of  $T_p F',''$  as the tangent space at  $p$ .

If  $F$  is representable, then it is actually the Zariski tangent space.

$$\begin{array}{ccc}
 M(\mathrm{Spec} k[\varepsilon]) & \longrightarrow & M(\mathrm{Spec} k) \\
 \uparrow \subset & & \uparrow \subset \\
 T_p M & \longrightarrow & p \\
 & & \\
 & \mathrm{Spec} k & \\
 & \swarrow \quad \searrow & \\
 \mathrm{Spec} k[\varepsilon] & \longrightarrow & \mathrm{Spec} \mathcal{O}_{M,p} \subset M
 \end{array}$$

$$\begin{array}{ccc}
 & & k \\
 & \nearrow & \uparrow \\
 \mathcal{O}_{M,p} & \longrightarrow & k[\varepsilon] \\
 \uparrow & & \uparrow \\
 \mathfrak{m}_p & & (\varepsilon) \\
 \uparrow & & \uparrow \\
 \mathfrak{m}_p^2 & & 0
 \end{array}$$

Moreover,  $T_p M = (\mathfrak{m}_p / \mathfrak{m}_p^2)^\vee$ , and in particular this is a  $k$ -vector space. To see the scaling structure, take  $\lambda \in k$ .

$$\begin{aligned}\lambda : k[\varepsilon] &\longrightarrow k[\varepsilon] \\ \varepsilon &\mapsto \lambda\varepsilon\end{aligned}$$

$$\lambda^* : \operatorname{Spec} (k[\varepsilon]) \longrightarrow \operatorname{Spec} (k[\varepsilon])$$

$$\begin{aligned}\lambda : M(\operatorname{Spec} (k[\varepsilon])) &\longrightarrow M(\operatorname{Spec} (k[\varepsilon])) \\ \cup &\qquad \qquad \cup \\ T_p M &\longrightarrow T_p M.\end{aligned}$$

**Conclusion:** If  $F$  is representable, for each  $p \in F(\operatorname{Spec} k)$  there exists a unique point of  $T_p F$  that are invariant under scaling.

1. If  $F, F', G \in \operatorname{Fun}((\operatorname{Sch}/S)^{\operatorname{op}}, \operatorname{Set})$ , there exists a fiber product

$$\begin{array}{ccc} F \times_G F' & \xrightarrow{\quad \quad \quad} & F' \\ \downarrow & & \downarrow \\ F & \longrightarrow & G \end{array}$$

where

$$(F \times_G F')(T) = F(T) \times_{G(T)} F'(T).$$

2. This works with the functor of points over a fiber product of schemes  $X \times_T Y$  for  $X, Y \longrightarrow T$ , where

$$h_{X \times_T Y} = h_X \times_{h_t} h_Y.$$

3. If  $F, F', G$  are representable, then so is the fiber product  $F \times_G F'$ .
4. For any functor

$$F : (\operatorname{Sch}/S)^{\operatorname{op}} \longrightarrow \operatorname{Set},$$

for any  $T \xrightarrow{f} S$  there is an induced functor

$$\begin{aligned}F_T : (\operatorname{Sch}/T) &\longrightarrow \operatorname{Set} \\ x &\mapsto F(x).\end{aligned}$$

5.  $F$  is representable by  $M/S$  implies that  $F_T$  is representable by  $M_T = M \times_S T/T$ .

## 2.2 Projective Space

Consider  $\mathbb{P}_{\mathbb{Z}}^n$ , i.e. “rank 1 quotient of an  $n + 1$  dimensional free module”.

**Proposition 2.3.**
 $\mathbb{P}_{\mathbb{Z}}^n$  represents the following functor

$$F : \text{Sch}^{\text{op}} \longrightarrow \text{Set}$$

$$F(S) = \mathcal{O}_S^{n+1} \longrightarrow L \longrightarrow 0 / \sim .$$

where  $\sim$  identifies diagrams of the following form:

$$\begin{array}{ccccc} \mathcal{O}_S^{n+1} & \longrightarrow & L & \longrightarrow & 0 \\ \parallel & & \downarrow \cong & & \\ \mathcal{O}_S^{n+1} & \longrightarrow & M & \longrightarrow & 0 \end{array}$$

and  $F(f)$  is given by pullbacks.

**Remark**  $\mathbb{P}_S^n$  represents the following functor:

$$F_S : (\text{Sch}/S)^{\text{op}} \longrightarrow \text{Set}$$

$$T \mapsto F_S(T) = \left\{ \mathcal{O}_T^{n+1} \longrightarrow L \longrightarrow 0 \right\} / \sim .$$

This gives us a cleaner way of gluing affine data into a scheme.

*Proof (of Proposition).*

Note:  $\mathcal{O}^{n+1} \longrightarrow L \longrightarrow 0$  is the same as giving  $n+1$  sections  $s_1, \dots, s_n$  of  $L$ , where surjectivity ensures that they are not the zero section.

So

$$F_i(S) = \left\{ \mathcal{O}_S^{n+1} \longrightarrow L \longrightarrow 0 \right\} / \sim ,$$

with the additional condition that  $s_i \neq 0$  at any point.

There is a natural transformation  $F_i \longrightarrow F$  by forgetting the latter condition, and is in fact a subfunctor.

$F \leq G$  is a subfunctor iff  $F(s) \hookrightarrow G(s)$ .


**Claim:** It is enough to show that each  $F_i$  and each  $F_{ij}$  are representable, since we have natural transformations:

$$\begin{array}{ccc} F_i & \longrightarrow & F \\ \uparrow & & \uparrow \\ F_{ij} & \longrightarrow & F_j \end{array}$$


and each  $F_{ij} \longrightarrow F_i$  is an open embedding (on the level of their representing schemes).

**Example .**

For  $n = 1$ , we can glue along open subschemes



For  $n = 2$ , we get overlaps of the following form:



This claim implies that we can glue together  $F_i$  to get a scheme  $M$ . We want to show that  $M$  represents  $F$ .  $F(s)$  (LHS) is equivalent to an open cover  $U_i$  of  $S$  and sections of  $F_i(U_i)$  satisfying the gluing (RHS).

Going from LHS to RHS isn't difficult, since for  $\mathcal{O}_S^{n+1} \rightarrow L \rightarrow 0$ ,  $U_i$  is the locus where  $s_i \neq 0$  and by surjectivity, this gives a cover of  $S$ .

RHS to LHS comes from gluing.

■

*Proof (of Claim).*

$$F_i(S) = \left\{ \mathcal{O}_S^{n+1} \rightarrow L \cong \mathcal{O}_S \rightarrow 0, s_i \neq 0 \right\},$$

but there are no conditions on the sections other than  $s_i$ .

So specifying  $F_i(S)$  is equivalent to specifying  $n - 1$  functions  $f_1 \cdots \widehat{f_i} \cdots f_n \in \mathcal{O}_S(s)$  with  $f_k \neq 0$ . We know this is representable by  $\mathbb{A}^n$ .

We also know  $F_{ij}$  is obviously the same set of sequences, where now  $s_j \neq 0$  as well, so we need to specify  $f_0 \cdots \widehat{f_i} \cdots f_j \cdots f_n$  with  $f_j \neq 0$ . This is representable by  $\mathbb{A}^{n-1} \times \mathbb{G}_m$ , i.e.  $\text{Spec } k[x_1, \dots, \widehat{x_i}, \dots, x_n, x_j^{-1}]$ . Moreover,  $F_{ij} \hookrightarrow F_i$  is open.

What is the compatibility we are using to glue? For any subset  $I \subset \{0, \dots, n\}$ , we can define

$$F_I = \left\{ \mathcal{O}_S^{n+1} \rightarrow L \rightarrow 0, s_i \neq 0 \text{ for } i \in I \right\} = \bigtimes_{i \in I} F_i,$$

and  $F_I \rightarrow F_J$  when  $I \supset J$ .

■



### 3 Tuesday January 14th

Last time: Representability of functors, and specifically projective space  $\mathbb{P}_{\mathbb{Z}}^n$  constructed via a functor of points, i.e.

$$h_{\mathbb{P}_{\mathbb{Z}}^n} : \mathbb{P}_{\mathbb{Z}}^n \text{Sch}^{\text{op}} \longrightarrow \text{Set}$$

$$s \mapsto \mathbb{P}_{\mathbb{Z}}^n(s) = \left\{ \mathcal{O}_s^{n+1} \longrightarrow L \longrightarrow 0 \right\}.$$

for  $L$  a line bundle, up to isomorphisms of diagrams:

$$\begin{array}{ccccc} \mathcal{O}_s^{n+1} & \longrightarrow & L & \longrightarrow & 0 \\ \parallel & & \downarrow \cong & & \\ \mathcal{O}_s^{n+1} & \longrightarrow & M & \longrightarrow & 0 \end{array}$$

That is, line bundles with  $n + 1$  sections that globally generate it, up to isomorphism.

The point was that for  $F_i \subset \mathbb{P}_{\mathbb{Z}}^n$  where

$$F_i(s) = \left\{ \mathcal{O}_s^{n+1} \longrightarrow L \longrightarrow 0 \mid s_i \text{ is invertible} \right\}$$

are representable and can be glued together, and projective space represents this functor.

**Remark** Because projective space represents this functor, there is a universal object:

$$\begin{array}{ccccc} \mathcal{O}_{\mathbb{P}_{\mathbb{Z}}^n}^{n+1} & \longrightarrow & L & \longrightarrow & 0 \\ & & \parallel & & \\ & & \mathcal{O}_{\mathbb{P}_{\mathbb{Z}}^n}(1) & & \end{array}$$


and other functors are pullbacks of the universal one. (Moduli Space)

**Exercise** Show that  $\mathbb{P}_{\mathbb{Z}}^n$  is proper over  $\text{Spec } \mathbb{Z}$ . Use the evaluative criterion, i.e. there is a unique lift

$$\begin{array}{ccc} \text{Spec } k & \longrightarrow & \mathbb{P}_{\mathbb{Z}}^n \\ \downarrow & \nearrow \text{dashed} & \downarrow \\ \text{Spec } R & \longrightarrow & \text{Spec } \mathbb{Z} \end{array}$$

**Definition 3.0.1** (Equalizer).

For a category  $C$ , we say a diagram  $X \longrightarrow Y \rightrightarrows Z$  is an *equalizer* iff it is universal with respect to the property:




Note that  $X$  is the universal object here.

**Example 3.1.**

For sets,  $X = \{y \mid f(y) = g(y)\}$  for  $Y \xrightarrow{f,g} Z$ .

**Definition 3.0.2 (Coequalizer).**

A **coequalizer** is the dual notion,



**Example 3.2.**

Take  $C = \text{Sch}/S$ ,  $X/S$  a scheme, and  $X_\alpha \subset X$  an open cover. We can take two fiber products,  $X_{\alpha\beta}, X_{\beta\alpha}$ :



These are canonically isomorphic.

In  $\text{Sch}/S$ , we have


$$\coprod_{\alpha\beta} X_{\alpha\beta} \begin{matrix} \xrightarrow{f_{\alpha\beta}} \\ \xrightarrow{g_{\alpha\beta}} \end{matrix} \coprod_{\alpha} X_{\alpha} \longrightarrow X$$

where

$$\begin{aligned} f_{\alpha\beta} &: X_{\alpha\beta} \longrightarrow X_{\alpha} \\ g_{\alpha\beta} &: X_{\alpha\beta} \longrightarrow X_{\beta}; \end{aligned}$$

this is a coequalizer.

Conversely, we can glue schemes. Given  $X_\alpha \longrightarrow X_{\alpha\beta}$  (schemes over open subschemes), we need to check triple intersections:



Then  $\varphi_{\alpha\beta} : X_{\alpha\beta} \xrightarrow{\cong} X_{\beta\alpha}$  must satisfy the **cocycle condition**:

1.

$$\varphi_{\alpha\beta}^{-1}(X_{\beta\alpha} \cap X_{\beta\gamma}) = X_{\alpha\beta} \cap X_{\alpha\gamma},$$

noting that the intersection is exactly the fiber product  $X_{\beta\alpha} \times_{X_\beta} X_{\beta\gamma}$ .

2. The following diagram commutes:

$$\begin{array}{ccc} X_{\alpha\beta} \cap X_{\alpha\gamma} & \xrightarrow{\varphi_{\alpha\gamma}} & X_{\gamma\alpha} \cap X_{\gamma\beta} \\ & \searrow \varphi_{\alpha\beta} \quad \nearrow \varphi_{\beta\gamma} & \\ & X_{\beta\alpha} \cap X_{\beta\gamma} & \end{array}$$

Then there exists a scheme  $X/S$  such that  $\coprod_{\alpha\beta} X_{\alpha\beta} \rightrightarrows \coprod X_\alpha \longrightarrow X$  is a coequalizer; this is the gluing.

Subfunctors satisfy a patching property because morphisms to schemes are locally determined. Thus representable functors (e.g. functors of points) have to be (Zariski) sheaves.

**Definition 3.0.3** (Zariski Sheaf).

A functor  $F : (\text{Sch}/S)^{\text{op}} \longrightarrow \text{Set}$  is a *Zariski sheaf* iff for any scheme  $T/S$  and any open cover  $T_\alpha$ , the following is an equalizer:

$$F(T) \longrightarrow \prod F(T_\alpha) \rightrightarrows \prod_{\alpha\beta} F(T_{\alpha\beta})$$

where the maps are given by restrictions.

---

**Example 3.3.**

Any representable functor is a Zariski sheaf precisely because the gluing is a coequalizer. Thus if you take the cover

$$\coprod_{\alpha\beta} T_{\alpha\beta} \longrightarrow \coprod_{\alpha} T_{\alpha} \longrightarrow T,$$

since giving a local map to  $X$  that agrees on intersections is enough to specify a map from  $T \longrightarrow X$ .

Thus any functor represented by a scheme automatically satisfies the sheaf axioms.

**Definition 3.0.4** (Subfunctors, Open/Closed Functors).

Suppose we have a morphism  $F' \longrightarrow F$  in the category  $\text{Fun}(\text{Sch}/S, \text{Set})$ .

- This is a **subfunctor** if  $\iota(T)$  is injective for all  $T/S$ .
- $\iota$  is **open/closed/locally closed** iff for any scheme  $T/S$  and any section  $\xi \in F(T)$  over  $T$ , then there is an open/closed/locally closed set  $U \subset T$  such that for all maps of schemes  $T' \xrightarrow{f} T$ , we can take the pullback  $f^*\xi$  and  $f^*\xi \in F'(T')$  iff  $f$  factors through  $U$ .

I.e. we can test if pullbacks are contained in a subfunctors by checking factorization.

**Note** This is the same as asking if the subfunctor  $F'$ , which maps to  $F$  (noting a section is the same as a map to the functor of points), and since  $T \longrightarrow F$  and  $F' \longrightarrow F$ , we can form the fiber product  $F' \times_F T$ :

$$\begin{array}{ccc} F' & \longrightarrow & F \\ \uparrow & & \uparrow \xi \\ F' \times_F T & \xrightarrow{g} & T \end{array}$$

and  $F' \times_F T \cong U$ .

Note: this is almost tautological!

Thus  $F' \longrightarrow F$  is open/closed/locally closed iff  $F' \times_F T$  is representable and  $g$  is open/closed/locally closed.

I.e. base change is representable, and (?).

**Exercise (Tautologous)**

1. If  $F' \longrightarrow F$  is open/closed/locally closed and  $F$  is representable, then  $F'$  is representable as an open/closed/locally closed subscheme
2. If  $F$  is representable, then open/etc subschemes yield open/etc subfunctors

Mantra: Treat functors as spaces. We have a definition of open, so now we'll define coverings.

**Definition 3.0.5** (Open Covers).

A collection of open subfunctors  $F_{\alpha} \subset F$  is an **open cover** iff for any  $T/S$  and any section  $\xi \in F(T)$ , i.e.  $\xi : T \longrightarrow F$ , the  $T_{\alpha}$  in the following diagram are an open cover of  $T$ :

$$\begin{array}{ccc}
F_\alpha & \longrightarrow & F \\
\uparrow & & \uparrow \xi \\
T_\alpha & \longrightarrow & T
\end{array}$$

**Example 3.4.**

Given

$$F(s) = \{ \mathcal{O}_s^{n+1} \longrightarrow L \longrightarrow 0 \}$$

and  $F_i(s)$  given by those where  $s_i \neq 0$  everywhere, the  $F_i \longrightarrow F$  are an open cover. Because the sections generate everything, taking the  $T_i$  yields an open cover.

**Proposition 3.1.**

A Zariski sheaf  $F : (\text{Sch}/S)^{\text{op}} \longrightarrow \text{Set}$  with a representable open cover is representable.

*Proof.*

Let  $F_\alpha \subset F$  be an open cover, say each  $F_\alpha$  is representable by  $x_\alpha$ . Form the fiber product  $F_{\alpha\beta} = F_\alpha \times_F F_\beta$ . Then  $x_\beta$  yields a section (plus some openness condition?), so  $F_{\alpha\beta} = x_{\alpha\beta}$  representable. Because  $F_\alpha \subset F$ , the  $F_{\alpha\beta} \longrightarrow F_\alpha$  have the correct gluing maps.

This follows from Yoneda (schemes embed into functors), and we get maps  $x_{\alpha\beta} \longrightarrow x_\alpha$  satisfying the gluing conditions. Call the gluing scheme  $x$ ; we'll show that  $x$  represents  $F$ .

First produce a map  $x \longrightarrow F$  from the sheaf axioms. We have a map  $\xi \in \prod_{\alpha} F(x_\alpha)$ , and because we can pullback, we get a unique element  $\xi \in F(X)$  coming from the diagram

$$F(x) \longrightarrow \prod F(x_\alpha) \rightrightarrows \prod_{\alpha\beta} F(x_{\alpha\beta}).$$

■

**Lemma 3.2.**

If  $E \longrightarrow F$  is a map of functors and  $E, F$  are Zariski sheaves, where there are open covers  $E_\alpha \longrightarrow E, F_\alpha \longrightarrow F$  with commutative diagrams

$$\begin{array}{ccc}
E & \longrightarrow & F \\
\uparrow & & \uparrow \\
E_\alpha & \xrightarrow{\cong} & F_\alpha
\end{array}$$

(i.e. these are isomorphisms locally) then the map is an isomorphism.

With the following diagram, we're done by the lemma:

$$\begin{array}{ccc}
X & \longrightarrow & F \\
\uparrow & & \uparrow \\
X_\alpha & \xrightarrow{\cong} & F_\alpha
\end{array}$$

**Example 3.5.**

For  $S$  and  $E$  a locally free coherent  $\mathcal{O}_S$  module,

$$\mathbb{P}E(T) = \{f^*E \longrightarrow L \longrightarrow 0\} / \sim$$

is a generalization of projectivization, then  $S$  admits a cover  $U_i$  trivializing  $E$ .

Then the restriction  $F_i \longrightarrow \mathbb{P}E$  where  $F_i(T)$  is the above set if  $f$  factors through  $U_i$  and empty otherwise. On  $U_i$ ,  $E \cong \mathcal{O}_{U_i}^{n_i}$ , so  $F_i$  is representable by  $\mathbb{P}_{U_i}^{n_i-1}$  by the proposition. (Note that this is clearly a sheaf.)

**Example 3.6.**

For  $E$  locally free over  $S$  of rank  $n$ , take  $r < n$  and consider the functor  $\text{Gr}(k, E)(T) = \{f^*E \longrightarrow Q \longrightarrow 0\} / \sim$  (a Grassmannian) where  $Q$  is locally free of rank  $k$ .

**Exercise**

- Show that this is representable
- For the Plucker embedding

$$\text{Gr}(k, E) \longrightarrow \mathbb{P} \wedge^k E,$$

a section over  $T$  is given by  $f^*E \longrightarrow Q \longrightarrow 0$  corresponding to

$$\wedge^k f^*E \longrightarrow \wedge^k Q \longrightarrow 0,$$

noting that the left-most term is  $f^* \wedge^k E$ .

Show that this is a closed subfunctor. (That it's a functor is clear, that it's closed is not.)

Take  $S = \text{Spec } k$ , then  $E$  is a  $k$ -vector space  $V$ , then sections of the Grassmannian are quotients of  $V \otimes \mathcal{O}$  that are free of rank  $n$ .

Take the subfunctor  $G_w \subset \text{Gr}(k, V)$  where

$$G_w(T) = \{\mathcal{O}_T \otimes V \longrightarrow Q \longrightarrow 0\} \text{ with } Q \cong \mathcal{O}_T \otimes W \subset \mathcal{O}_T \otimes V.$$

If we have a splitting  $V = W \oplus U$ , then  $G_w = \mathbb{A}(\text{hom}(U, W))$ . If you show it's closed, it follows that it's proper by the exercise at the beginning.

Thursday: Define the Hilbert functor, show it's representable. The Hilbert scheme functor gives e.g. for  $\mathbb{P}^n$  of all flat families of subschemes.

---

## 4 Thursday January 16th

### 4.1 Subfunctors

A functor  $F' \subset F : (\text{Sch}/S)^{\text{op}} \rightarrow \text{Set}$  is **open** iff for all  $T \xrightarrow{\xi} F$  where  $T = h_T$  and  $\xi \in F(T)$ .

We can take fiber products:

$$\begin{array}{ccc} F' & \longrightarrow & F \\ \uparrow & & \uparrow \\ F' \times_F T & \xrightarrow{\text{Open}} & T \\ \text{Representable} & & \end{array}$$

So we can think of “inclusion in  $F$ ” as being an *open condition*: for all  $T/S$  and  $\xi \in F(T)$ , there exists an open  $U \subset T$  such that for all covers  $f : T' \rightarrow T$ , we have

$$F(f)(\xi) = f^*(\xi) \in F'(T')$$

iff  $f$  factors through  $U$ .

Suppose  $U \subset T$  in  $\text{Sch}/T$ , we then have

$$h_{U/T}(T') = \begin{cases} \emptyset & T' \rightarrow T \text{ doesn't factor} \\ \{\text{pt}\} & \text{otherwise} \end{cases}.$$

which follows because the literal statement is  $h_{U/T}(T') = \text{hom}_T(T', U)$ .

By the definition of the fiber product,

$$(F' \times_F T)(T') = \left\{ (a, b) \in F'(T) \times T(T) \mid \xi(b) = \iota(a) \text{ in } F(T) \right\},$$

where  $F' \xrightarrow{\iota} F$  and  $T \xrightarrow{\xi} F$ .

So note that the RHS diagram here is exactly given by pullbacks, since we identify sections of  $F/T'$  as sections of  $F$  over  $T/T'$  (?).

$$\begin{array}{ccc} F' & \xrightarrow{\iota} & F \\ \uparrow & & \uparrow \xi \\ F' \times_F T & \longrightarrow & T \\ & & \nwarrow f \\ & & T' \end{array}$$

$f \circ \xi$

We can thus identify

$$(F' \times_F T)(T') = h_{U/S}(T'),$$

and so for  $U \subset T$  in  $\text{Sch}/S$  we have  $h_{U/S} \subset h_{T/S}$  is the functor of maps that factor through  $U$ . We just identify  $h_{U/S}(T') = \text{hom}_S(T', U)$  and  $h_{T/S}(T') = \text{hom}_S(T', T)$ .

**Example 4.1.**

$\mathbb{G}_m, \mathbb{G}_a$ .  $\mathbb{G}_a$  represents giving a global function,  $\mathbb{G}_m$  represents giving an invertible function.

$$\begin{array}{ccc} \mathbb{G}_m & \longrightarrow & \mathbb{G}_a \\ \uparrow & \lrcorner & \uparrow \\ T' & \longrightarrow & T \end{array} \quad \begin{array}{c} \\ \\ f \in \mathcal{O}_T(T) \end{array}$$

where  $T' = \{f \neq 0\}$  and  $\mathcal{O}_T(T)$  are global functions.

## 4.2 Actual Geometry: Hilbert Schemes

The best moduli space!

Want to parameterize families of subschemes over a fixed object. Fix  $k$  a field,  $X/k$  a scheme; we'll parameterize subschemes of  $X$ .

**Definition 4.0.1** (Hilbert Functor).

The hilbert functor is given by

$$\text{Hilb}_{X/S} : (\text{Sch}/S)^{op} \longrightarrow \text{Set}$$

which sends  $T$  to closed subschemes  $Z \subset X \times_S T \longrightarrow T$  which are flat over  $T$ .

Here flatness replaces the Cartier condition.

**Definition 4.0.2** (Flatness).

For  $X \xrightarrow{f} Y$  and  $\mathbb{F}$  a coherent sheaf on  $X$ ,  $f$  is flat over  $Y$  iff for all  $x \in X$  the stalk  $F_x$  is a flat  $\mathcal{O}_{y, f(x)}$ -module.

Note that  $f$  is flat if  $\mathcal{O}_x$  is.

Flatness corresponds to varying continuously.

**Warning:** Unless otherwise stated, assume schemes are Noetherian.

Note that everything works out if we only path with finite covers.

**Remark** If  $X/k$  is projective, so  $X \subset \mathbb{P}_k^n$ , we have line bundles  $\mathcal{O}_x(1) = \mathcal{O}(1)$ . For any sheaf  $F$  over  $X$ , there is a hilbert polynomial  $P_F(n) = \chi(F(n)) \in \mathbb{Z}[n]$ . (i.e. we twist by  $\mathcal{O}(1)$   $n$  times.)



The cohomology of  $F$  isn't changed by the pushforward into  $\mathbb{P}_n$  since it's a closed embedding, i.e.

$$\chi(X, F) = \chi(\mathbb{P}^n, i_* F) = \sum (-1)^i \dim_k H^i(\mathbb{P}^n, i_* F(n)).$$

**Fact (First)** For  $n \gg 0$ ,  $\dim_k H^0 = \dim M_n$ , the  $n$ th graded piece of  $M$ , which is a graded module over the homogeneous coordinate ring whose  $i_* F = \tilde{M}$ .

In general, for  $L$  ample of  $X$  and  $F$  coherent on  $X$ , we can define a **Hilbert polynomial**,

$$P_F(n) = \chi(F \otimes L^n).$$

This is an invariant of a polarized projective variety, and in particular subschemes. Over irreducible bases, flatness corresponds to this invariant being constant.

**Proposition 4.1.**

For  $f : X \rightarrow S$  projective, i.e. there is a factorization:

$$\begin{array}{ccc} X & \xrightarrow{\quad} & \mathbb{P}^n \times S \ni \mathcal{O}(1) \\ & \searrow f & \swarrow \\ & S & \end{array}$$

If  $S$  is reduced, irreducible, locally Noetherian, then  $f$  is flat  $\iff P_{\mathcal{O}_{x_s}}$  is constant for all  $s \in S$ .

To be more precise, look the base change to  $X_1$ , and the pullback of the fiber?  $\mathcal{O} \Big|_{x_i} ?$

Note: not using the word “integral” here!  $S$  is flat  $\iff$  the hilbert polynomial over the fibers are constant.

**Example 4.2.**

The zero-dimensional subschemes  $Z \in \mathbb{P}_k^n$ , then  $P_Z$  is the length of  $Z$ , i.e.  $\dim_k(\mathcal{O}_Z)$ , and

$$P_Z(n) = \chi(\mathcal{O}_Z \otimes \mathcal{O}(n)) = \chi(\mathcal{O}_Z) = \dim_k H^0(Z; \mathcal{O}_Z) = \dim_k \mathcal{O}_Z(Z).$$

For two closed points in  $\mathbb{P}^2$ ,  $P_Z = 2$ .

Consider the affine chart  $\mathbb{A}^2 \subset \mathbb{P}^2$ , which is given by

$$\text{Spec } k[x, y]/(y, x^2) \cong k[x]/(x^2)$$

and  $P_Z = 2$ . I.e. in flat families, it has to record how the tangent directions come together.

**Example 4.3.**

Consider the flat family  $xy = 1$  (flat because it's an open embedding) over  $k[x]$ , here we have points running off to infinity.

**Proposition 4.2 (Modified Characterization of Flatness for Sheaves).**

A sheaf  $F$  is flat iff  $P_{F_S}$  is constant.

**4.2.1 Proof**

Assume  $S = \text{Spec } A$  for  $A$  a local Noetherian domain.

**Lemma 4.3.**

For  $F$  a coherent sheaf on  $X/A$  is flat, we can take the cohomology via global sections  $H^0(X; F(n))$ . This is an  $A$ -module, and is a free  $A$ -module for  $n \gg 0$ .

*Proof (of Lemma).*

Assumed  $X$  was projective, so just take  $X = \mathbb{P}_A^n$  and let  $F$  be the pushforward. There is a correspondence sending  $F$  to its ring of homogeneous sections constructed by taking the sheaf associated to the graded module  $\sum_{n \gg 0} H^0(\mathbb{P}_A^m; F(n))$ . This is equal to  $\oplus_{n \gg 0} H^0(\mathbb{P}_A^m; F(n))$  and taking the associated sheaf ( $Y \mapsto \tilde{Y}$ , as per Hartshorne's notation) which is free, and thus  $F$  is free.

See tilde construction in Hartshorne, essentially amounts to localizing free tings.

Conversely, take an affine cover  $U_i$  of  $X$ . We can compute the cohomology using Čech cohomology, i.e. taking the Čech resolution. We can also assume  $H^i(\mathbb{P}^m; F(n)) = 0$  for  $n \gg 0$ , and the Čech complex vanishes in high enough degree. But then there is an exact sequence

$$0 \longrightarrow H^0(\mathbb{P}^m; F(n)) \longrightarrow \mathcal{C}^0(\underline{U}; F(n)) \longrightarrow \cdots \longrightarrow \mathcal{C}^m(\underline{U}; F(n)) \longrightarrow 0.$$

Assuming  $F$  is flat, and using the fact that flatness is a 2 out of 3 property, the images of these maps are all flat by induction from the right.

Finally, local Noetherian + finitely generated flat implies free. ■

By the lemma, we want to show  $H^0(\mathbb{P}^m; F(n))$  is free for  $n \gg 0$  iff the hilbert polynomials on the fibers  $P_{F_S}$  are all constant.

**Claim 1 (1).**

It suffices to show that for each point  $s \in \text{Spec } A$ , we have

$$H^0(X_s; F_S(n)) = H^0(X; F(n)) \otimes k(S)$$

for  $k(S)$  the residue field, for  $n \gg 0$ .

Note that  $P_{F_s}$  measures the rank of the LHS.

$\implies$  : The dimension of RHS is constant, whereas the LHS equals  $P_{F_S}(n)$ .

$\impliedby$  : If the dimension of the RHS is constant, so the LHS is free.

For a f.g. module over a local ring, testing if localization at closed point and generic point have the same rank.

For  $M$  a finitely generated module over  $A$ , find  $0 \rightarrow A^n \rightarrow M \rightarrow Q$  is surjective after tensoring with  $\text{Frac}(A)$ , and tensoring with  $k(S)$  for a closed point, if  $\dim A^n = \dim M$  then  $Q = 0$ .

*Proof (of Claim 1).*

By localizing, we can assume  $s$  is a closed point. Since  $A$  is Noetherian, its ideal is f.g. and we have

$$A^m \rightarrow A \rightarrow k(S) \rightarrow 0.$$

We can tensor with  $F$  (viewed as restricting to fiber) to obtain

$$F(n)^m \rightarrow F(n) \rightarrow F_S(n) \rightarrow 0.$$

Because  $F$  is flat, this is still exact.

We can take  $H^*(x, \cdot)$ , and for  $n \gg 0$  only  $H^0$  survives. This is the same as tensoring with  $H^0(x, F(n))$ . ■

**Definition 4.3.1** (Hilbert Polynomial Subfunctor).

Given a polynomial  $P \in \mathbb{Z}[n]$  for  $X/S$  projective, we define a subfunctor by picking only those with Hilbert polynomial  $p$  fiberwise as  $\text{Hilb}_{X/S}^P \subset \text{Hilb}_{X/S}$ . This is given by  $Z \subset X \times_S T$  with  $P_Z = P$ .

**Theorem 4.4 (Grothendieck).**

If  $S$  is Noetherian and  $X/S$  projective, then  $\text{Hilb}_{X/S}^P$  is representable by a projective  $S$ -scheme.

See cycle spaces in analytic geometry.

## 5 Thursday January 23

Some facts about the Hilbert polynomial:

1. For a subscheme  $Z \subset \mathbb{P}_k^n$  with  $\deg P_z = \dim Z = n$ , then

$$p_z(t) = \deg z t^n / (n!) + O(t^{n-1}).$$

2. We have  $p_z(t) = \chi(\mathcal{O}_z(t))$ , consider the sequence


$$0 \rightarrow I_z(t) \rightarrow \mathcal{O}_{\mathbb{P}^n}^{(t)} \rightarrow \mathcal{O}_z^{(t)} \rightarrow 0,$$

then  $\chi(I_z(t)) = \dim H^0(\mathbb{P}^n, J_z(t))$  for  $t \gg 0$ , and  $p_z(0)$  is the Euler characteristic of  $\mathcal{O}_Z$ .

Serre vanishing, Riemann-Roch, ideal sheaf.

---

**Example 5.1** (Good to keep in mind).  
The twisted cubic:



Then

$$p_C(t) = (\deg C)t + \chi(\mathcal{O}_{\mathbb{P}^1}) = 3t + 1.$$

### 5.0.1 Hypersurfaces

Recall that length 2 subschemes of  $\mathbb{P}^1$  are the same as specifying quadratics that cut them out, each such  $Z \subset \mathbb{P}^1$  satisfies  $Z = V(f)$  where  $\deg f = d$  and  $f$  is homogeneous. So we'll be looking at  $\mathbb{P}H^0(\mathbb{P}_k^n, \mathcal{O}(d))^\vee$ , and the guess would be that this is  $\text{Hilb}_{\mathbb{P}_k^n}$

Resolve the structure sheaf

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^n}(-d) \longrightarrow \mathcal{O}_{\mathbb{P}^n}(t) \longrightarrow \mathcal{O}_D(t) \longrightarrow 0.$$

so we can twist to obtain

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^n}(t-d) \longrightarrow \mathcal{O}_{\mathbb{P}^n}(t) \longrightarrow \mathcal{O}_D(t) \longrightarrow 0.$$

Then

$$\chi(\mathcal{O}_D(t)) = \chi(\mathcal{O}_{\mathbb{P}^n}(t)) - \chi(\mathcal{O}_{\mathbb{P}^n}(t-d)),$$

which is

$$\binom{n+t}{n} - \binom{n+t-d}{n} = \frac{dt^{n-1}}{(n-1)!} + O(t^{n-2}).$$

---

**Lemma 5.1.**

Anything with the Hilbert polynomial of a degree  $d$  hypersurface is in fact a degree  $d$  hypersurface.

We want to write a morphism of functors

$$\mathrm{Hilb}_{\mathbb{P}_k^n}^{P_{n,d}} \longrightarrow \mathbb{P}H^0(\mathbb{P}^n, \mathcal{O}(d))^\vee.$$

which sends flat families to families of equations cutting them out.

Want

$$Z \subset \mathbb{P}^n \times S \longrightarrow \mathcal{O}_S \otimes H^0(\mathbb{P}^n, \mathcal{O}(d))^\vee \longrightarrow L \longrightarrow 0.$$

This happens iff

$$0 \longrightarrow L^\vee \longrightarrow \mathcal{O}_S \otimes H^0(\mathbb{P}^n, \mathcal{O}(d))$$

with torsion-free quotient.

Note that we use  $L^\vee$  instead of  $\mathcal{O}_S$  because of scaling.

We have

$$\begin{aligned} 0 &\longrightarrow I_Z \longrightarrow \mathcal{O}_{\mathbb{P}^n \times S} \longrightarrow \mathcal{O}_Z \longrightarrow 0 \\ 0 &\longrightarrow I_Z(d) \longrightarrow \mathcal{O}_{\mathbb{P}^n \times S}(d) \longrightarrow \mathcal{O}_Z(d) \longrightarrow 0 \quad \text{by twisting.} \end{aligned}$$

We then consider  $\pi_S : \mathbb{P}^n \times S \longrightarrow S$ , and apply the pushforward to the above sequence noting that it is not right-exact.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_{S*} I_Z(d) & \longrightarrow & \pi_{S*} \mathcal{O}_{\mathbb{P}^n \times S}(d) & \longrightarrow & \pi_{S*} \mathcal{O}_Z(d) \longrightarrow 0 \\ \parallel & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & L^\vee = \mathcal{O}_S \otimes H^0(\mathbb{P}^n, \mathcal{O}(d)) & \longrightarrow & \text{locally free} & \longrightarrow & 0 \end{array}$$

Note: above diagram may be off horizontally? Todo: check.

This equality follows from flatness, cohomology, and base change. In particular, we need the following facts.

The scheme-theoretic fibers, given by  $H^0(\mathbb{P}^n, I_Z(d))$  and  $H^0(\mathbb{P}^n, \mathcal{O}_Z(d))$ , are all the same dimension.

Using

1. Cohomology and base change, i.e. for  $X \xrightarrow{f} Y$  a map of Noetherian schemes (or just finite-type) and  $F$  a sheaf on  $X$  which is flat over  $Y$ , there is a natural map (not usually an isomorphism)

$$R^i f_* f \otimes k(y) \longrightarrow H^i(x_y, F|_{x_y}),$$

but is an isomorphism if  $\dim H^i(x_y, F|_{x_y})$  is constant, in which case  $R^i f_* f$  is locally free.

---

2. If  $Z \subset \mathbb{P}_k^n$  is a degree  $d$  hypersurface, then independently we know

$$\dim H^0(\mathbb{P}^n, I_Z(d)) = 1 \text{ and } \dim H^0(\mathbb{P}^n, \mathcal{O}_Z(d)) = \binom{d+n}{n} - 1.$$

To get a map going backwards, we take the universal degree 2 polynomial and form

$$V(a_{00}x_0^2 + a_{11}x_1^2 + a_{12}x_2^2 + a_{01}x_0x_1 + a_{02}x_0x_2 + a_{12}x_1x_2) \subset \mathbb{P}^2 \times \mathbb{P}^5.$$

### 5.0.2 Example: Twisted Cubics

Consider a map  $\mathbb{P}^1 \rightarrow \mathbb{P}^3$  obtained by taking a basis of a homogeneous cubic polynomial. The canonical example is

$$(x, y) \rightarrow (x^3, x^2y, xy^2, y^3).$$

Then  $P_C(t) = 3t + 1$ , and  $\text{Hilb}_{\mathbb{P}_k^3}^{3t+1}$  has a component with generic point a twisted cubic, and another component with points a curve disjoint union a point, and the overlap are nodal curves with a “fat” 3-dimensional point:



Then  $P_{C'} = 1 + \tilde{P}$ , the hilbert polynomial of just the base without the disjoint point, so this equals  $1 + P_{2,3} = 1 + (3t + 0) = 3t + 1$ . For  $P_{C''}$ , we take the sequence

$$0 \longrightarrow k \longrightarrow \mathcal{O}_{C''} \longrightarrow \mathcal{O}_{C''\text{reduced}} \longrightarrow 0,$$

so

$$P_{C''} = 1 + P_{C''\text{red}} = 3t + 1.$$

Note: flat families have to have the same constant Hilbert polynomial.

Note that we can get paths in this space from  $C \longrightarrow C''$  and  $C' \longrightarrow C''$  by collapsing a twisted cubic onto a plane, and sending a disjoint point crashing into the node on a nodal cubic.

We're mapping  $\mathbb{P}^1 \longrightarrow \mathbb{P}^3$ , and there is a natural action of  $\mathbb{P}\text{GL}(4) \curvearrowright \mathbb{P}^3$ , so we get a map

---


$$\mathbb{PGL}(4) \times \mathbb{P}^3 \longrightarrow \mathbb{P}^3.$$

Let  $c \in \mathbb{P}^3$  and let  $\mathcal{C}$  be the preimage. This induces (?) a map

$$\mathbb{PGL}(4) \longrightarrow \mathrm{Hilb}_{\mathbb{P}^3}^{3t+1}$$

where the fiber over  $[C]$  in the latter is  $\mathbb{PGL}(2) = \mathrm{Aut}(\mathbb{P}^1)$ . By dimension counting, we find that the dimension of the twisted cubic component is  $15 - 3 = 12$ .

The 15 in the other component comes from 3-dim choices of plane, 3-dim choices of a disjoint point, and

$$\mathbb{P}H^0(\mathbb{P}^2, \mathcal{O}(3))^\vee \cong \mathbb{P}^9,$$

yielding 15 dimensions.

To show that these are actually different components, we use Zariski tangent spaces. Let  $T_1$  be the tangent space of the twisted cubic component, then

$$\dim T_1 \mathrm{Hilb}_{\mathbb{P}_k^3}^{3t+1} = 12,$$

and similarly the dimension of the tangent space over the  $C'$  component is 15.

**Fact (from Algebra)** Let  $A$  be Noetherian and local, then the dimension of the Zariski tangent space,  $\dim \mathfrak{m}/\mathfrak{m}^2 \geq \dim A$ , the Krull dimension. If this is an equality, then  $A$  is regular.

Thus dimensions of the tangent spaces give an upper bound.

Proposition: If  $X/k$  is projective and  $P$  is a Hilbert polynomial, then  $[Z] \in \mathrm{Hilb}_{X/k}^P$ , i.e. a closed subscheme of  $X$  with hilbert polynomial  $p$  (note there's an ample bundle floating around) then the tangent space is  $\mathrm{hom}_{\mathcal{O}_x}(I_z, \mathcal{O}_z)$ .

## 6 Tuesday January 28th

Last time: Twisted cubics, given by  $\mathrm{Hilb}_{\mathbb{P}_k^3}^{3t+1}$ .





We got lower (?) bounds on the dimension by constructing families, but want an exact dimension.

Key: Let  $Z \subset X$  be a closed  $k$ -dimensional subspace.

Proposition: For  $[z] \in \text{Hilb}_{X/k}^P(k)$ , we have an identification of the Zariski tangent space  $T_{[z]}\text{Hilb}_{X/k}^P = \text{hom}_{\mathcal{O}_x}(I_z, \mathcal{O}_z)$ .

Say  $F : (\text{Sch}/K)^{\text{op}} \rightarrow \text{Set}$  is a function and let  $x \in F(k)$ . There is an inclusion  $i : \text{Spec } k \hookrightarrow \text{Spec } k[\varepsilon]$ . Then there is an induced map  $F(\text{Spec } k[\varepsilon]) \xrightarrow{i^*} F(\text{Spec } k)$  where  $T_x F := (i^*)^{-1}(x) \mapsto x$ . So if  $F$  is represented by a scheme  $H/k$ , then  $T_x h_J = T_x H = (\mathfrak{m}_x/\mathfrak{m}_x^2)^\vee$  over  $k$ .

Will need a criterion for flatness later, esp. for Artinian thickenings.

Lemma: Assume  $A'$  is a Noetherian ring and  $0 \rightarrow J \rightarrow A' \rightarrow A \rightarrow 0$  with  $J^2 = 0$ . Assume we have  $X'/\text{Spec } A'$ , and a coherent sheaf  $F'$  on  $X'$ , where  $X'$  is Noetherian.

Then  $F'$  is flat over  $A'$  iff



1.  $F$  is flat
2.  $0 \rightarrow F \otimes_A J \rightarrow F'$  is exact.

Take the first exact sequence and tensor with  $F'$  (which is right-exact), then  $J \otimes_{A'} F' = J \otimes_A$  canonically. This follows because  $J = J \otimes_{A'} A$ , and there is an isomorphism  $J \otimes_{A'} A' \rightarrow J \otimes_{A'} A$ . And  $F = F' \otimes_{A'} A$  is a pullback of  $F'$ . If flat, then tensoring is exact.

Proof: Both conditions are necessary since pullbacks of flats are flat (1), and (2) gives the flatness condition.)

Recall that for a module over a Noetherian ring,  $M/A$ ,  $M$  is flat over  $A$  iff  $\text{Tor}_1^A(M, A/p) = 0$  for all prime  $p$ . Reason: Tor commutes with direct limits, so  $M$  is flat iff  $\text{Tor}_1^A(M, N) = 0$  for all finitely generated  $N$ . Since  $A$  is Noetherian,  $N$  has a finite filtration  $N$  where  $N_i/N_{i+1} \cong A/p_i$ . Use the fact that every ideal is contained in a prime ideal.

Take  $x \in N$ , this yields a map  $A \rightarrow N$  which factors through  $A/I$ . If we make such a filtration on  $A/I$ , then we can quotient  $N$  by  $\text{im } f$  where  $f: A/I \rightarrow N$ . Continuing inductively, the resulting filtration must stabilize. So we can assume  $N = A/I$ .

Then  $I$  is contained in a maximal.

Exercise: Finish proof. See Aatiyah Macdonald.

So it's enough to show that  $\text{Tor}_1^{A'}(F', A'/p') = 0$  for all primes  $p' \subset A'$ .

Observation: Since  $J$  is nilpotent,  $J \subset p'$ . Let  $p = p'/J$ , this is a prime ideal.

We have an exact diagram by taking quotients:

---


$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & J & \longrightarrow & p' & \longrightarrow & p \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & J & \longrightarrow & A' & \longrightarrow & A \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \\
& & & & A'/p' & & A/p \\
& & & & \downarrow & & \downarrow \\
& & & & 0 & & 0
\end{array}$$

So we can tensor with  $F'$  everywhere, and get a map from kernels to cokernels using the snake lemma:

$$\begin{array}{ccccccc}
& & 0 & & \text{Tor}(A, F) = 0 & & \\
& & \downarrow & & \downarrow & & \\
& & \text{Tor}_1^{A_1}(A'/p', F') & & \text{Tor}_1^{A_1}(A/p, F') & & \\
& \xrightarrow{\text{snake}} & & & & & \\
0 & \longrightarrow & J \otimes_{A'} F' & \xrightarrow{\text{by commuting square}} & p' \otimes_{A'} F' & \longrightarrow & p \otimes_{A'} F' \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & J \otimes_{A'} F' & \xleftarrow{\text{by (2)}} & A' \otimes_{A'} F' & \longrightarrow & A \otimes_{A'} F' \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& \xrightarrow{\text{snake}} & & & A'/p' \otimes_{A'} F' & \xrightarrow{=} & A/p \otimes_{A'} F' \\
& & & & \downarrow & & \downarrow \\
& & & & 0 & & 0
\end{array}$$

Then by (1), we have  $\text{Tor}_1^{A'}(A'/p', F') = \text{Tor}_1^{A'}(A/p, F') = 0$ .

■

We will just need this for  $A' = k[\varepsilon]$  and  $A = k$ .

Proposition:  $T_z \text{Hilb}_{X/k} = \text{hom}_{\mathcal{O}_x}(I_z, \mathcal{O}_z)$ .

Proof: Again we have  $T_z \text{Hilb}_{X/k} \subset \text{Hilb}_{X/k}(k[\varepsilon])$ , and is given by  $\left\{ Z' \subset X \times_{\text{Spec } k} \text{Spec } k[\varepsilon]_{\text{flat}/k[\varepsilon]} \mid Z' \times_{\text{Spec } k} k \right\}$ .

We have an exact diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & I_{Z'} & \longrightarrow & \mathcal{O}_{X[\varepsilon]} & \longrightarrow & \mathcal{O}_{Z'} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & & \downarrow & & \downarrow & & \downarrow \\
 & & k & & I_Z & \longrightarrow & \mathcal{O}_x & \longrightarrow & \mathcal{O}_z & \longrightarrow & \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 k[\varepsilon] & & \longrightarrow & I_{Z'} & \longrightarrow & \mathcal{O}_{x[\varepsilon]} & \longrightarrow & \mathcal{O}_{Z'} & \longrightarrow & \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 & & k & & I_Z & \longrightarrow & \mathcal{O}_x & \longrightarrow & \mathcal{O}_Z & \longrightarrow & \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 0 & & & & & & & & & & 
 \end{array}$$

Note the existence of a splitting above.

Given  $\phi \in \text{hom}_{\mathcal{O}_x}(I_Z, \mathcal{O}_Z)$ . We have

$$I_{Z'} = \left\{ f + \varepsilon g \mid f \in I_Z, \phi(f) = g \pmod{I_Z}, \phi(f) \in \mathcal{O}_Z, g \pmod{I_Z} \in \mathcal{O}_x/I_Z = \mathcal{O}_Z \right\}.$$

It's easy to see that  $Z' \subset x'$ , and

1.  $Z' \times k = Z$
2. It's flat over  $k[\varepsilon]$ , looking at  $0 \longrightarrow k \otimes I_{Z'} \longrightarrow I_{Z'}$ .

For the converse, take  $f \in I_Z$  and lift to  $f' = f + \varepsilon g \in I_{Z'}$ , then  $g \in \mathcal{O}_x$  is well-defined wrt  $I_Z$ . Then  $g \in \text{hom}_{\mathcal{O}_x}(I_Z, \mathcal{O}_Z)$ .

The main point: these hom sets are extremely computable.

Example: Let  $Z$  be a twisted cubic in  $\text{Hilb}_{\mathbb{P}^3/k}^{3t+1}(k)$ .

Observation:  $\text{hom}_{\mathcal{O}_x}(I_Z, \mathcal{O}_Z) = \text{hom}_{\mathcal{O}_X}(I_Z/I_Z^2, \mathcal{O}_Z) = \text{hom}_{\mathcal{O}_Z}(I_Z/I_Z^2, \mathcal{O}_Z)$ . If  $I_Z/I_Z^2$  is locally free, these are global sections of the dual, i.e.  $H^0((I_Z/I_Z^2)^\vee)$ .

In this case,  $Z \hookrightarrow X$  is regularly embedded, and thus  $(I_Z/I_Z^2)^\vee$  should be regarded as the normal bundle. Sections of the normal bundle match up with directions to take first-order deformations:

For  $i : C \hookrightarrow \mathbb{P}^3$ , there is an exact sequence  $0 \longrightarrow I/I^2 \longrightarrow i^* \Omega_{\mathbb{P}^3} \longrightarrow \Omega_\varepsilon \longrightarrow 0$ , taking duals, which induces  $0 \longrightarrow T_C \longrightarrow i^* T_{\mathbb{P}^3} \longrightarrow N_{C/\mathbb{P}^3} \longrightarrow 0$ .

How do we compute  $T_{\mathbb{P}^3}$ ? Fit into the exact sequence  $0 \longrightarrow \mathcal{O} \longrightarrow i^* \mathcal{O}(1)^4 \longrightarrow i^* T_{\mathbb{P}^3} \longrightarrow 0$ , which we can restrict to  $C$ . We have  $i^* \mathcal{O}(1) \cong \mathcal{O}_{\mathbb{P}^1}(3)$ , so  $0 \longrightarrow H^0 \mathcal{O}_C \longrightarrow H^*(\mathcal{O}(3)^4) \longrightarrow H^0(i^* T_{\mathbb{P}^3}) \longrightarrow 0$ ,



Figure 1: Image



Figure 2: Image

which looks like  $k \rightarrow k^{16} \rightarrow k^{15}$ . This yields  $0 \rightarrow H^0(T_c) \rightarrow H^0(i^*T_{\mathbb{P}^3}) \rightarrow H^0(N_{C/\mathbb{P}^3}) \rightarrow H^1T_c$ , which reduces to  $0 \rightarrow k^3 \rightarrow k^{15} \rightarrow k^{12} \rightarrow 0$ .

Example:  $\text{Hilb}_{\mathbb{P}^n}^{P_?} \cong \mathbb{P}H^0(\mathbb{P}^n, \mathcal{O}(d))^\vee$  which has dimension  $\binom{n+1}{n} - 1$ .

Pick  $Z$  a  $k$  point in this Hilbert scheme, then  $T_Z H = \text{hom}(I_Z, \mathcal{O}_Z)$ . Since  $I_Z \cong \mathcal{O}_{\mathbb{P}^n}(-d)$  which fits into  $0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-d) \rightarrow \mathcal{O}_{\mathbb{P}^n} \rightarrow \mathcal{O}_Z \rightarrow 0$ .

We can identify  $\text{hom}(I_Z, \mathcal{O}_Z) = H^0((I_Z/I_Z^2)^\vee) = H^0(\mathcal{O}_Z(d))$ .

$$0 \quad \mathcal{O}_{\mathbb{P}^n} \quad \mathcal{O}_{\mathbb{P}^n}(d) \quad \mathcal{O}_Z(d) \quad 0$$

$$0 \quad H^0(\cdot) \quad H^0(\cdot) \quad H^0(\cdot) \quad 0$$

$$k \quad k^{\binom{n+d}{n}} \quad k^{\binom{n+d}{n}-1}$$

Example: The tangent space of the following cubic:

---

We can identify  $\text{hom}_{\mathcal{O}_k}(I_Z, \mathcal{O}_Z) = H^0((I_Z/I_Z^2)^\vee) = 3 + H^0((I_{Z_0}/I_{Z_0}^2)^\vee)$ , where the latter equals  $H^0(\mathcal{O}_1 \big|_{z_0} \oplus \mathcal{O}(\zeta) \big|_{z_0})$  yielding  $3 + 9 = 12$ .

## 7 Thursday January 30th

Recall how we constructed the hilbert scheme of hypersurfaces

$$\text{Hilb}_{\mathbb{P}_k^n}^{P_{m,d}} = \mathbb{P}H^0(\mathbb{P}^n; \mathcal{O}(d))^\vee$$

A section  $\text{Hilb}_{\mathbb{P}_k^n}^P(s)$  corresponds to  $z \in \mathbb{P}_s^n$ . We can look at the exact sequence

$$0 \longrightarrow I_Z(m) \longrightarrow \mathcal{O}_{\mathbb{P}_S^n} \xrightarrow{\text{restrict}} \mathcal{O}_z(m) \longrightarrow 0.$$

as  $\mathbb{P}_s^n \xrightarrow{\pi_s} S$ , so we can pushforward along  $\pi$ , which is left-exact, so

$$0 \longrightarrow \pi_{s*} I_Z(m) \longrightarrow \pi_{s*} \mathcal{O}_{\mathbb{P}_S^n} = \mathcal{O}_S \otimes H^0(\mathbb{P}^n; \mathcal{O}(m)) \longrightarrow \mathcal{O}_z(m) \longrightarrow R^1 \pi_{s*} I_Z(m) \longrightarrow \cdots$$

*Idea:*  $Z \subset \mathbb{P}_k^n$  will be determined (in families!) by the space of degree  $d$  polynomials vanishing on  $Z$  (?), i.e.  $H^0(\mathbb{P}^n, I_Z(m)) \subset H^0(\mathbb{P}^n, \mathcal{O}(m))$  for  $m$  very large. This would give a map of functors  $\text{Hilb}_{\mathbb{P}_k^n}^P \longrightarrow \text{Gr}(N, H^0(\mathbb{P}^n, \mathcal{O}(m)))$ . If this is a closed subfunctor, a closed subfunctor of a representable functor is representable and we're done.

Note: We need to get an  $m$  uniform in  $Z$ , and more concretely:

1. First need to make sense of what it means for  $Z$  to be determined by  $H^0(\mathbb{P}^n, I_Z(m))$  for  $m$  only depending on  $P$ .
2. This works point by point, but we need to do this in families. I.e. we'll use the previous exact sequence, and want the  $R^1$  to vanish.

Slogan: We need uniform vanishing statements. There is a convenient way to package the vanishing requirements needed here. From now on, take  $k = \bar{k}$  and  $\mathbb{P}^n = \mathbb{P}_k^n$ .

### Definition 7.0.1.

A coherent sheaf  $F$  on  $\mathbb{P}^n$  is  $m$ -regular if  $H^i(\mathbb{P}^n; F(m-i)) = 0$  for all  $i > 0$ .

### Example 7.1.

Consider  $\mathcal{O}_{\mathbb{P}^n}$ , this is 0-regular. Line bundles on  $\mathbb{P}^n$  only have 0 and top cohomology. Just need to check that  $H^n(\mathbb{P}^n; \mathcal{O}(-n)) = 0$ , but by Serre duality this is  $H^0(\mathbb{P}^n; \mathcal{O}(n) \otimes \omega_{\mathbb{P}^n})^\vee$  which is  $H^0(\mathbb{P}^n; \mathcal{O}(-1))^\vee = 0$ .

**Proposition:** Assume  $F$  is  $m$ -regular. Then

1. There is a natural multiplication map from linear forms on  $\mathbb{P}^n$ ,  $H^0(\mathbb{P}^n; \mathcal{O}(1)) \otimes H^0(\mathbb{P}^n; F(k)) \rightarrow H^0(\mathbb{P}^n; F(k+1))$  which is surjective for  $k \geq n$ .

Think of this as a graded module, this tells you the lowest number of small grade pieces needed to determine the entire thing.

2.  $F$  is  $m'$ -regular for  $m' \geq m$ .
3.  $F(k)$  is globally generated for  $k \geq m$ , i.e. the restriction  $H^0(\mathbb{P}^n; F(k)) \otimes \mathcal{O}_{\mathbb{P}^n} \rightarrow F(k) \rightarrow 0$  is exact (i.e. surjective).

Example:  $\mathcal{O}$  is  $m$ -regular for  $m \geq 0$  implies  $\mathcal{O}(k)$  is  $-k$ -regular and is also  $m$ -regular for  $m \geq -k$ .

*Proof of (1) and (2):* Induction on dimension of  $n$  in  $\mathbb{P}^n$ . Coherent sheaves on  $\mathbb{P}^0$  are vector spaces, so no higher cohomology.

Step 1:

Take a generic hyperplane  $H \subset \mathbb{P}^n$ , there is an exact sequence  $0 \rightarrow \mathcal{O}(-1) \rightarrow \mathcal{O} \rightarrow \mathcal{O}_H \rightarrow 0$  where  $\mathcal{O}_H$  is the structure sheaf. Tensoring with  $F$  remains exact, so we get  $0 \rightarrow F(-1) \rightarrow F \rightarrow F_H \rightarrow 0$ . Why?  $\mathbb{A}^n \subset \mathbb{P}^n$ , let  $A = \mathcal{O}_{\mathbb{P}^n}(\mathbb{A}^n)$  be the polynomial ring over  $\mathbb{A}^n$ . Then the restriction of the first sequence to  $\mathbb{A}^n$  yields  $0 \rightarrow A \xrightarrow{f} A \rightarrow A/f \rightarrow 0$ , and thus we want  $F \xrightarrow{f} F \rightarrow F/fF \rightarrow 0$  which results after restricting the second sequence to  $\mathbb{A}^n$ .

Thus we just want  $f$  to not be a zero divisor. If we take  $f$  not vanishing on any associated point of  $F$ , then this will be exact. Associated points: generic points arising by supports of sections of  $F$ .  $F$  is coherent, so it has finitely many associated points.

If  $H$  does not contain any of the associated points of  $F$ , then the second sequence is indeed exact.

Step 2:

Twist up by  $k$  to obtain  $0 \rightarrow F(k-1) \rightarrow F(k) \rightarrow F_H(k) \rightarrow 0$ . Look at the LES in cohomology to get  $H^i(F(m-i)) \rightarrow H^i(F_H(m-i)) \rightarrow H^{i+1}(F(m-(i+1)))$ . So  $F_H$  is  $m$ -regular. By induction, this proves statements 1 and 2 for all  $F_H$ .

So take  $k = m + 1 - i$  and consider  $H^i(F(m-i)) \rightarrow H^i(F(m+1-i)) \rightarrow H^i(F_H(m+1-i))$ . We know 2 is satisfied, so the RHS is zero, and we know the LHS is zero, so the middle term is zero. Thus  $F$  itself is  $m+1$  regular, and by inducting on  $m$  we get statement 2.

By multiplication maps, we get a commutative diagram:

$$\begin{array}{ccccc}
 & & H^0(\mathcal{O}(1)) \otimes H^0(F(k)) & \xrightarrow{\quad} & H^0(\mathcal{O}(1)) \otimes H^0(F_H(k)) \\
 & \nearrow H \otimes \text{id} & \downarrow \beta & \searrow & \downarrow \\
 H^0(F(k)) & \xrightarrow{H} & H^0(F(k+1)) & \xrightarrow{\alpha} & H^0(F_H(k+1))
 \end{array}$$

We'd like to show the diagonal map is surjective.

Observation 1: The top map is a surjection, since  $H^0(F(k)) \rightarrow H^0(F_H(k)) \rightarrow H^1(F(k-1)) = 0$  for  $k \geq m$  by (2).

Observation 2: The right-hand map is surjective for  $k \geq m$ .



---

Observation 3:  $\ker(\alpha) \subset \text{im}(\beta)$  by a small diagram chase, so  $\beta$  is surjective.

This shows (1) and (2) completely.

*Proof of (3):* We know  $F(k)$  is globally generated for  $k \gg 0$ . Thus for all  $k \geq m$ ,  $F(k)$  is globally generated by (1). ■

**Theorem:** Let  $P \in \mathbb{Q}[t]$  be a Hilbert polynomial. There exists an  $m_0$  only depending on  $P$  such that for all subschemes  $Z \subset \mathbb{P}_k^n$  with hilbert polynomial  $P_Z = P$ , the ideal sheaf  $I_Z$  is  $m_0$ -regular.

*Proof:* Induct on  $n$ . For  $n = 0$ , again clear because higher cohomology vanishes and there are no nontrivial subschemes.

For a fixed  $Z$ , pick  $H$  in  $\mathbb{P}^n$  (and setting  $I := I_Z$  for notation) such that  $0 \rightarrow I(-1) \rightarrow I \rightarrow I_H \rightarrow 0$  is exact. Note that the hilbert polynomial  $P_{I_H}(t) = P_I(t) - P_I(t-1)$  and  $P_I = P_{\mathcal{O}_{\mathbb{P}^n}} - P_Z$ . By induction, there exists some  $m_1$  depending only on  $P$  such that  $I_H$  is  $m_1$ -regular. We get  $H^{i-1}(I_H(k)) \rightarrow H^i(I(k-1)) \rightarrow H^i(I(k)) \rightarrow H^i(I_H(k))$ , and for  $k \geq m_1 - i$  the LHS and RHS vanish so we get an isomorphism in the middle. By Serre vanishing, for  $k \gg 0$  we have  $H^i(I(k)) = 0$  and thus  $H^i(I(k)) = 0$  for  $k \geq m_i - i$ . This works for all  $i > 1$ , we have  $H^i(I(m_i - i)) = 0$ .

We now need to find  $m_0 \geq m_1$  such that  $H^1(I(m_0 - 1)) = 0$  (trickiest part of the proof).

**Lemma:** The sequence  $\left( \dim H^1(I(k)) \right)_{k \geq m_i - 1}$  is strictly decreasing.

Note:  $h^1 = \dim H^1$ .

Given the lemma, it's enough to take  $m_0 \geq m_1 + h^1(I(m_1 - 1))$ . Consider the LES we have a surjection  $H^0(\mathcal{O}_Z(m_1 - 1)) \rightarrow H^1(I(m_1 - 1)) \rightarrow 0$ . So the dimension of the LHS is equal to  $P_Z(m_1 - 1)$ , using the fact that terms vanish and make the Euler characteristic equal to  $P_Z$ . Thus we can take  $m_0 = m_1 + P(m_1 - 1)$ .

*Proof of Lemma:* Considering the LES  $H^0(I(k+1)) \xrightarrow{\alpha_{k+1}} H^0(I_H(k+1)) \rightarrow H^1(I(k)) \rightarrow H^1(I(k+1)) \rightarrow 0$ , where the last term is zero because  $I_H$  is  $m_1$ -regular. So the sequence  $h^1(I(k))$  is non-increasing.

Observation: If it does *not* strictly decrease for some  $k$ , then there is an equality on the RHS, which makes  $\alpha_{k+1}$  surjective. This means that  $\alpha_{k+2}$  is surjective, since  $H^0(\mathcal{O}(1)) \otimes H^0(I_H(k+1)) \rightarrow H^0(I_H(k+2))$ . So if one is surjective, everything above it is surjective, but by Serre vanishing we eventually get zeros. So  $\alpha_{k+i}$  is surjective for all  $i \geq 1$ , contradicting Serre vanishing, since the RHS are isomorphisms for all  $k$ . ■

Thus for any  $Z \subset \mathbb{P}_k^n$  with  $P_Z = P$ , we uniformly know that  $I_Z$  is  $m_0$ -regular for some  $m_0$  depending only on  $P$ .

**Claim 2.**

$Z$  is determined by the degree  $m_0$  polynomials vanishing on  $Z$ , i.e.  $H^0(I_Z(m_0))$  as a subspace of all degree  $m_0$  polynomials  $H^0(\mathcal{O}(m_0))$  and has fixed dimension. We have  $H^i(I_Z(m_0)) = 0$  for all  $i > 0$ , and in particular  $h^0(I_Z(m_0)) = P(m_0)$  is constant.

---

It is determined by these polynomials because we have a sequence

$$0 \longrightarrow I_Z(m_0) \longrightarrow \mathcal{O}(m_0) \longrightarrow \mathcal{O}_Z(m_0) \longrightarrow 0.$$

We can get a commuting diagram over it  $0 \longrightarrow H^0(I_Z(m_0)) \otimes \mathcal{O}_{\mathbb{P}^n} \longrightarrow H^0(\mathcal{O}(m_0)) \otimes \mathcal{O}_{\mathbb{P}^n} \longrightarrow \cdots$  where the middle map down is just evaluation and the first map down is a surjection. Hence  $I_Z(m_0)$ , hence  $\mathcal{O}_Z$ , hence  $Z$  is determined by  $H^0(I_Z(m_0))$ .

Next time: we'll show that this is a subfunctor that is locally closed.