

Linearization and Transversality

Sections 8.3 and 8.4

D. Zack Garza

April 2020

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
 H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Review 8.2

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
 H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Section 8.3: The Space of Perturbations of H

Goal

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Goal: Given a fixed Hamiltonian $H \in C^\infty(W \times S^1; \mathbb{R})$, perturb it (without modifying the periodic orbits) so that $\mathcal{M}(x, y)$ are manifolds of the expected dimension.

Goal

Linearization and
Transversality


D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
 H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Start by trying to construct a subspace $\mathcal{C}_\varepsilon^\infty(H) \subset \mathcal{C}^\infty(W \times S^1; \mathbb{R})$, the space of perturbations of H depending on a certain sequence $\varepsilon = \{\varepsilon_k\}$, and show it is a dense subspace.



Idea

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Idea: similar to how you build $L^2(\mathbb{R})$, define a norm $\|\cdot\|_\varepsilon$ on $C_\varepsilon^\infty(H)$ and take the subspace of finite-norm elements.

- Let $h(\mathbf{x}, t) \in C_\varepsilon^\infty(H)$ denote a perturbation of H .
- Fix $\varepsilon = \{\varepsilon_k \mid k \in \mathbb{Z}^{\geq 0}\} \subset \mathbb{R}^{>0}$ a sequence of real numbers, which we will choose carefully later.
- For a fixed $\mathbf{x} \in W$, $t \in \mathbb{R}$ and $k \in \mathbb{Z}^{\geq 0}$, define

$$|d^k h(\mathbf{x}, t)| = \max \left\{ d^\alpha h(\mathbf{x}, t) \mid |\alpha| = k \right\},$$

the maximum over all sets of multi-indices α of length k .

Note: I interpret this as

$$d^{\alpha_1, \alpha_2, \dots, \alpha_k} h = \frac{\partial^k h}{\partial x_{\alpha_1} \partial x_{\alpha_2} \cdots \partial x_{\alpha_k}},$$

the partial derivatives wrt the corresponding variables.

Linearization and
Transversality

D. Zack Garza

Review 8.2

Section 8.3: The
Space of
Perturbations of
 H

Section 8.4:
Linearizing the
Floer Equation:
The Differential
of F

Section 8.4: Linearizing the Floer Equation: The Differential of F