# **Algebra**

## D. Zack Garza

## August 17, 2019

#### **Contents**

1	Maj	or Theorems	1	
2	Lecture 1 (Thu 15 Aug 2019)			
	2.1	Definitions	1	
	2.2	Preliminaries	3	
	2.3	Cyclic Groups	3	
	2.4	Homomorphisms	4	
	2.5	Direct Products	4	
	2.6	Finitely Generated Abelian Groups	4	
	2.7	Fundamental Homomorphism Theorem	5	
		2.7.1 The First Homomorphism Theorem		
		2.7.2 The Second Theorem		
3	Lect	ture 2	5	

### 1 Major Theorems

**Theorem 1** (Cauchy). For any prime p dividing the order of G, there is an element x of order p (and thus a subgroup  $H = \langle x \rangle$ ).

**Theorem 2** (Lagrange). If  $H \leq G$  is a subgroup, then  $|H| \mid |G|$ .

**Theorem 3** (Sylow 1). If |G| = n has prime factorization  $n = \prod p_i^{a_i}$ , then G has subgroups of order  $p_i^{a_i}$  for every i.

## 2 Lecture 1 (Thu 15 Aug 2019)

We'll be using Hungerford's Algebra text.

#### 2.1 Definitions

The following definitions will be useful to know by heart:

- The order of a group
- Cartesian product
- Relations
- Equivalence relation
- Partition
- Binary operation
- Group
- Isomorphism
- Abelian group
- Cyclic group
- Subgroup
- Greatest common divisor
- Least common multiple
- Permutation
- Transposition
- Orbit
- Cycle
- The symmetric group  $S^n$
- The alternating group  $A_n$
- Even and odd permutations
- Cosets
- Index
- The direct product of groups
- Homomorphism
- Image of a function
- Inverse image of a function
- Kernel
- Normal subgroup
- Factor group
- Simple group

Here is a rough outline of the course:

- Group Theory
  - Groups acting on sets
  - Sylow theorems and applications
  - Classification
  - Free and free abelian groups
  - Solvable and simple groups
  - Normal series
- Galois Theory
  - Field extensions
  - Splitting fields
  - Separability
  - Finite fields
  - Cyclotomic extensions
  - Galois groups
  - Solvability by radicals
- Module theory

- Free modules
- Homomorphisms
- Projective and injective modules
- Finitely generated modules over a PID
- Linear Algebra
  - Matrices and linear transformations
  - Rank and determinants
  - Canonical forms
  - Characteristic polynomials
  - Eigenvalues and eigenvectors

#### 2.2 Preliminaries

**Definition 1.** A group is an ordered pair  $(G, \cdot : G \times G \to G)$  where G is a set and  $\cdot$  is a binary operation, which satisfies the following axioms:

- Associativity:  $(g_1g_2)g_3 = g_1(g_2g_3)$ ,
- Identity:  $\exists e \in G \ni ge = eg = g$ ,
- Inverses:  $g \in G \implies \exists h \in G \ni gh = gh = e$ .

#### Example 1.

- $\bullet$   $(\mathbb{Z},+)$
- $(\mathbb{Q}, +)$
- $(\mathbb{Q}^{\times}, \times)$
- $(\mathbb{R}^{\times}, \times)$
- $(GL(n, \mathbb{R}), \times) = \{A \in Mat_n \ni det(A) \neq 0\}$
- $\bullet$   $(S_n, \circ)$

**Definition 2.** A subset  $S \subseteq G$  is a subgroup of G iff

- 1.  $s_1, s_2 \in S \implies s_1 s_2 \in S$
- $2. e \in S$
- $3. \ s \in S \implies s^{-1} \in S$

We denote such a subgroup  $S \leq G$ .

Examples of subgroups:

- $(\mathbb{Z},+) \leq (\mathbb{Q},+)$
- $SL(n,\mathbb{R}) \leq GL(n,\mathbb{R})$ , where  $SL(n,\mathbb{R}) = \{A \in GL(n,\mathbb{R}) \ni \det(A) = 1\}$

#### 2.3 Cyclic Groups

**Definition 3.** A group G is cyclic iff G is generated by a single element.

**Exercise 1.** Show  $\langle g \rangle = \{g^n \ni n \in \mathbb{Z}\} \cong \bigcap \{H \leq G \ni g \in H\}.$ 

**Theorem 4.** Let G be a cyclic group, so  $G\langle g \rangle$ .

- If  $|G| = \infty$ , then  $G \cong \mathbb{Z}$ .
- If  $|G| = n < \infty$ , then  $G \cong \mathbb{Z}_n$ .

**Definition 4.** Let  $H \leq G$ , and define a **right coset of** G by  $aH = \{ah \ni H \in H\}$ . A similar definition can be made for **left cosets**.

Then  $aH = bH \iff b^{-1}a \in G$  and  $Ha = Hb \iff ab^{-1} \in H$ .

Some facts:

- Cosets partition H, i.e.  $b \notin H \implies aH \cap bH = \{e\}$ .
- |H| = |aH| = |Ha| for all  $a \in G$ .

**Theorem 5** (Lagrange). If G is a finite group and  $H \leq G$ , then  $|H| \mid |G|$ .

**Definition 5.** A subgroup  $N \leq G$  is **normal** iff gN = Ng for all  $g \in G$ , or equivalently  $gNg^{-1} \subseteq N$ . I denote this  $N \leq G$ .

When  $N \leq G$ , the set of left/right cosets of N themselves have a group structure. So we define

$$G/N = \{gN \ni g \in G\}$$
 where  $(g_1N)(g_2N) = (g_1g_2)N$ .

Given  $H, K \leq G$ , define  $HK = \{hk \ni h \in H, k \in K\}$ . We have a general formula,

$$|HK| = \frac{|H||K|}{|H \cap K|}.$$

#### 2.4 Homomorphisms

**Definition 6.** Let G, G' be groups, then  $\varphi : G \to G'$  is a homomorphism if  $\varphi(ab) = \varphi(a)\varphi(b)$ .

**Example 2.** •  $\exp: (\mathbb{R}, +) \to (\mathbb{R}^{>0}, \cdot)$  where  $\exp(a+b) = e^{a+b} = e^a e^b = \exp(a) \exp(b)$ .

- det:  $(GL(n,\mathbb{R}),\times) \to (\mathbb{R}^{\times},\times)$  where det(AB) = det(A) det(B).
- Let  $N \leq G$  and  $\varphi G \to G/N$  given by  $\varphi(g) = gN$ .
- Let  $\varphi : \mathbb{Z} \to \mathbb{Z}_n$  where  $\phi(g) = [g] = g \mod n$  where  $\mathbb{Z}_n \cong \mathbb{Z}/n\mathbb{Z}$

**Definition 7.** Let  $\varphi : G \to G'$ . Then  $\varphi$  is a **monomorphism** iff it is injective, an **epimorphism** iff it is surjective, and an **isomorphism** iff it is bijective.

#### 2.5 Direct Products

Let  $G_1, G_2$  be groups, then define

$$G_1 \times G_2 = \{(g_1, g_2) \ni g_1 \in G, g_2 \in G_2\}$$
 where  $(g_1, g_2)(h_1, h_2) = (g_1h_1, g_2, h_2)$ .

We have the formula  $|G_1 \times G_2| = |G_1||G_2|$ .

#### 2.6 Finitely Generated Abelian Groups

**Definition 8.** We say a group is **abelian** if G is commutative, i.e.  $g_1, g_2 \in G \implies g_1g_2 = g_2g_1$ .

**Definition 9.** A group is **finitely generated** if there exist  $\{g_1, g_2, \dots g_n\} \subseteq G$  such that  $G = \langle g_1, g_2, \dots g_n \rangle$ .

This generalizes the notion of a cyclic group, where we can simply intersect all of the subgroups that contain the  $g_i$  to define it.

We know what cyclic groups look like – they are all isomorphic to  $\mathbb{Z}$  or  $\mathbb{Z}_n$ . So now we'd like a structure theorem for abelian finitely generated groups.

**Theorem 6.** Let G be a finitely generated abelian group. Then

$$G \cong \mathbb{Z}^r \times \prod_{i=1}^s \mathbb{Z}_{p_i^{\alpha_i}}$$

for some finite  $r, s \in \mathbb{N}$  and  $p_i$  are (not necessarily distinct) primes.


**Example 3.** Let G be a finite abelian group of order 4. Then  $G \cong \mathbb{Z}_4$  or  $\mathbb{Z}_2^2$ , which are not isomorphic because every element in  $\mathbb{Z}_2^2$  has order 2 where  $\mathbb{Z}_4$  contains an element of order 4.

#### 2.7 Fundamental Homomorphism Theorem

Let  $\varphi: G \to G'$  be a group homomorphism and define  $\ker \varphi = \{g \in G \ni \varphi(g) = e'\}$ .

#### 2.7.1 The First Homomorphism Theorem

**Theorem 7.** There exists a map  $\varphi': G/\ker \varphi \to G'$  such that the following diagram commutes:



That is,  $\varphi = \varphi' \circ \eta$ , and  $\varphi'$  is an isomorphism onto its image, so  $G/\ker \varphi = \operatorname{im} \varphi$ . This map is give by  $\varphi'(g(\ker \varphi)) = \varphi(g)$ .

**Exercise 2.** Check that  $\varphi$  is well-defined.

#### 2.7.2 The Second Theorem

**Theorem 8.** Let  $K, N \leq G$  where  $N \leq G$ . Then

$$\frac{K}{N \bigcap K} \cong \frac{NK}{N}$$

*Proof.* Define a map  $K \xrightarrow{\varphi} NK/N$  by  $\varphi(k) = kN$ . You can show that  $\varphi$  is onto by looking at  $\ker \varphi$ ; note that  $kN = \varphi(k) = N \iff k \in N$ , and so  $\ker \varphi = N \cap K$ .

#### 3 Lecture 2