

Title

Contents

1 Tuesday, October 20	2
1.1 Gluing Two Opens	2
1.2 More General Gluing	3

1 | Tuesday, October 20

1.1 Gluing Two Opens

Recall that a *prevariety* is a ringed space that is locally isomorphic to an affine variety, where we recall that (X, \mathcal{O}_X) is *locally isomorphic* to an affine variety iff there exists an open cover $U_i \rightrightarrows X$ such that (U_i, \mathcal{O}_{U_i}) .

We found one way of producing these: the gluing construction. Given two ringed spaces (X_1, \mathcal{O}_{X_1}) and (X_2, \mathcal{O}_{X_2}) and open sets $U_{12} \in X_1$ and $U_{21} \in X_2$ and an isomorphism $(U_{12}, \mathcal{O}_{U_{12}}) \xrightarrow{f} (U_{21}, \mathcal{O}_{U_{21}})$, we defined

- The topological space as $X_1 \coprod_f X_2$
- The sheaf of rings as $\mathcal{O}_X = \left\{ \varphi : U \rightarrow k \mid \varphi|_{U \cap X_i} \text{ is regular for } i = 1, 2 \right\}$.

Example 1.1.1.

$\mathbb{P}^1/k = X_1 \cup X_2$ where $X_1 \cong \mathbb{A}^1, X_2 \cong \mathbb{A}^1$. Take $U_{12} = D(x)$ and $U_{21} = D(y)$ with

$$f : U_{12} \rightarrow U_{21}$$

$$x \mapsto \frac{1}{x} = y.$$




Figure 1: Supposing $\text{char}(k) \neq 2$. Note that for \mathbb{C} this recovers S^2 in the classical topology.

Example 1.1.2.

Let $X_i = \mathbb{A}^1$ and $U_{12} = D(x), U_{21} = D(y)$ with

$$\begin{aligned} f : U_{12} &\rightarrow U_{21} \\ x &\mapsto x = y. \end{aligned}$$




Figure 2: Line with the doubled origin.

Then $\mathcal{O}_X = \left\{ \varphi : X \rightarrow k \mid \varphi|_{X_i} \text{ is regular} \right\} \cong k[x]$.

1.2 More General Gluing

Now we want to glue more than two open sets. Let I be an indexing set for prevarieties X_i . Suppose that for an ordered pair (i, j) we have open sets $U_{ij} \subset X_i$ and isomorphisms $f_{ij} : U_{ij} \xrightarrow{\sim} U_{ji}$ such that

- a. $f_{ji} = f_{ij}^{-1}$
- b. $f_{jk} \circ f_{ij} = f_{ik}$ (cocycle condition)



Figure 3: Opens with isomorphisms.

Then the gluing construction is given by

1. $X := \coprod X_i / \sim$ where $x \sim f_{ij}(x)$ for all i, j and all $x \in U_{ij}$.
2. $\mathcal{O}_x(U) := \left\{ \varphi : U \rightarrow k \mid \varphi|_{U \cap X_i} \in \mathcal{O}_{X_i} \right\}$.

Every prevariety arises from the gluing construction applied to X_i affine varieties, since a prevariety (X, \mathcal{O}_X) by definition has an open affine cover $X_i \rightrightarrows X$ and X is the result of gluing the X_i s by the identity.

Example 1.2.1.

Let $X_1 = X_2 = X_3 = \mathbb{A}^2/k$. Glue by the following instructions:





Figure 4: The map not shown is whatever formula is necessary to make the diagram commute.

Here

- $(y_1, y_2) = (1/x_1, x_2/x_1)$
- $(z_1, z_2) = (1/x_2, x_1/x_2)$
- $U_{12} = D(x_1)$
- $U_{21} = D(x_2)$.

Figure 5: Yields \mathbb{P}^2

Here $X_1 = [1 : y/x : z/x]$, $X_2 = [x/y : 1 : z/y]$.

Example 1.2.2.

From Gathmann 5.10, open and closed subprevarieties.