

Title

D. Zack Garza

Table of Contents

Contents

Table of Contents	2
1 Lecture 10	3
1.1 Representability and Local Triviality	4

1 | Lecture 10

Remark 1.0.1: What we've been calling a *torsor* (a sheaf with a group action plus conditions) is called by some sources a **pseudotorsor** (e.g. the Stacks Project), and what we've been calling a *locally trivial torsor* is referred to as a *torsor* instead.

Recall that statement of ??; we'll now continue with the proof:

Proof (of Hilbert 90).

Observation 1.0.2: Let $\tau = X_{\text{zar}}, X_{\text{ét}}, X_{\text{fppf}}$, then the data of a GL_n -torsor split by a τ -cover $U \rightarrow X$ is the same as descent data for a vector bundle relative to U/X .

This descent data comes from the following:

$$\begin{array}{c} U \times_X U \\ \pi_1 \downarrow \quad \downarrow \pi_2 \\ U \\ \downarrow \\ X \end{array}$$

That U trivializes our torsor means that $\pi^*T = \pi^*G$ as a G -torsor, where G acts on itself by left-multiplication. We have two different ways of pulling back, and identifications with G in both, yielding

$$\begin{array}{ccc} \pi_1^* \pi^* T & \xrightarrow{\sim} & \pi_2^* \pi^* T \\ \downarrow & & \downarrow \\ \pi_1^* \pi^* G & \xrightarrow{\sim} & \pi_2^* \pi^* G \end{array}$$

Both of the bottom objects are isomorphic to $G|_{U \times U}$.

Claim: The top horizontal map is descent data for T , and the bottom horizontal map is an automorphism of a G -torsor and thus is a section to G . I.e. a section to GL_n is an invertible matrix on double intersections (satisfying the cocycle condition) and a cover, which is precisely descent data for a vector bundle.

Using fppf descent, proved previously, we know that descent data for vector bundles is effective. So if we have a locally trivial GL_n -torsor on the fppf site, it's also trivial on the other two sites, yielding the desired maps back and forth. Thus $H^1(X_{\text{ét}}, \text{GL}_n)$ is in bijection with n -dimensional vector bundles on X . ■

Exercise 1.0.3(?): See if Hilbert 90 is true for groups other than GL_n .

1.1 Representability and Local Triviality

Question 1.1.1: Suppose G is an affine flat X -group scheme. Are all G -torsors representable by a X -scheme?

Answer 1.1.2: Yes, by the same proof as last time, try working out the details. Idea: you can trivialize a G -torsor flat locally and use fppf descent.

Question 1.1.3: Given a G -torsor T that is fppf locally trivial, is it étale locally trivial?

Answer 1.1.4: In general no, but yes if G is smooth.

Proof (Sketch).

You can take an fppf local trivialization, trivialize by p itself, then slice to get an étale trivialization. Given a torsor $T \rightarrow X$, we can base change it to itself:

$$\begin{array}{ccc} T \times_X T & \longrightarrow & T \\ \downarrow \exists \uparrow & & \downarrow \\ T & \longrightarrow & X \end{array}$$

The torsor $T \times_X T \rightarrow T$ is trivial since there exists the indicated section.

■

Example 1.1.5 (non-smooth group schemes):

- α_p , the kernel of Frobenius on \mathbb{A}^1 or \mathbb{G}_a ,
- μ_p in characteristic p , representing p th roots of unity, the kernel of Frobenius on \mathbb{G}_m ,
- The kernel of Frobenius on any positive dimensional affine group scheme.
- $\mu_p \times \mathrm{GL}_n$, etc.