

Title

D. Zack Garza

Friday 20th March, 2020

Contents

1	Friday February 21st	1
1.1	Singularities	1
1.2	Singularities at Infinity	2

1 Friday February 21st

1.1 Singularities

Recall that there are three types of singularities:

- Removable
- Poles
- Essential

Recall that a function g is holomorphic at z_0 iff

$$\lim_{z \rightarrow z_0} (z - z_0)g(z) = 0$$

Theorem 1.1 (3.2).

An isolated singularity z_0 of f is a pole $\iff \lim_{z \rightarrow z_0} f(z) = \infty$.

Theorem 1.2 (3.3, Casorati-Weierstrass).

If f is holomorphic in $D_r(z_0) \setminus \{z_0\}$ and has an essential singularity z_0 , then there exists a radius r such that $f(D_r(\{z_0\}) \setminus \{z_0\})$ is dense in \mathbb{C} .

Proof .

Proceed by contradiction. Suppose there exists a $w \in \mathbb{C}$ and a $\delta > 0$ such that

$$D_\delta(w) \cap f(D_r(\{z_0\}) \setminus \{z_0\}) = \emptyset.$$

If $z \in D_r(w) \setminus z_0$, then $|f(z) - w| > \delta$. Define $g(z) = \frac{1}{f(z) - w}$ on $D_r(z_0) \setminus \{z_0\}$; then $|g(z)| < \frac{1}{\delta}$.

Note that this implies that $g(z)$ is holomorphic on $D_r(z_0) \setminus \{z_0\}$. $g(z)$ being holomorphic here follows from f being holomorphic here.

Then $g(z)$ has a removable singularity at $z = z_0$ by theorem 3.1.

If $g(z_0) \neq 0$, then $f(z) - w$ is holomorphic at z_0 , contradicting the fact that z_0 is an essential singularity.

If instead $g(z_0) = 0$, then z_0 is a pole, again a contradiction. ■

Note: revisit why this is a contradiction.

1.2 Singularities at Infinity

The point $z = \infty$ can be one of three types of singularities:

1. *Removable* $\iff f(z) = \sum_{k=-1}^{\infty} c_k \frac{1}{z^k}$.

- I.e. only one positive exponent.

2. *Pole* $\iff f(z) = \sum_{k=-\infty}^n c_k z^k$

- I.e. there are finitely many positive exponents.

3. *Essential* $\iff f(z) = \sum_{k=-\infty}^{\infty} c_k z^k$

- There are infinitely many positive exponents.

Definition 1.2.1 (Meromorphic).

A function f is **meromorphic** on Ω iff there exists a sequence $\{z_i\} \subset \Omega$ with no limit point in Ω such that

1. f is holomorphic on $\Omega \setminus \{z_i\}$, and
2. f has poles at each z_i .

Theorem 1.3(3.4, Meromorphic Functions are Rational).

f is meromorphic on \mathbb{CP}^1 iff f is a rational function.

Proof.

\implies : By part 1 of the definition above, the point $z = 0$ is either a pole or a removable singularity of the function $F(z) = f\left(\frac{1}{z}\right)$. By part 2, F has finitely many poles. So write

$$f(z) = f_k(z) + g_k(z)$$

where f_k is the principal part and g_k is regular at $z = z_k$.

\impliedby : asdsda

