

# Title

D. Zack Garza

Thursday 20<sup>th</sup> August, 2020

## Contents

### 1 Thursday, August 20

1

### 1 Thursday, August 20

References: <https://www.daniellitt.com/etale-cohomology>

Prerequisites:

- Homological Algebra
  - Abelian Categories
  - Derived Functors
  - Spectral Sequences (just exposure!)
- Sheaf theory and sheaf cohomology
- Schemes (Hartshorne II and III)

Outline/Goals:

- Basics of etale cohomology
    - Etale morphism
    - Grothendieck topologies
    - The etale topology
    - Etale cohomology and the basis theorems
    - Etale cohomology of curves
    - Comparison theorems to singular cohomology
    - Focused on the case where coefficients are a constructible sheaf.
  - Prove the Weil Conjectures (more than one proof)
    - Proving the Riemann Hypothesis for varieties over finite fields
- One of the greatest pieces of 20th century mathematics!
- Topics
    - Weil 2 (Strengthening of RH, used in practice)
    - Formality of algebraic varieties (topological features unique to varieties)
    - Other things (monodromy, refer to Katz' AWS notes)

What is Etale Cohomology? Suppose  $X/\mathbb{C}$  is a quasiprojective variety: a finite type separated integral  $\mathbb{C}$ -scheme.

---

If you take the complex points, it naturally has the structure of a complex analytic space  $X(\mathbb{C})^{\text{an}}$ : you can give it the Euclidean topology, which is much finer than the Zariski topology.

For a nice topological space, we can associate the singular cohomology  $H^i(X(\mathbb{C})^{\text{an}}, \mathbb{Z})$ , which satisfies several nice properties:

- Finitely generated  $\mathbb{Z}$ -modules
- Extra Hodge structure when tensored up to  $\mathbb{C}$  (same as  $\mathbb{C}$  coefficients)
- Cycle classes (i.e. associate to a subvariety a class in cohomology)

Goal of etale cohomology: do something similar for much more general “nice” schemes. Note that some of these properties are special to complex varieties

E.g. finitely generated: not true for a random topological space

We’ll associate  $X$  a “nice scheme”  $\rightsquigarrow H^i(X_{\text{et}}, \mathbb{Z}/\ell^n \mathbb{Z})$ . Take the inverse limit over all  $n$  to obtain the  $\ell$ -adic cohomology  $H^i(X_{\text{et}}, \mathbb{Z}_\ell)$ . You can tensor with  $\mathbb{Q}$  to get something with  $\mathbb{Q}_\ell$  coefficients. And as in singular cohomology, you can a “twisted coefficient system”.

What are nice schemes:

- $X = \text{Spec } \mathcal{O}_k$ , the ring of integers over a number field.
- $X$  a variety over an algebraically closed field
  - Typical, most analogous to taking a variety over  $\mathbb{C}$ .
- $X$  a variety over a non-algebraically closed field

Some comparisons between the last two cases:

- For  $\mathbb{C}$ - variety,  $H_{\text{sing}}^i$  will vanish above  $i = 2d$ .
- Over a finite field,  $H^i$  will vanish for  $i > 2d + 1$  but generally not vanish for  $i = 2d + 1$ .

In good situations, these are finitely generated  $\mathbb{Z}/\ell^n \mathbb{Z}$ -modules, have Mayer-Vietoris and excision sequences, spectral sequences, etc.

Related invariants: for a scheme with a geometric point  $(X, \bar{x}) \rightsquigarrow \pi_1^{\text{étale}}(X, \bar{x})$ , which is a profinite topological group, which is a profinite topological group.

Note: a geometric point is a map from  $\text{Spec } X$  to an algebraically closed field.

More invariants beyond the scope of this course:

- Higher homotopy groups
- Homotopy type (equivalence class of spaces)

So we want homotopy-theoretic invariants for varieties.

**Remark 1.**

This cohomology theory is necessarily weird!

**Theorem 1.1 (Serre).**

There does not exist a cohomology theory for schemes over  $\bar{\mathbb{F}}_q$  with the following properties:

1. Functorial
2. Satisfies the Kunneth formula
3. For  $E$  an elliptic curve,  $H^1(E) = \mathbb{Q}^2$ .

Slogan: No cohomology theory with  $\mathbb{Q}$  coefficients.

---

*Proof .*

Take  $E$  to be a supersingular elliptic curve. Then  $\text{End}(E) \otimes \mathbb{Q}$  is a quaternion algebra.

Fact: There are no algebra morphisms  $R \rightarrow \text{Mat}_{2 \times 2}(\mathbb{Q})$

**Exercise .**

Functoriality and Kunneth implies that  $\text{End}(E) \curvearrowright E$  yields an action on  $H^1(E)$ , which is precisely an algebra morphism  $\text{End}(E) \rightarrow \text{Mat}_{2 \times 2}(\mathbb{Q})$ , a contradiction.

The content: the sum of two endomorphisms act via their sum on  $H^1$ .

**Exercise .**

Prove the same thing for  $\mathbb{Q}_p$  coefficients, where  $p$  divides the characteristic of the ground field.

Proof the same, just need to know what quaternion algebras show up.

■

This forces using some funky type of coefficients.

What are the Weil Conjectures?

Suppose  $X/\mathbb{F}_q$  is a variety, then

$$\zeta_x(t) = \exp \left( \sum_{n>0} \frac{1}{n} |X(\mathbb{F}_{q^n})| t^n \right).$$

Some comments: