

# Title

D. Zack Garza

Friday 25<sup>th</sup> September, 2020

## Contents

1 Friday, September 25

1

# 1 | Friday, September 25

- For  $X, Y$  topological spaces, consider

$$Y^X = C(X, Y) = \text{hom}_{\text{Top}}(X, Y) := \left\{ f : X \rightarrow Y \mid f \text{ is continuous} \right\}.$$

- Topologize with the *compact-open* topology:  $U \in \text{hom}_T(X, X)$  open iff for every  $f \in U$ ,  $f(K)$  is open for every compact  $K \subseteq X$ .
  - \* If  $Y = (Y, d)$  is a metric space, this is the topology of “uniform convergence on compact sets”: for  $f_n \rightarrow f$  in this topology iff

$$\|f_n - f\|_{\infty, K} := \sup \left\{ d(f_n(x), f(x)) \mid x \in K \right\} \xrightarrow{n \rightarrow \infty} 0 \quad \forall K \subseteq X \text{ compact}.$$

In words:  $f_n \rightarrow f$  uniformly on every compact set.

- If  $X$  itself is compact and  $Y$  is a metric space,  $C(X, Y)$  can be promoted to a metric space with  $d(f, g) = \sup_{x \in X} (f(x), g(x))$ .
- So define  $\text{Map}(X, Y) = \text{hom}_{\text{Top}}(X, Y)$  equipped with the compact-open topology.
  - Can immediately consider a lot of interesting spaces by considering  $\text{Map}(\cdot, Y)$ :

$$X = I := [0, 1] \rightsquigarrow P(Y; x_0) := \left\{ f : I \rightarrow Y \mid f(0) = x_0 \right\} = Y^I$$

$$X = S^1 \rightsquigarrow \Omega(Y; x_0) = \mathcal{L}(Y; x_0) := \left\{ f : S^1 \rightarrow Y \right\} = Y^{S^1}.$$

- Importance in homotopy theory: the path space fibration  $\Omega(Y) \hookrightarrow P(Y) \xrightarrow{\gamma \mapsto \gamma(1)} Y$  (plays a role in “homotopy replacement”, allows you to assume everything is a fibration and use homotopy long exact sequences).
- Adjoint property: there is a homeomorphism

$$\text{Map}(X \times Z, Y) \leftrightarrow \cong \text{Map}(Z, \text{Map}(X, Y))$$

$$H : X \times Z \rightarrow Y \iff \tilde{H} : Z \rightarrow \text{Map}(X, Y)$$

$$(x, z) \mapsto H(x, z) \iff z \mapsto H(\cdot, z).$$

---

Categorically,  $\text{hom}(X, \cdot) \leftrightarrow (X \times \cdot)$  form an adjoint pair in  $\text{Top}$ .

- Fun fact: with some mild point-set conditions (Locally compact and Hausdorff),

$$\pi_0 \text{Map}(X, Y) = \{[f], \text{homotopy classes of maps } f : X \rightarrow Y\},$$

i.e. two maps  $f, g$  are homotopic  $\iff$  there is a path in  $\text{Map}(X, Y)$  connecting them.

\* Proof:

$$\text{Map}(I, \text{Map}(X, Y)) \cong \text{Map}(Y \times I, X),$$

and just check that  $\gamma(0) = f \iff H(x, 0) = f$  and  $\gamma(1) = g \iff H(x, 1) = g$ .

- Since these are homeomorphisms, everything is invertible, so equip with function composition to form a group.

–