

Discussion Notes

D. Zack Garza

August 26, 2019

Contents

1 Discussion 1	1
1.1 Uniform Convergence	2

1 Discussion 1

If X is an F_σ set, then

$$X = \bigcup_{i=1}^{\infty} F_i \quad \text{with each } F_i \text{ closed.}$$

If X is a G_δ set, then

$$X = \bigcap_{i=1}^{\infty} G_i \quad \text{with each } G_i \text{ open.}$$

A set A is *nowhere dense* iff $(\overline{A})^\circ = \emptyset$ iff for any interval I , there exists a subinterval S such that $S \cap A = \emptyset$. This is a set that is not dense in any nonempty open set. If the closure of a subset of \mathbb{R} contains no open intervals, it will be nowhere dense.

A set A is *meager* or *first category* if it can be written as

$$A = \bigcup_{i \in \mathbb{N}} A_i \quad \text{with each } A_i \text{ nowhere dense}$$

A set A is *null* if for any ε , there exists a cover of A by countably many intervals of total length less than ε , i.e. there exists $\{I_k\}_{k \in \mathbb{N}}$ such that $A \subseteq \bigcup_{k \in \mathbb{N}} I_k$ and $\sum_{k \in \mathbb{N}} \mu(I_k) < \varepsilon$. If A is null, we say $\mu(A) = 0$.

Some facts:

- If $f_n \rightarrow f$ and each f_n is continuous, then D_f is meager.
- If $f \in \mathcal{R}(a, b)$ and f is bounded, then D_f is null.
- If f is monotone, then D_f is countable.
- If f is monotone and differentiable on (a, b) , then D_f is null.

We define the *oscillation of f* as

$$\omega_f(x) := \lim_{\delta \rightarrow 0^+} \sup_{y, z \in B_\delta(x)} |f(y) - f(z)|$$

1.1 Uniform Convergence

We say that $f_n \rightarrow f$ *converges uniformly on A* if $\|f_n - f\|_\infty = \sup_{x \in A} |f_n(x) - f(x)| \rightarrow 0$. (Note that this defines a sequence of *numbers* in \mathbb{R} .)

This means that one can find an n large enough that for every $x \in A$, we have $|f_n(x) - f(x)| \leq \varepsilon$ for any ε .

- Showing uniform convergence: find some M_n , independent of x , such that $|f_n(x) - f(x)| \leq M_n$ where $M_n \rightarrow 0$.
- Negating: Fix ε , let n be arbitrary, and find a bad x (which can depend on n) such that $|f_n(x) - f(x)| \geq \varepsilon$.

Example: $\frac{1}{1+nx} \rightarrow 0$ pointwise on $(0, \infty)$, which can be seen by fixing x and taking $n \rightarrow \infty$. To see the convergence is not uniform, choose $x = \frac{1}{n}$ and $\varepsilon = \frac{1}{2}$.