

Algebra

D. Zack Garza

August 15, 2019

Contents

1	Lecture 1 (Thu 15 Aug 2019)	1
1.1	Cyclic Groups	2
1.2	Homomorphisms	3
1.3	Direct Products	3
1.4	Finitely Generated Abelian Groups	3
1.5	Fundamental Homomorphism Theorem	4
1.5.1	The First Homomorphism Theorem	4
1.5.2	The Second Theorem	4
2	Lecture 2	4

1 Lecture 1 (Thu 15 Aug 2019)

Definition 1. $\{ \{ \{ \{ \} \} \}$ A **group** is an ordered pair $(G, \cdot : G \times G \rightarrow G)$ where G is a set and \cdot is a binary operation, which satisfies the following axioms:

- Associativity: $(g_1 g_2) g_3 = g_1 (g_2 g_3)$,
- Identity: $\exists e \in G \ni ge = eg = g$,
- Inverses: $g \in G \implies \exists h \in G \ni gh = gh = e$.

$\{ \{ \{ \{ \} \} \}$

Example 1. $\{ \{ \{ \{ \} \} \}$ $(\mathbb{Z}, +)$ - $(\mathbb{Q}, +)$ - $(\mathbb{Q}^\times, \times)$ - $(\mathbb{R}^\times, \times)$ - $(\text{GL}(n, \mathbb{R}), \times) = \{A \in \text{Mat}_n \ni \det(A) \neq 0\}$ - (S_n, \circ)

$\{ \{ \{ \{ \} \} \}$

Definition 2. $\{ \{ \{ \{ \} \} \}$ A subset $S \subseteq G$ is a **subgroup** of G iff

1. $s_1, s_2 \in S \implies s_1 s_2 \in S$
2. $e \in S$
3. $s \in S \implies s^{-1} \in S$

$\{ \{ \{ \{ \} \} \}$

We denote such a subgroup $S \leq G$.

Examples of subgroups:

- $(\mathbb{Z}, +) \leq (\mathbb{Q}, +)$
- $\text{SL}(n, \mathbb{R}) \leq \text{GL}(n, \mathbb{R})$, where $\text{SL}(n, \mathbb{R}) = \{A \in \text{GL}(n, \mathbb{R}) \mid \det(A) = 1\}$

1.1 Cyclic Groups

Definition 3. > {{{{}}} A group G is **cyclic** iff G is generated by a single element.

> {{{{}}}

Exercise 1. > {{{{}}} Show $\langle g \rangle = \{g^n \mid n \in \mathbb{Z}\} \cong \bigcap \{H \leq G \mid g \in H\}$.

> {{{{}}}

Theorem 1. > {{{{}}} Let G be a cyclic group, so $G \cong \langle g \rangle$.

- If $|G| = \infty$, then $G \cong \mathbb{Z}$.
- If $|G| = n < \infty$, then $G \cong \mathbb{Z}_n$.

> {{{{}}}

Definition 4. > {{{{}}} Let $H \leq G$, and define a **right coset** of G by $aH = \{ah \mid h \in H\}$. A similar definition can be made for **left cosets**.

> {{{{}}}

Then $aH = bH \iff b^{-1}a \in H$ and $Ha = Hb \iff ab^{-1} \in H$.

Some facts:

- Cosets partition H , i.e. $b \notin H \implies aH \cap bH = \emptyset$.
- $|H| = |aH| = |Ha|$ for all $a \in G$.

Theorem 2 (Lagrange). > {{{{}}} If G is a finite group and $H \leq G$, then $|H| \mid |G|$.

> {{{{}}}

Definition 5. > {{{{}}} A subgroup $N \leq G$ is **normal** iff $gN = Ng$ for all $g \in G$, or equivalently $gNg^{-1} \subseteq N$. I denote this $N \trianglelefteq G$.

> {{{{}}}

When $N \trianglelefteq G$, the set of left/right cosets of N themselves have a group structure. So we define

$$G/N = \{gN \mid g \in G\} \text{ where } (g_1N)(g_2N) = (g_1g_2)N.$$

Given $H, K \leq G$, define $HK = \{hk \mid h \in H, k \in K\}$. We have a general formula,

$$|HK| = \frac{|H||K|}{|H \cap K|}.$$

1.2 Homomorphisms

Definition 6. > {{{}}} Let G, G' be groups, then $\varphi : G \rightarrow G'$ is a **homomorphism** if $\varphi(ab) = \varphi(a)\varphi(b)$.

> {{{}}}

Example 2. > {{{}}}

- $\exp : (\mathbb{R}, +) \rightarrow (\mathbb{R}^{>0}, \cdot)$ where $\exp(a+b) = e^{a+b} = e^a e^b = \exp(a)\exp(b)$.
- $\det : (\text{GL}(n, \mathbb{R}), \times) \rightarrow (\mathbb{R}^\times, \times)$ where $\det(AB) = \det(A)\det(B)$.
- Let $N \trianglelefteq G$ and $\varphi G \rightarrow G/N$ given by $\varphi(g) = gN$.
- Let $\varphi : \mathbb{Z} \rightarrow \mathbb{Z}_n$ where $\phi(g) = [g] = g \bmod n$ where $\mathbb{Z}_n \cong \mathbb{Z}/n\mathbb{Z}$

> {{{}}}

Definition 7. > {{{}}} Let $\varphi : G \rightarrow G'$. Then φ is a **monomorphism** iff it is injective, an **epimorphism** iff it is surjective, and an **isomorphism** iff it is bijective.

> {{{}}}

1.3 Direct Products

Let G_1, G_2 be groups, then define

$$G_1 \times G_2 = \{(g_1, g_2) \mid g_1 \in G_1, g_2 \in G_2\} \text{ where } (g_1, g_2)(h_1, h_2) = (g_1 h_1, g_2 h_2).$$

We have the formula $|G_1 \times G_2| = |G_1||G_2|$.

1.4 Finitely Generated Abelian Groups

Definition 8. > {{{}}} We say a group is **abelian** if G is commutative, i.e. $g_1, g_2 \in G \implies g_1 g_2 = g_2 g_1$.

> {{{}}}

Definition 9. > {{{}}} A group is **finitely generated** if there exist $\{g_1, g_2, \dots, g_n\} \subseteq G$ such that $G = \langle g_1, g_2, \dots, g_n \rangle$.

> {{{}}}

This generalizes the notion of a cyclic group, where we can simply intersect all of the subgroups that contain the g_i to define it.

We know what cyclic groups look like – they are all isomorphic to \mathbb{Z} or \mathbb{Z}_n . So now we'd like a structure theorem for abelian finitely generated groups.

Theorem 3. > {{{}}} Let G be a finitely generated abelian group. Then

$$G \cong \mathbb{Z}^r \times \prod_{i=1}^s \mathbb{Z}_{p_i^{\alpha_i}}$$

for some finite $r, s \in \mathbb{N}$ and p_i are (not necessarily distinct) primes.

> {{{}}}

Example 3. > {{{}}} Let G be a finite abelian group of order 4. Then $G \cong \mathbb{Z}_4$ or \mathbb{Z}_2^2 , which are not isomorphic because every element in \mathbb{Z}_2^2 has order 2 where \mathbb{Z}_4 contains an element of order 4.

> {{{}}}

1.5 Fundamental Homomorphism Theorem

Let $\varphi : G \rightarrow G'$ be a group homomorphism and define $\ker \varphi = \{g \in G \mid \varphi(g) = e'\}$.

1.5.1 The First Homomorphism Theorem

Theorem 4. > {{{}}} There exists a map $\varphi' : G/\ker \varphi \rightarrow G'$ such that the following diagram commutes:

$$\begin{array}{ccc} G & \xrightarrow{\varphi} & G' \\ \eta \downarrow & \nearrow \varphi' & \\ G/\ker \varphi & & \end{array} \quad >$$

That is, $\varphi = \varphi' \circ \eta$, and φ' is an isomorphism onto its image, so $G/\ker \varphi = \text{im } \varphi$. This map is given by $\varphi'(g(\ker \varphi)) = \varphi(g)$.

> {{{}}}

Exercise 2. > {{{}}} Check that φ is well-defined.

> {{{}}}

1.5.2 The Second Theorem

Theorem 5. > {{{}}} Let $K, N \leq G$ where $N \trianglelefteq G$. Then

$$\frac{K}{N \cap K} \cong \frac{NK}{N}$$

> {{{}}}

Proof. > {{{}}} Define a map $K \xrightarrow{\varphi} NK/N$ by $\varphi(k) = kN$. You can show that φ is onto by looking at $\ker \varphi$; note that $kN = \varphi(k) = N \iff k \in N$, and so $\ker \varphi = N \cap K$. \square

> {{{}}}

2 Lecture 2