

# Title

D. Zack Garza

January 20, 2020

## Contents

<b>1 Chapter 1</b>	<b>1</b>
1.1 Within Chapter . . . . .	1
1.2 End of Chapter Exercises . . . . .	4

## 1 Chapter 1

### 1.1 Within Chapter

Proposition 1.1: Fix an ideal  $\mathfrak{a} \trianglelefteq R$ . There is a correspondence

$$\left\{ \mathfrak{b} \mid \mathfrak{a} \subseteq \mathfrak{b} \trianglelefteq R \right\} \iff \left\{ \tilde{\mathfrak{b}} \trianglelefteq R/\mathfrak{a} \right\}.$$

Proof: Let  $f : R \rightarrow T$  be any ring homomorphism and let  $S(R), S(T)$  denote the sets of subrings of  $R, T$  respectively. Then  $f$  induces two maps:

$$\begin{aligned} \Phi : S(R) &\rightarrow S(T) \\ H &\mapsto f(H) \end{aligned}$$

$$\begin{aligned} \Psi : S(T) &\rightarrow S(R) \\ K &\mapsto f^{-1}(K). \end{aligned}$$

It can be shown that

- $\Phi \circ \Psi(K) = K \bigcap \text{im } f$
- $\Psi \circ \Phi(H) = H \bigcap \ker f$ .

Proposition 1.2: TFAE

- $R$  is a field
- $R$  is simple, i.e. the only ideals of  $R$  are  $0, R$ .

- Every homomorphism  $\phi : R \rightarrow S$  for  $S$  an arbitrary ring is injective.

Proof: ?

Proposition: Maximal ideals are prime.

Proof: ?

Proposition: If  $\mathfrak{p} \trianglelefteq R$  is prime,  $R/\mathfrak{p}$  is a domain. If  $\mathfrak{m} \trianglelefteq R$  is maximal,  $R/\mathfrak{m}$  is a field.

Proof: ?

Theorem 1.3: Every ring  $R$  has a nontrivial maximal ideal  $I \neq 0$ , and every ideal is contained in a maximal ideal.

Proof: ?

Corollary 1.5: Every non-unit of  $R$  is contained in a maximal ideal.

Proof: ?

Proposition 1.6: If  $A \setminus \mathfrak{m} \subset R^\times$ , then  $A$  is a local ring with  $\mathfrak{m}$  its maximal ideal. If  $\mathfrak{m}$  is maximal and  $1 + m \in R^\times$  for all  $m \in \mathfrak{m}$ , then  $A$  is a local ring.

Proof: ?

Proposition: If  $f \in k[x_1, \dots, x_n]$  is irreducible over  $k$ , then  $(f)$  is prime.

Proposition:  $\mathbb{Z}$  is a PID, and  $(p)$  is prime iff  $p$  is zero or a prime number, and every such ideal is maximal.

Proposition:  $k[\{x_i\}]$  has maximal ideals that are not principal iff  $n > 1$ .

Exercise: Characterize the maximal and prime ideals of  $k[x_1, \dots, x_n]$ ? Is this a field, domain, PID, UFD, a local ring, ...?

Proposition: Every nonzero prime ideal in a PID is maximal.

Proof: ?

Definition: The set  $\text{nil}(A)$  of all nilpotent elements in a ring  $A$  is the nilradical of  $A$ . The set  $J(A) = \bigcap_{\mathfrak{m} \in \text{Spec}_{\max}(A)} \mathfrak{m}$  is the Jacobson radical.

Proposition 1.7:  $\text{nil}(A) \trianglelefteq A$  is an ideal and  $A/J(A)$  has no nonzero nilpotent elements.

Proof: ?

Proposition 1.8:  $\text{nil}(A) = \bigcap_{\mathfrak{p} \in \text{Spec}(A)} \mathfrak{p}$  is the intersection of all prime ideals of  $A$ .

Proof: ?

Proposition 1.9:  $x \in J(A)$  iff  $1 - xa \in A^\times$  for all  $a \in A$ .

Proposition: If  $(m), (n) \trianglelefteq \mathbb{Z}$  then  $(m) \cap (n) = (\text{gcd}(m, n))$  and  $(m)(n) = (mn)$ .

Exercise: If  $\mathfrak{a} \trianglelefteq k[x_1, \dots, x_m]$ , characterize  $\mathfrak{a}^n$ .

Exercise: Show that  $\mathfrak{a}, \mathfrak{b} \trianglelefteq A$  are coprime iff there exist  $a \in \mathfrak{a}, b \in \mathfrak{b}$  such that  $a + b = 1$ .

Proposition 1.10: Let  $\{mfa_i\} \trianglelefteq A$  be a family of ideals and define  $\phi : A \rightarrow \prod A/\mathfrak{a}_i$ .

1. If  $\{\mathfrak{a}_i\}$  are pairwise coprime, then  $\prod \mathfrak{a}_i = \bigcap \mathfrak{a}_i$
2.  $\phi$  is surjective iff  $\{\mathfrak{a}_i\}$  are pairwise coprime.
3.  $\phi$  is injective iff  $\bigcap \mathfrak{a}_i = (0)$ .

Exercise: Show that the union of ideals is not necessarily an ideal.

Proposition 1.11:

- a. Let  $\{\mathfrak{p}_i\}$  be a set of prime ideals and let  $\mathfrak{a} \in \bigcup \mathfrak{p}_i$ . Then  $\mathfrak{a} \subseteq \mathfrak{p}_i$  for some  $i$ .
- b. Let  $\{\mathfrak{a}_i\}$  be ideals and  $\mathfrak{p} \supseteq \bigcap \mathfrak{a}_i$  be prime.  $\mathfrak{p} \supseteq \mathfrak{a}_i$  for some  $i$ , and if  $\mathfrak{p} = \bigcap \mathfrak{a}_i$ , then  $\mathfrak{p} = \mathfrak{a}_i$  for some  $i$ .

Exercise: Let  $A = \mathbb{Z}$ , and characterize the ideal quotient  $(m : n)$ .

Exercise 1.12:

1.  $\mathfrak{a} \subseteq (\mathfrak{a} : \mathfrak{b})$
2.  $(\mathfrak{a} : \mathfrak{b})\mathfrak{b} \subseteq \mathfrak{a}$
3.  $((\mathfrak{a} : \mathfrak{b}) : \mathfrak{c}) = (\mathfrak{a} : \mathfrak{b}\mathfrak{c}) = ((\mathfrak{a} : \mathfrak{c}) : \mathfrak{b})$
4.  $(\bigcap \mathfrak{a}_i : \mathfrak{b}) = \bigcap (\mathfrak{a}_i : \mathfrak{b})$
5.  $(\mathfrak{a} : \sum \mathfrak{b}_i) = \bigcap (\mathfrak{a} : \mathfrak{b}_i)$

Proposition: For  $\mathfrak{a} \trianglelefteq A$ ,  $\sqrt{\mathfrak{a}}$  is an ideal.

Exercise 1.13:

1.  $\sqrt{\mathfrak{a}} \supseteq \mathfrak{a}$
2.  $\sqrt{\sqrt{\mathfrak{a}}} = \sqrt{\mathfrak{a}}$
3.  $\sqrt{\mathfrak{a}\mathfrak{b}} = \sqrt{\mathfrak{a} \bigcap \mathfrak{b}} = \sqrt{\mathfrak{a}} \bigcap \sqrt{\mathfrak{b}}$
4.  $\sqrt{\mathfrak{a}} = (1) \iff \mathfrak{a} = (1)$
5.  $\sqrt{\mathfrak{a} + \mathfrak{b}} = \sqrt{\sqrt{\mathfrak{a}} + \sqrt{\mathfrak{b}}}$
6. For  $\mathfrak{p}$  prime,  $\sqrt{\mathfrak{p}^n} = \mathfrak{p}$  for all  $n \geq 1$ .

Proposition 1.14:  $\sqrt{\mathfrak{a}} = \bigcap_{\mathfrak{a} \subseteq \mathfrak{p} \in \text{Spec}(A)} \mathfrak{p}$

Proposition 1.15: Let  $D$  be the set of zero-divisors in  $A$ . Then  $D = \bigcup_{x \neq 0} \sqrt{\text{Ann}(x)}$ .

Exercise: Let  $(m) \trianglelefteq \mathbb{Z}$  where  $m = \prod p_i^{k_i}$ , and show that  $\sqrt{(m)} = (p_1 p_2 \cdots) = \bigcap (p_i)$ .

Proposition 1.16: If  $\sqrt{\mathfrak{a}}, \sqrt{\mathfrak{b}}$  are coprime then  $\mathfrak{a}, \mathfrak{b}$  are coprime.

Exercise: Show that if  $f : A \rightarrow B$  and  $\mathfrak{a} \trianglelefteq A$ , it is not necessarily the case that  $f(\mathfrak{a}) \trianglelefteq B$ .

Exercise: Show that if  $\mathfrak{b}$  is prime then  $A \cdot f^{-1}(\mathfrak{b})$  is prime, but if  $\mathfrak{a}$  is prime then  $B \cdot f(\mathfrak{a})$  need not be prime.

Exercise: Write  $\mathfrak{a}^e := \langle f(\mathfrak{a}) \rangle$  and  $\mathfrak{b}^e = \langle f^{-1}(\mathfrak{b}) \rangle$ . Let  $f : \mathbb{Z} \rightarrow \mathbb{Z}[i]$  be the inclusion, and show that

- $(2)^e = \langle (1+i)^2 \rangle$ , which is not prime in  $\mathbb{Z}[i]$
- (Nontrivial) If  $p \equiv 1 \pmod{4}$ , then  $\mathfrak{p}^e$  is the product of two distinct prime ideals

- If  $p = 3 \pmod{4}$  then  $\mathfrak{p}^e$  is prime.

Proposition: Let  $C = \{\mathfrak{b}^c \mid \mathfrak{b} \trianglelefteq B\}$  and  $E = \{\mathfrak{a}^e \mid \mathfrak{a} \trianglelefteq A\}$ . Then

1.  $\mathfrak{a} \subseteq \mathfrak{a}^{ec}$  and  $\mathfrak{b} \supset \mathfrak{b}^{ce}$ ,
2.  $\mathfrak{b}^c = \mathfrak{b}^{cec}$  and  $\mathfrak{a}^e = \mathfrak{a}^{eee}$
3.  $C = \{\mathfrak{a} \trianglelefteq A \mid \mathfrak{a}^{ec} = \mathfrak{a}\}$  and  $E = \{\mathfrak{b} \trianglelefteq B \mid \mathfrak{b}^{ce} = \mathfrak{b}\}$ .
4. The map  $\phi : C \rightarrow E$  given by  $\phi(\mathfrak{a}) = \mathfrak{a}^{ec}$  is a bijection with inverse  $\mathfrak{b} \mapsto \mathfrak{b}^c$ .
5. If  $\mathfrak{a} \in C$  then  $\mathfrak{a} = \mathfrak{b}^c = \mathfrak{b}^{cec} = \mathfrak{a}^{ec}$ , and if  $\mathfrak{a} = \mathfrak{a}^{ec}$  then  $\mathfrak{a}$  is the contraction of  $\mathfrak{a}^e$ .

Exercise 1.18:

$$\begin{array}{ll}
(\mathfrak{a}_1 + \mathfrak{a}_2)^e = \mathfrak{a}_1^e + \mathfrak{a}_2^e, & (\mathfrak{b}_1 + \mathfrak{b}_2)^c \geq \mathfrak{b}_1^c + \mathfrak{b}_2^c \\
(\mathfrak{a}_1 \cap \mathfrak{a}_2)^e \subseteq \mathfrak{a}_1^e \cap \mathfrak{a}_2^e, & (\mathfrak{b}_1 \cap \mathfrak{b}_2)^c = \mathfrak{b}_1^c \cap \mathfrak{b}_2^c \\
(\mathfrak{a}_1 \mathfrak{a}_2)^e = \mathfrak{a}_1^e \mathfrak{a}_2^e, & (\mathfrak{b}_1 \mathfrak{b}_2)^c \supseteq \mathfrak{b}_1^c \mathfrak{b}_2^c \\
(\mathfrak{a}_1 : \mathfrak{a}_2)^e \subseteq (\mathfrak{a}_1^e : \mathfrak{a}_2^e), & (\mathfrak{b}_1 : \mathfrak{b}_2)^c \subseteq (\mathfrak{b}_1^c : \mathfrak{b}_2^c) \\
r(\mathfrak{a})^e \subseteq r(\mathfrak{a}^e), & r(\mathfrak{b})^c = r(\mathfrak{b}^c)
\end{array}
.$$

## 1.2 End of Chapter Exercises