Interesting Topological Spaces in Algebraic Geometry

D. Zack Garza

Tuesday 28^{th} July, 2020

Contents

1 Ideas for Spaces

- Curves
  - Elliptic Curves
  - Higher genus
  - Hyperelliptic curves
  - The modular curve
- Surfaces
  - Compact Riemann surfaces
    * Bolza Surface (Genus 2)
    * Klein Quartic (Genus 3)
    * Hurwizt Surfaces
  - Kummer surfaces
- Compact Complex Surfaces
  - Rational ruled
  - Enriques Surfaces
  - $K3$
    * Kahler Manifolds
  - Kodaira
  - Toric
  - Hyperelliptic
  - Properly quasi-elliptic
  - General type
  - Type VII
- Fake projective planes
- Conics
- Calabi-Yau manifolds
  - Dimension 1: All elliptic curves (up to homeomorphism)

- Dimension 2: $K3$ surfaces
- Dimension 3 (threefolds): 500 million +, unknown if infinitely many
- The bananafold
- Hyperkähler

- Hurwitz schemes
- Topological galois groups, e.g. $G(\overline{F}/F)$ for $F = \mathbb{Q}, \mathbb{F}_p$.
- Spec (R) for R a DVR (a Sierpinski space)
- Quiver Grassmannians
- Rigid analytic spaces
- Affine line with two origins
- Moduli stack of elliptic curves $\mathcal{M}_{1,1}$.
- Abelian Surface
- Fano Varieties
- Curves: isomorphic to $\mathbb{P}^1$
- Surfaces: Del Pezzo surfaces
- Weighted projective space
- Toric Varieties
- Grassmannian
- Flag Varieties
- Moduli Spaces

> Due to Kunihiko Kodaira's classification of complex surfaces, we know that any compact hyperkähler 4-manifold is either a K3 surface or a compact torus T^{4}. (Every Calabi–Yau manifold in 4 (real) dimensions is a hyperkähler manifold, because SU(2) is isomorphic to Sp(1).)

> As was discovered by Beauville, the Hilbert scheme of k points on a compact hyperkähler 4-manifold is a hyperkähler manifold of dimension 4k. This gives rise to two series of compact examples: Hilbert schemes of points on a K3 surface and generalized Kummer varieties.

2 Analogies

- 2-manifolds: Uniformization
  - Simply connected Riemann surfaces are conformally equivalent to one of $\mathbb{H}, \mathbb{D}^\circ, \mathbb{CP}^1$.
- 3-manifolds: Thurston's Geometrization
  - Oriented prime 3-manifolds can be decomposed into geometric "pieces" of 8 possible types
  - Geometric structure: a diffeo $M \cong \tilde{M}/\Gamma$ where Γ is a discrete Lie group acting freely/transitively on X