

Algebra

D. Zack Garza

September 10, 2019

Contents

| | | |
|----------|---------------------------------------------|-----------|
| 1 | Summary | 2 |
| 2 | Major Theorems | 2 |
| 3 | Lecture 1 (Thu 15 Aug 2019) | 3 |
| 3.1 | Definitions | 3 |
| 3.2 | Preliminaries | 5 |
| 3.3 | Cyclic Groups | 5 |
| 3.4 | Homomorphisms | 6 |
| 3.5 | Direct Products | 6 |
| 3.6 | Finitely Generated Abelian Groups | 6 |
| 3.7 | Fundamental Homomorphism Theorem | 7 |
| 3.7.1 | The First Homomorphism Theorem | 7 |
| 3.7.2 | The Second Theorem | 7 |
| 4 | Lecture 2 | 7 |
| 4.1 | Permutation Groups | 8 |
| 4.2 | Orbits | 8 |
| 4.3 | Groups Acting on Sets | 9 |
| 5 | Lecture 3 (Aug 22) | 10 |
| 5.1 | Burnside's Theorem | 10 |
| 5.2 | Sylow Theory | 11 |
| 5.2.1 | Class Functions | 11 |
| 5.2.2 | Cauchy's Theorem | 12 |
| 5.2.3 | Normalizers | 12 |
| 6 | Appendix | 13 |
| 6.0.1 | Big List of Notation | 13 |
| 7 | Lecture 4: TODO | 13 |
| 8 | Lecture 5 (Tuesday 8/27) | 13 |
| 8.1 | Sylow Theorems | 13 |
| 8.1.1 | Sylow 1 | 13 |
| 8.1.2 | Sylow 2 | 14 |

| | | |
|-----------|---------------------------------------------|-----------|
| 8.1.3 | Sylow 3 | 14 |
| 8.1.4 | Applications | 15 |
| 8.2 | Classification of groups of a certain order | 16 |
| 9 | Lecture 6 | 16 |
| 9.1 | Internal Direct Products | 16 |
| 9.2 | Determination of groups of a given order | 17 |
| 9.3 | Free Groups | 17 |
| 9.4 | Generators and Relations | 18 |
| 10 | Lecture 7 (Thursday 29th) | 19 |
| 10.1 | Groups of Order 6 | 19 |
| 10.2 | Groups of Order 8 | 21 |
| 10.3 | Some Nice Facts | 21 |
| 10.4 | Simple Groups | 22 |
| 10.5 | Series of Groups | 22 |
| 11 | Lecture 8: Series of Groups | 23 |
| 11.1 | The Commutator Subgroup | 24 |
| 11.2 | Free Abelian Groups | 25 |
| 12 | Another Lecture: On to Rings | 25 |

1 Summary

Groups and rings, including Sylow theorems, classifying small groups, finitely generated abelian groups, Jordan-Holder theorem, solvable groups, simplicity of the alternating group, euclidean domains, principal ideal domains, unique factorization domains, noetherian rings, Hilbert basis theorem, Zorn's lemma, and existence of maximal ideals and vector space bases.

Previous course web pages:

- Fall 2017, Asilata Bapat

2 Major Theorems

Theorem 1 (Cauchy). For any prime p dividing the order of G , there is an element x of order p (and thus a subgroup $H = \langle x \rangle$ of order p as well).

Theorem 2 (Lagrange). If $H \leq G$ is a subgroup, then $|H| \mid |G|$. Moreover,

$$|G| = [G : H] |H|.$$

Theorem 3 (Sylow 1). If $|G| = n = \prod p_i^{a_i}$ as a prime factorization, then G has subgroups of order $p_i^{a_i}$ for every i and for every $1 \leq r \leq a_i$. In particular, $\text{Syl}(p, G) \neq \emptyset$.

Moreover, every subgroup H of order p^k is normal in a subgroup of order p^{k+1} for $1 \leq k \leq \alpha_i$, and thus $H \leq P$ for some $P \in \text{Syl}(p, G)$.

Theorem 4 (Sylow 2). If $P_1, P_2 \in \text{Syl}(p, G)$, then there exists a $g \in G$ such that $gP_1g = P_2$.

Theorem 5 (Sylow 3). Let $|G| = p^n m$ and $r_p = |\text{Syl}(p, G)|$. Then

- $r_p \equiv 1 \pmod{p}$,
- $r_p \mid m$,
- $r_p = [G : N_G(P)]$.

Theorem 6 (Classification of finitely generated abelian groups). If G is a finitely generated abelian group, then $G \cong F \oplus T$, where F is free abelian and T is a torsion group. If $|T| = n$, then $T \cong \bigoplus \mathbb{Z}_{p_i^{\alpha_i}}$ where $n = \prod p_i^{\alpha_i}$ is some factorization of n with the p_i **not necessarily distinct**.

Theorem 7. Conjugacy classes partition G

$$|G| = |Z(G)| + \sum_{\text{One representative in each orbit}} |C_G(g_i)| = \sum_{\text{asdsa}} [G : C(g_i)].$$

Theorem 8 (Orbit Stabilizer). If $G \curvearrowright X$, then for any $x \in X$

$$[G : \text{Stab}(x)] = |\mathcal{O}_x|, \quad \text{i.e.} \quad |G| = |\mathcal{O}_x| |\text{Stab}(x)|$$

where $\mathcal{O}_x = \{g \curvearrowright x \mid g \in G\} \subseteq X$ and $\text{Stab}(x) = \{x \in X \mid \forall g \in G, g \curvearrowright x = x\} \leq G$.

Some nice lemmas:

- Every subgroup of a cyclic group is itself cyclic.
- $aH = bH \iff b^{-1}a \in H$.
- $A \leq G$ and $B \leq G \implies (A \cap B) \leq G$.
 - Corollary: $\#A = p, \#B = q \implies A \cap B = \{e\}$.
 - Corollary: $\#A = p, \#B = p \implies A = B$ or $A \cap B = \{e\}$.

3 Lecture 1 (Thu 15 Aug 2019)

We'll be using Hungerford's Algebra text.

3.1 Definitions

The following definitions will be useful to know by heart:

- The order of a group
- Cartesian product
- Relations
- Equivalence relation
- Partition
- Binary operation
- Group

- Isomorphism
- Abelian group
- Cyclic group
- Subgroup
- Greatest common divisor
- Least common multiple
- Permutation
- Transposition
- Orbit
- Cycle
- The symmetric group S^n
- The alternating group A_n
- Even and odd permutations
- Cosets
- Index
- The direct product of groups
- Homomorphism
- Image of a function
- Inverse image of a function
- Kernel
- Normal subgroup
- Factor group
- Simple group

Here is a rough outline of the course:

- Group Theory
 - Groups acting on sets
 - Sylow theorems and applications
 - Classification
 - Free and free abelian groups
 - Solvable and simple groups
 - Normal series
- Galois Theory
 - Field extensions
 - Splitting fields
 - Separability
 - Finite fields
 - Cyclotomic extensions
 - Galois groups
 - Solvability by radicals
- Module theory
 - Free modules
 - Homomorphisms
 - Projective and injective modules
 - Finitely generated modules over a PID
- Linear Algebra
 - Matrices and linear transformations
 - Rank and determinants

- Canonical forms
- Characteristic polynomials
- Eigenvalues and eigenvectors

3.2 Preliminaries

Definition 9. A **group** is an ordered pair $(G, \cdot : G \times G \rightarrow G)$ where G is a set and \cdot is a binary operation, which satisfies the following axioms:

- Associativity: $(g_1 g_2) g_3 = g_1 (g_2 g_3)$,
- Identity: $\exists e \in G \ni ge = eg = g$,
- Inverses: $g \in G \implies \exists h \in G \ni gh = gh = e$.

Example 10.

- $(\mathbb{Z}, +)$
- $(\mathbb{Q}, +)$
- $(\mathbb{Q}^\times, \times)$
- $(\mathbb{R}^\times, \times)$
- $(\text{GL}(n, \mathbb{R}), \times) = \{A \in \text{Mat}_n \ni \det(A) \neq 0\}$
- (S_n, \circ)

Definition 11. A subset $S \subseteq G$ is a **subgroup** of G iff

1. $s_1, s_2 \in S \implies s_1 s_2 \in S$
2. $e \in S$
3. $s \in S \implies s^{-1} \in S$

We denote such a subgroup $S \leq G$.

Examples of subgroups:

- $(\mathbb{Z}, +) \leq (\mathbb{Q}, +)$
- $\text{SL}(n, \mathbb{R}) \leq \text{GL}(n, \mathbb{R})$, where $\text{SL}(n, \mathbb{R}) = \{A \in \text{GL}(n, \mathbb{R}) \ni \det(A) = 1\}$

3.3 Cyclic Groups

Definition 12. A group G is **cyclic** iff G is generated by a single element.

Exercise 1. Show $\langle g \rangle = \{g^n \ni n \in \mathbb{Z}\} \cong \bigcap_{g \in G} \{H \ni H \leq G \text{ and } g \in H\}$.

Theorem 13. Let G be a cyclic group, so $G = \langle g \rangle$.

- If $|G| = \infty$, then $G \cong \mathbb{Z}$.
- If $|G| = n < \infty$, then $G \cong \mathbb{Z}_n$.

Definition 14. Let $H \leq G$, and define a **right coset** of G by $aH = \{ah \ni H \in H\}$. A similar definition can be made for **left cosets**.

Then $aH = bH \iff b^{-1}a \in H$ and $Ha = Hb \iff ab^{-1} \in H$.

Some facts:

- Cosets partition H , i.e. $b \notin H \implies aH \cap bH = \{e\}$.
- $|H| = |aH| = |Ha|$ for all $a \in G$.

Theorem 15 (Lagrange). If G is a finite group and $H \leq G$, then $|H| \mid |G|$.

Definition 16. A subgroup $N \leq G$ is **normal** iff $gN = Ng$ for all $g \in G$, or equivalently $gNg^{-1} \subseteq N$. I denote this $N \trianglelefteq G$.

When $N \trianglelefteq G$, the set of left/right cosets of N themselves have a group structure. So we define

$$G/N = \{gN \mid g \in G\} \text{ where } (g_1N)(g_2N) = (g_1g_2)N.$$

Given $H, K \leq G$, define $HK = \{hk \mid h \in H, k \in K\}$. We have a general formula,

$$|HK| = \frac{|H||K|}{|H \cap K|}.$$

3.4 Homomorphisms

Definition 17. Let G, G' be groups, then $\varphi : G \rightarrow G'$ is a **homomorphism** if $\varphi(ab) = \varphi(a)\varphi(b)$.

Example 18. • $\exp : (\mathbb{R}, +) \rightarrow (\mathbb{R}^{>0}, \cdot)$ where $\exp(a+b) = e^{a+b} = e^a e^b = \exp(a) \exp(b)$.

- $\det : (\text{GL}(n, \mathbb{R}), \times) \rightarrow (\mathbb{R}^\times, \times)$ where $\det(AB) = \det(A) \det(B)$.
- Let $N \trianglelefteq G$ and $\varphi G \rightarrow G/N$ given by $\varphi(g) = gN$.
- Let $\varphi : \mathbb{Z} \rightarrow \mathbb{Z}_n$ where $\phi(g) = [g] = g \bmod n$ where $\mathbb{Z}_n \cong \mathbb{Z}/n\mathbb{Z}$

Definition 19. Let $\varphi : G \rightarrow G'$. Then φ is a **monomorphism** iff it is injective, an **epimorphism** iff it is surjective, and an **isomorphism** iff it is bijective.

3.5 Direct Products

Let G_1, G_2 be groups, then define

$$G_1 \times G_2 = \{(g_1, g_2) \mid g_1 \in G_1, g_2 \in G_2\} \text{ where } (g_1, g_2)(h_1, h_2) = (g_1h_1, g_2h_2).$$

We have the formula $|G_1 \times G_2| = |G_1||G_2|$.

3.6 Finitely Generated Abelian Groups

Definition 20. We say a group is **abelian** if G is commutative, i.e. $g_1, g_2 \in G \implies g_1g_2 = g_2g_1$.

Definition 21. A group is **finitely generated** if there exist $\{g_1, g_2, \dots, g_n\} \subseteq G$ such that $G = \langle g_1, g_2, \dots, g_n \rangle$.

This generalizes the notion of a cyclic group, where we can simply intersect all of the subgroups that contain the g_i to define it.

We know what cyclic groups look like – they are all isomorphic to \mathbb{Z} or \mathbb{Z}_n . So now we'd like a structure theorem for abelian finitely generated groups.

Theorem 22. Let G be a finitely generated abelian group. Then

$$G \cong \mathbb{Z}^r \times \prod_{i=1}^s \mathbb{Z}_{p_i^{\alpha_i}}$$

for some finite $r, s \in \mathbb{N}$ and p_i are (not necessarily distinct) primes.

Example 23. Let G be a finite abelian group of order 4. Then $G \cong \mathbb{Z}_4$ or \mathbb{Z}_2^2 , which are not isomorphic because every element in \mathbb{Z}_2^2 has order 2 where \mathbb{Z}_4 contains an element of order 4.

3.7 Fundamental Homomorphism Theorem

Let $\varphi : G \rightarrow G'$ be a group homomorphism and define $\ker \varphi = \{g \in G \mid \varphi(g) = e'\}$.

3.7.1 The First Homomorphism Theorem

Theorem 24. There exists a map $\varphi' : G/\ker \varphi \rightarrow G'$ such that the following diagram commutes:

$$\begin{array}{ccc} G & \xrightarrow{\varphi} & G' \\ \eta \downarrow & \nearrow \varphi' & \\ G/\ker \varphi & & \end{array}$$

That is, $\varphi = \varphi' \circ \eta$, and φ' is an isomorphism onto its image, so $G/\ker \varphi = \text{im } \varphi$. This map is given by $\varphi'(g(\ker \varphi)) = \varphi(g)$.

Exercise 2. Check that φ is well-defined.

3.7.2 The Second Theorem

Theorem 25. Let $K, N \leq G$ where $N \trianglelefteq G$. Then

$$\frac{K}{N \cap K} \cong \frac{NK}{N}$$

Proof. Define a map $K \xrightarrow{\varphi} NK/N$ by $\varphi(k) = kN$. You can show that φ is onto, then look at $\ker \varphi$; note that $kN = \varphi(k) = N \iff k \in N$, and so $\ker \varphi = N \cap K$. \square

4 Lecture 2

Last time: the fundamental homomorphism theorems.

Theorem 1: Let $\varphi : G \rightarrow G'$ be a homomorphism. Then there is a canonical homomorphism $\eta : G \rightarrow G/\ker \varphi$ such that the usual diagram commutes. Moreover, this map induces an isomorphism $G/\ker \varphi \cong \text{im } \varphi$.

Theorem 2: Let $K, N \leq G$ and suppose $N \trianglelefteq G$. Then there is an isomorphism

$$\frac{K}{K \cap N} \cong \frac{KN}{N}$$

(Show that $K \cap N \trianglelefteq K$, and KN is a subgroup exactly because N is normal).

Theorem 3: Let $H, K \trianglelefteq G$ such that $H \leq K$.

1. H/K is normal in G/K .

2. The quotient $(G/K)/(H/K) \cong G/H$.

Proof: We'll use the first theorem. First make a map

$$\begin{aligned} G/K &\rightarrow G/H \\ \phi(gk) &= gH \end{aligned}$$

Exercise: Show that this map is onto, and that $\ker \phi \cong H/K$.

4.1 Permutation Groups

Let A be a set, then a *permutation* on A is a bijective map $A \rightarrow A$. This can be made into a group with a binary operation given by composition of functions. Denote S_A the set of permutations on A .

Theorem: S_A is in fact a group. Check associativity, inverses, identity, etc.

In the special case that $A = \{1, 2, \dots, n\}$, then $S_n := S_A$.

Recall two line notation

$$\begin{pmatrix} 1 & 2 & \cdots & n \\ \sigma(1) & \sigma(2) & \cdots & \sigma(n) \end{pmatrix}$$

Moreover, $|S_n| = n!$ by a combinatorial counting argument.

Example: S_3 is the symmetries of a triangle (see notes).

Example: The symmetries of a square are *not* given by S_4 , it is instead D_4 (see notes).

4.2 Orbits

Permutations S_A “acts” on A , and if $\sigma \in S_A$, then $\langle \sigma \rangle$ also acts on A .

Define $a \sim b$ iff there is some n such that $\sigma^n(a) = b$. This is an equivalence relation, and thus induces a partition of A . See notes for diagram. The equivalence classes under this relation are called the *orbits* under σ .

Example:

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 8 & 2 & 6 & 3 & 7 & 4 & 5 & 1 \end{pmatrix} = (18)(2)(364)(57).$$

Definition: A permutation $\sigma \in S_n$ is a *cycle* iff it contains at most one orbit with more than one element. The *length* of a cycle is the number of elements in the largest orbit.

Recall cycle notation: $\sigma = (\sigma(1)\sigma(2)\cdots\sigma(n))$. Note that this is read right-to-left by convention!

Theorem: Every permutation $\sigma \in S_n$ can be written as a product of disjoint cycles.

Definition: A *transposition* is a cycle of length 2. Moreover, we have

$$(a_1 a_2 \cdots a_n) = (a_1 a_n)(a_1 a_{n-1}) \cdots (a_1 a_2),$$

and so every permutation is a product of transpositions. This is not a unique decomposition, however, as e.g. $\text{id} = (12)^2 = (34)^2$.

Theorem: Any $\sigma \in S_n$ can be written as **either** an even number of transpositions or an odd number of transpositions.

Define $A_n = \{\sigma \in S_n \mid \sigma \text{ is even}\}$. We claim that $A_n \trianglelefteq S_n$.

1. Closure: If τ_1, τ_2 are both even, then $\tau_1\tau_2$ also has an even number of transpositions.
2. The identity has an even number of transpositions, since zero is even.
3. Inverses: If $\sigma = \prod_{i=1}^s \tau_i$ where s is even, then $\sigma^{-1} = \prod_{i=1}^s \tau_{s-i}$. But each τ is order 2, so $\tau^{-1} = \tau$, so there are still an even number of transpositions.

So A_n is a subgroup. It is normal because it is index 2, or the kernel of a homomorphism, or by a direct computation.

4.3 Groups Acting on Sets

Think of this as a generalization of a G -module.

Definition: A group G is said to *act* on a set X if there exists a map $G \times X \rightarrow X$ such that

1. $e \curvearrowright x = x$
2. $(g_1g_2) \curvearrowright x = g_1 \curvearrowright (g_2 \curvearrowright x)$.

Examples:

1. $G = S_A \curvearrowright A$
2. $H \leq G$, then $G \curvearrowright X = G/H$ where $g \curvearrowright xH = (gx)H$.
3. $G \curvearrowright G$ by conjugation, i.e. $g \curvearrowright x = gxg^{-1}$.

Definition: Let $x \in X$, then define the *stabilizer subgroup*

$$G_x = \{g \in G \mid g \curvearrowright x = x\} \leq G$$

We can also look at the dual thing,

$$X_g = \{x \in X \mid g \curvearrowright x = x\}.$$

We then define the *orbit* of an element x as

$$Gx = \{g \curvearrowright x \mid g \in G\}$$

and we have a similar result where $x \sim y \iff x \in Gy$, and the orbits partition X .

Theorem: Let G act on X . We want to know the number of elements in an orbit, and it turns out that

$$|Gx| = [G : G_x] \tag{1}$$

Proof: Construct a map $Gx \xrightarrow{\psi} G/G_x$ where $\psi(g \curvearrowright x) = gG_x$. Exercise: Show that this is well-defined, so if 2 elements are equal then they go to the same coset. Exercise: Show that this is surjective.

Injectivity: $\psi(g_1x) = \psi(g_2x)$, so $g_1Gx = g_2Gx$ and $(g_2^{-1}g_1)Gx = Gx$ so $g_2^{-1}g_1 \in Gx \iff g_2^{-1}g_1 \curvearrowright x = x \iff g_1x = g_2x$.

Next time: Burnside's theorem, proving the Sylow theorems.

5 Lecture 3 (Aug 22)

Last time: let G be a group and X be a set; we say G acts on X (or that X is a G -set) when there is a map $G \times X \rightarrow X$ such that $ex = x$ and $(gh) \curvearrowright x = g \curvearrowright (h \curvearrowright x)$. We then define the stabilizer of x as

$$G_x = \{g \in G \mid g \curvearrowright x = x\} \leq G,$$

and the orbit

$$G.x = \mathcal{O}_x = \{g \curvearrowright x \mid x \in X\} \subseteq X.$$

When G is finite, we have

$$\#G.x = \frac{\#G}{\#G_x}.$$

We can also consider the fixed points of X ,

$$X_g = \{x \in X \mid g \curvearrowright x = x \forall g \in G\} \subseteq X$$

5.1 Burnside's Theorem

Theorem (Burnside): Let X be a G -set and v be the number of orbits. Then

$$v\#G = \sum_{g \in G} \#X_g.$$

Proof:

Define $N = \{(g, x) \mid g \curvearrowright x = x\} \subseteq G \times X$, we then have

$$\begin{aligned} |N| &= \sum_{g \in G} |X_g| \\ &= \sum_{x \in X} |G_x| \\ &= \sum_{x \in X} \frac{|G|}{|G.x|} \\ &= |G| \left(\sum_{x \in X} \frac{1}{|G.x|} \right) \\ &= |G|v. \end{aligned}$$

Since the orbits partition X , say into $X = \bigcup_{i=1}^v \sigma_i$, let $\sigma = \{\sigma_i \ni 1 \leq i \leq v\}$ and abuse notation slightly by replacing each orbit in σ with a representative element $x_i \in \sigma_i \subset X$. We then have

$$\sum_{x \in \sigma} \frac{1}{|G \cdot x|} = \frac{1}{|Gx|} |\sigma| = 1.$$

Application: Consider seating 10 people around a circular table. How many distinct seating arrangements are there?

Let X be the set of configurations, $G = S_{10}$, and let $G \curvearrowright X$ by permuting configurations. Then v , the number of orbits under this action, yields the number of distinct seating arrangements. By Burnside, we have

$$v = \frac{1}{|G|} \sum_{g \in G} |Xg| = \frac{1}{10} (10!) = 9!,$$

since $Xg = \{x \in X \ni gx = x\} = \emptyset$ unless $g = e$, and $X_e = X$.

5.2 Sylow Theory

Recall Lagrange's theorem: If $H \leq G$ and G is finite, then $\#H \mid \#G$.

Consider the converse: if $n \mid \#G$, does there exist a subgroup of size n ? The answer is no in general, and a counterexample is A_4 which has $4!/2 = 12$ elements but no subgroup of order 6.

5.2.1 Class Functions

Let X be a G -set, and choose orbit representatives $x_1 \cdots x_v$. Then

$$|X| = \sum_{i=1}^v |Gx_i|.$$

We can then separately count all orbits with exactly one element, which is exactly $X_G = \{x \in G \ni g \curvearrowright x = x \forall g\}$

We then have

$$|X| = |X_G| + \sum_{i=j}^v$$

for some j where $|Gx_i| > 1$ for all $i \geq j$.

Theorem: Let G be a group of order p^n for p a prime, then

$$|X| \equiv |X_G| \pmod{p}$$

Proof: We know that $|Gx_i| = [G : Gx_i]$ for $j \leq i \leq v$, and $|Gx_i| > 1$ implies that $Gx_i \neq G$ and thus $p \mid [G : Gx_i]$. The result follows.

Application: If $|G| = p^n$, then the center $Z(G)$ is nontrivial. Let $X = G$ act on itself by conjugation, so $g \curvearrowright x = gxg^{-1}$. Then

$$X_G = \{x \in G \ni gxg^{-1} = x\} = \{x \in G \ni gx = xg\} = Z(G)$$

But then, by the previous theorem, we have $|Z(G)| \equiv |X| \equiv |G| \pmod{p}$, but since $Z(G) \leq G$ we have $|Z(G)| \equiv 0 \pmod{p}$, and so in particular, $Z(G) \neq \{e\}$.

Definition: A group G is a p -group iff every element in G has order p^k for some k . A subgroup is a p -group exactly when it is a p -group in its own right.

5.2.2 Cauchy's Theorem

Theorem (Cauchy): Let G be a finite group, where $p \mid |G|$ is a prime. Then G is an element (and thus a subgroup) of order p .

Proof: Consider $X = \{(g_1, g_2, \dots, g_p) \in G^{\oplus p} \mid g_1 g_2 \cdots g_p = e\}$. Given any $p - 1$ elements, say $g_1 \cdots g_{p-1}$, the remaining element is completely determined by $g_p = (g_1 \cdots g_{p-1})^{-1}$. So $|X| = |G|^{p-1}$.

Since $p \mid |G|$, we have $p \mid |X|$. Now let $\sigma \in S_p$ the symmetric group act on X by index permutation, i.e. $\sigma \curvearrowright (g_1, g_2, \dots, g_p) = (g_{\sigma(1)}, g_{\sigma(2)}, \dots, g_{\sigma(p)})$.

Exercise: Check that this gives a well-defined group action.

Let $\sigma = (1 \ 2 \ \cdots \ p) \in S_p$, and note $\langle \sigma \rangle \leq S_p$ also acts on X where $|\langle \sigma \rangle| = p$. Therefore we have

$$|X| = |X_{\langle \sigma \rangle}| \pmod{p}.$$

Since $p \mid |X|$, it follows that $|X_{\langle \sigma \rangle}| \equiv 0 \pmod{p}$, and thus $p \mid |X_{\langle \sigma \rangle}|$.

If $\langle \sigma \rangle$ fixes (g_1, g_2, \dots, g_p) , then $g_1 = g_2 = \cdots = g_p$.

Note that $(e, e, \dots) \in X_{\langle \sigma \rangle}$, as is (a, a, \dots, a) since $p \mid |X_{\langle \sigma \rangle}|$. So there is some $a \in G$ such that $a^p = 1$. Moreover, $\langle a \rangle \leq G$ is a subgroup of size p .

5.2.3 Normalizers

Let G be a group and $X = S$ be the set of subgroups of G . Let G act on X by $g \curvearrowright H = gHg^{-1}$. What is the stabilizer? $G_x = G_H = \{g \in G \mid gHg^{-1} = H\}$, making G_H the largest subgroup such that $H \trianglelefteq G_H$. So we define $N_G(H) = G_H$.

Lemma: Let H be a p -subgroup of G of order p^n . Then $[N_G(H) : H] \equiv [G : H] \pmod{p}$.

Proof: Let $S = G/H$ be the set of left H -cosets in G . Now let H act on S by $H \curvearrowright x + H = (hx) + H$.

By a previous theorem, $|G/H| = |S| = |S_H| \pmod{p}$, where $|G/H| = [G : H]$. What is S_H ? Thus is given by $S_H = \{x + H \in S \mid xHx^{-1} \in H \forall h \in H\}$. Therefore $x \in N_G(H)$.

Corollary: Let $H \leq G$ be a subgroup of order p^n . If $p \mid [G : H]$ then $N_G(H) \neq H$. Proof: Exercise.

Theorem: Let G be a finite group, then G is a p -group iff $|G| = p^n$.

Proof: Suppose $|G| = p^n$ and $a \in G$. Then $|\langle a \rangle| = p^\alpha$ for some α . Conversely, suppose G is a p -group. Factor $|G|$ into primes and suppose $\exists q$ such that $q \mid |G|$ but $q \neq p$. By Cauchy, we can then get a subgroup $\langle c \rangle$ such that $|\langle c \rangle| \mid q$, but then $|G| \neq p^n$.

6 Appendix

6.0.1 Big List of Notation

| | | | |
|-------------------|-----------------------------------------------|---------------------------------|-----------------|
| $C(x) =$ | $\{g \in G : gxg^{-1} = x\}$ | $\subseteq G$ | Centralizer |
| $C_G(x) =$ | $\{gxg^{-1} : g \in G\}$ | $\subseteq G$ | Conjugacy Class |
| $G_x =$ | $\{g.x : x \in X\}$ | $\subseteq X$ | Orbit |
| $x_0 =$ | $\{g \in G : g.x = x\}$ | $\subseteq G$ | Stabilizer |
| $Z(G) =$ | $\{x \in G : \forall g \in G, gxg^{-1} = x\}$ | $\subseteq G$ | Center |
| $\text{Inn}(G) =$ | $\{\phi_g(x) = gxg^{-1}\}$ | $\subseteq \text{Aut}(G)$ | Inner Aut. |
| $\text{Out}(G) =$ | $\text{Aut}(G)/\text{Inn}(G)$ | $\hookrightarrow \text{Aut}(G)$ | Outer Aut. |
| $N(H) =$ | $\{g \in G : gHg^{-1} = H\}$ | $\subseteq G$ | Normalizer |

7 Lecture 4: TODO

8 Lecture 5 (Tuesday 8/27)

Let G be a finite group and p a prime. TFAE:

- $|H| = p^n$ for some n
- Every element of H has order p^α for some α .

If either of these are true, we say H is a p -group.

Let H be a p -group, last time we proved that if $p \mid [G : H]$ then $N_G(H) \neq H$.

8.1 Sylow Theorems

Let G be a finite group and suppose $|G| = p^n m$ where $(m, p) = 1$. Then

8.1.1 Sylow 1

Motto: take a prime factorization of $|G|$, then there are subgroups of order p^i for *every* prime power appearing, up to the maximal power.

1. G contains a subgroup of order p^i for every $1 \leq i \leq n$.
2. Every subgroup H of order p^i where $i < n$ is a normal subgroup in a subgroup of order p^{i+1} .

Proof: By induction on i . For $i = 1$, we know this by Cauchy's theorem. If we show (2), that shows (1) as a consequence. So suppose this holds for $i < n$. Let $H \leq G$ where $|H| = p^i$, we now want a subgroup of order p^{i+1} . Since $p \mid [G : H]$, by the previous theorem, $H < N_G(H)$ is a proper subgroup (?).

Now consider the canonical projection $N_G(H) \rightarrow N_G(H)/H$. Since $p \mid [N_G(H) : H] = |N_G(H)/H|$, by Cauchy there is a subgroup of order p in this quotient. Call it K . Then $\pi^{-1}(K) \leq N_G(H)$.

Exercise: $|\phi^{-1}(K)| = p^{i+1}$.

It now follows that $H \trianglelefteq \phi^{-1}(K)$. \square

Definition: For G a finite group and $|G| = p^n m$ where $p \nmid m$. Then a subgroup of order p^n is called a Sylow p -subgroup. (By Sylow 1, these exist.)

8.1.2 Sylow 2

If P_1, P_2 are Sylow p -subgroups of G , then P_1 and P_2 are conjugate.

Proof: Let \mathcal{L} be the left cosets of P_1 , i.e. $\mathcal{L} = G/P_1$. Then let P_2 act on \mathcal{L} by $p_2 \curvearrowright (g + P_1) := (p_2 g) + P_1$.

By a previous theorem about orbits and fixed points, we have

$$|\mathcal{L}_{P_2}| = |\mathcal{L}| \pmod{p}.$$

Since $p \nmid |\mathcal{L}|$, we have $p \nmid |\mathcal{L}_{P_2}|$. So \mathcal{L}_{P_2} is nonempty.

So there exists a coset xP_1 such that $xP_1 \in \mathcal{L}_{P_2}$, and so $yxP_1 = xP_1$ for all $y \in P_2$.

Then $x^{-1}yxP_1 = P_1$ for all $y \in P_2$, and so $x^{-1}P_2x = P_1$. But then P_1 and P_2 are conjugate. \square

8.1.3 Sylow 3

Let G be a finite group, and $p \mid |G|$. Let r_p be the number of Sylow p -subgroups of G . Then

- $r_p \equiv 1 \pmod{p}$.
- $r_p \mid |G|$.
- $r_p = [G : N_G(P)]$

Let $X = \mathcal{S}$ be the set of Sylow p -subgroups, and let $P \in X$ be a fixed Sylow p -subgroup. Let $P \curvearrowright \mathcal{S}$ by conjugation, so for $\bar{P} \in \mathcal{S}$ let $x \curvearrowright \bar{P} = x\bar{P}x^{-1}$.

By the same old theorem, we have

$$|\mathcal{S}| = \mathcal{S}_P \pmod{p}$$

What are the fixed points \mathcal{S}_P ?

$$\mathcal{S}_P = \left\{ T \in \mathcal{S} \mid xTx^{-1} = T \quad \forall x \in P \right\}.$$

Let $T \in \mathcal{S}_P$, so $xTx^{-1} = T$ for all $x \in P$. Then $P \leq N_G(T)$, so both P and T are Sylow p -subgroups in $N_G(H)$ as well as G .

Then there exists a $f \in N_G(T)$ such that $T = gPg^{-1}$. But the point is that in the normalizer, there is only **one** Sylow p -subgroup. But then T is the unique largest normal subgroup of $N_G(T)$, which forces $T = P$.

But then $\mathcal{S}_P = \{P\}$, and using the formula, we have $r_p \equiv 1 \pmod{p}$.

Now modify this slightly by letting G act on \mathcal{S} (instead of just P) by conjugation. Since all Sylows are conjugate, by Sylow (1) there is only one orbit, so $\mathcal{S} = GP$ for $P \in \mathcal{S}$. But then

$$r_p = |\mathcal{S}| = |GP| = [G : G_p] \mid |G|.$$

Note that this gives a precise formula for r_p , although the theorem is just an upper bound of sorts, and $G_p = N_G(P)$.

8.1.4 Applications

Of interest historically: classifying finite *simple* groups, where a group G is *simple* if $N \trianglelefteq G$ and $N \neq \{e\}$, then $N = G$.

Example: Let $G = \mathbb{Z}_p$, any subgroup would need to have order dividing p , so G must be simple.

Example: $G = A_n$ for $n \geq 5$ (see Galois theory)

One major application is proving that groups of a certain order are *not* simple.

Applications:

1. Let $|G| = p^n q$ with $p > q$. Then G is not simple.

Strategy: Find a proper normal nontrivial subgroup using Sylow theory. Can either show $r_p = 1$, or produce normal subgroups by intersecting distinct Sylow p -subgroups.

Consider r_p , then $r_p = p^\alpha q^\beta$ for some α, β . But since $r_p \equiv 1 \pmod{p}$, $p \nmid r_p$, we must have $r_p = 1, q$. But since $q < p$ and $q \not\equiv 1 \pmod{p}$, this forces $r_p = 1$.

So let P be a Sylow p -subgroup, then $P < G$. Then gPg^{-1} is also a Sylow, but there's only 1 of them, so P is normal.

2. Let $|G| = 45$, then G is not simple. (Exercise)
3. Let $|G| = p^n$, then G is not simple if $n > 1$.

By Sylow (1), there is a normal subgroup of order p^{n-1} in G .

4. Let $|G| = 48$, then G is not simple.

Note $48 = 2^4 3$, so consider r_2 , the number of Sylow 2-subgroups. Then $r_2 \equiv 1 \pmod{2}$ and $r_2 \mid 48$. So $r_2 = 1, 3$. If $r_2 = 1$, we're done, otherwise suppose $r_2 = 3$.

Let $H \neq K$ be Sylow 2-subgroups, so $|H| = |K| = 2^4 = 16$. Now consider $H \cap K$, which is a subgroup of G . How big is it?

Since $H \neq K$, $|H \cap K| < 16$. The order has to divide 16, so we in fact have $|H \cap K| \leq 8$. Suppose it is less than 4, towards a contradiction. Then

$$|HK| = \frac{|H||K|}{|H \cap K|} \geq \frac{(16)(16)}{4} = 64 > |G| = 48.$$

So we can only have $|H \cap K| = 8$. Since this is an index 2 subgroup in both H and K , it is in fact normal. But then $H, K \subseteq N_G(H \cap K) := X$. But then $|X|$ must be a multiple of 16 *and* divide 48,

so it's either 16 or 28. But $|X| > 16$, because $H \subseteq X$ and $K \subseteq X$. So then $N_G(H \cap K) = G$, and so $H \cap K \trianglelefteq G$.

8.2 Classification of groups of a certain order

We have a classification of finite abelian groups (see table)

9 Lecture 6

Recall the Sylow theorems:

- p groups exist for *every* p^i dividing $|G|$, and $H(p) \trianglelefteq H(p^2) \trianglelefteq \dots H(p^n)$.
- All Sylow p -subgroups are conjugate.
- Numerical constraints
 - $r_p \equiv 1 \pmod{p}$,
 - $r_p \mid |G|$ and $r_p \mid m$,

9.1 Internal Direct Products

Suppose $H, K \leq G$, and consider the smallest subgroup containing both H and K . Denote this $H \vee K$.

If either H or K is normal in G , then we have $H \vee K = HK$. There's a "recipe" for proving you have a direct product of groups:

Lemma: Let G be a group, $H \trianglelefteq G$ and $K \trianglelefteq G$, and

1. $H \vee K = HK = G$,
2. $H \cap K = \{e\}$.

Then $G \cong H \times K$.

Proof:

We first want to show that $hk = kh \ \forall k \in K, h \in H$. We then have

$$hkh^{-1}k^{-1} = (hkh^{-1})k^{-1} \in K = h(kh^{-1}k^{-1}) \in H \implies hkh^{-1}k^{-1} \in H \cap K = \{e\}.$$

So define

$$\begin{aligned} \phi : H \times K &\rightarrow G \\ (h, k) &\mapsto hk, \end{aligned}$$

and (exercise) check that this is a homomorphism, it is surjective, and injective.

Applications:

Theorem: Every group of order p^2 is abelian.

Proof: If G is cyclic, then it is abelian and $G \cong \mathbb{Z}_{p^2}$. So suppose otherwise. By Cauchy, there is an element of order p in G . So let $H = \langle a \rangle$, for which we have $|H| = p$.

Then $H \trianglelefteq G$ by Sylow 1, since it's normal in $H(p^2)$, which would have to equal G .

Now consider $b \notin H$. By Lagrange, we must have $o(b) = 1, p$, and since $e \in H$, we must have $o(b) = p$ (uses fact that G is not cyclic). Now let $K = \langle b \rangle$. Then $|K| = p$, and $K \trianglelefteq G$ by the same argument.

Theorem: Let $|G| = pq$ where $q \not\equiv 1 \pmod p$ and $p < q$. Then G is cyclic (and thus abelian).

Proof: Use Sylow 1. Let P be a sylow p -subgroup. We want to show that $P \trianglelefteq G$ to apply our direct product lemma, so it suffices to show $r_p = 1$.

We know $r_p \equiv 1 \pmod p$ and $r_p \mid |G| = pq$, and so $r_p = 1, q$. It can't be q because $p < q$.

Now let Q be a sylow q -subgroup. Then $r_q \equiv 1 \pmod q$ and $r_q \mid pq$, so $r_q = 1, p$. But since $p < q$, we must have $r_q = 1$. So $Q \trianglelefteq G$ as well.

We now have $P \cap Q = \emptyset$ (why?) and

$$|PQ| = \frac{|P||Q|}{|P \cap Q|} = |P||Q| = pq,$$

and so $G = PQ$, and $G \cong \mathbb{Z}_p \times \mathbb{Z}_q \cong \mathbb{Z}_{pq}$.

Example: every group of order $15 = 5^1 3^1$ is cyclic.

9.2 Determination of groups of a given order

| Order of G | Number of Groups | List of Distinct Groups |
|------------|------------------|--------------------------------------------------------------------------|
| 1 | 1 | $\{e\}$ |
| 2 | 1 | \mathbb{Z}_2 |
| 3 | 1 | \mathbb{Z}_3 |
| 4 | 2 | $\mathbb{Z}_4, \mathbb{Z}_2^2$ |
| 5 | 1 | \mathbb{Z}_5 |
| 6 | 2 | \mathbb{Z}_6, S_3 (*) |
| 7 | 1 | \mathbb{Z}_7 |
| 8 | 5 | $\mathbb{Z}_8, \mathbb{Z}_4 \times \mathbb{Z}_2, \mathbb{Z}_2^3, D_8, Q$ |
| 9 | 2 | $\mathbb{Z}_9, \mathbb{Z}_3^2$ |
| 10 | 2 | \mathbb{Z}_{10}, D_5 |
| 11 | 1 | \mathbb{Z}_{11} |

We still need to justify 6, 8, and 10.

9.3 Free Groups

Define an *alphabet* $A = \{a_1, a_2, \dots, a_n\}$, and let a *syllable* be of the form a_i^m for some m . A *word* is any expression of the form $\prod_{n_i} a_{n_i}^{m_i}$.

We have two operations,

- Concatenation, i.e. $(a_1a_2) \star (a_3^2a_5) = a_1a_2a_3^2a_5$.
- Contraction, i.e. $(a_1a_2^2) \star (a_2^{-1}a_5) = a_1a_2^2a_2^{-1}a_5 = a_1a_2a_5$.

If we've contracted a word as much as possible, we say it is *reduced*.

We let $F[A]$ be the set of reduced words and define a binary operation

$$f : F[A] \times F[A] \rightarrow F[A]$$

$$(w_1, w_2) \mapsto w_1w_2 \text{ (reduced) .}$$

Theorem: (A, f) is a group.

Definition: $F[A]$ is called the *free group generated by A*. A group G is called *free* on a subset $A \subseteq G$ iff $G \cong F[A]$.

Examples:

1. $A = \{x\} \implies F[A] = \{x^n \mid n \in \mathbb{Z}\} \cong \mathbb{Z}$.
2. $A = \{xy\} = \mathbb{Z} * \mathbb{Z}$ (not defined yet!). Note that there are no relations, i.e. $xyxyxy$ is reduced. To abelianize, we'd need to introduce a relation $xy = yx$.

Properties:

1. If G is free on A and free on B then we must have $|A| = |B|$.
2. Any (nontrivial) subgroup of a free group is free. (See Fraleigh or Hungerford for possible Algebraic proofs!)

Theorem: Let G be generated by some (possibly infinite) subset $A = \{A_i \mid i \in I\}$ and G' be generated by some $A'_i \subseteq A_i$. Then

- (a) There is at most one homomorphism $a_i \rightarrow a'_i$.
- (b) If $G \cong F[A]$, there is exactly *one* homomorphism.

Corollary: Every group G' is a homomorphic image of a free group.

Proof:

Let A be the generators of G' and $G = F[A]$, then define $\varphi(a_i) = a_i$. This is onto exactly because $G' = \langle a_i \rangle$, and using the theorem above we're done.

9.4 Generators and Relations

Let G be a group and $A \subseteq G$ be a generating subset so $G = \langle A \mid a \in A \rangle$. There exists a $\phi : F[A] \rightarrow G$, and by the first isomorphism theorem, we have $F[A]/\ker \phi \cong G$.

Let $R = \ker \phi$, these provide the *relations*.

Examples:

Let $G = \mathbb{Z}_3 = \langle [1]_3 \rangle$. Let $x = [1]_3$, then define $\phi : F[\{x\}] \rightarrow \mathbb{Z}_3$, then since $[1] + [1] + [1] = [0] \pmod{3}$, we have $\ker \phi = \langle x^3 \rangle$.

Let $G = \mathbb{Z} \oplus \mathbb{Z}$, then $G \cong \langle x, y \mid [x, y] = 1 \rangle$.

We'll use this for groups of order 6 – there will be only one presentation that is nonabelian, and we'll exhibit such a group.

10 Lecture 7 (Thursday 29th)

Recall the table of distinct small groups we had:

| Order of G | Number of Groups | List of Distinct Groups |
|------------|------------------|--------------------------------------------------------------------------|
| 1 | 1 | $\{e\}$ |
| 2 | 1 | \mathbb{Z}_2 |
| 3 | 1 | \mathbb{Z}_3 |
| 4 | 2 | $\mathbb{Z}_4, \mathbb{Z}_2^2$ |
| 5 | 1 | \mathbb{Z}_5 |
| 6 | 2 | \mathbb{Z}_6, S_3 (*) |
| 7 | 1 | \mathbb{Z}_7 |
| 8 | 5 | $\mathbb{Z}_8, \mathbb{Z}_4 \times \mathbb{Z}_2, \mathbb{Z}_2^3, D_4, Q$ |
| 9 | 2 | $\mathbb{Z}_9, \mathbb{Z}_3^2$ |
| 10 | 2 | \mathbb{Z}_{10}, D_5 |
| 11 | 1 | \mathbb{Z}_{11} |

Exercise: show that groups of order p^2 are abelian.

We still need to justify S_3, D_4, Q, D_5 .

Recall that for any group A , we can consider the free group on the elements of A , $F[A]$. (Note that we can also restrict A to just its generators.) There is then a homomorphism $F[A] \rightarrow A$, where the kernel is the relations.

Example: $\mathbb{Z} * \mathbb{Z} = \langle x, y \mid xyx^{-1}y^{-1} = e \rangle$ where $x = (1, 0), y = (0, 1)$.

10.1 Groups of Order 6

Let G be nonabelian of order 6. Idea: look at subgroups of index 2.

Let P be a Sylow 3-subgroup of G , then $r_3 = 1$ so $P \trianglelefteq G$. Moreover, P is cyclic since it is order 3, so $P = \langle a \rangle$. But since $|G/P| = 2$, it is also cyclic, so $G/P = \langle bP \rangle$.

Note that $b \notin P$, but $b^2 \in P$ since $(bP)^2 = P$, so $b^2 \in \{e, a, a^2\}$. If $b = a, a^2$ then b has order 6, but this would make $G = \langle b \rangle$ cyclic and thus abelian. So $b^2 = 1$.

Since $P \trianglelefteq G$, we have $bPb^{-1} = P$, and in particular bab^{-1} has order 3. So either $bab^{-1} = a$, or $bab^{-1} = a^2$. If $bab^{-1} = a$, then G is abelian, so $bab^{-1} = a^2$.

So $G = \langle a, b \mid a^3 = e, b^2 = e, bab^{-1} = a^2 \rangle$.

We've shown that *if* there is such a nonabelian group, then it must satisfy these relations – we still need to produce some group that actually realizes this.

Consider the symmetries of the triangle:



Figure 1: Image

You can check that a, b satisfy the appropriate relations.

For order 10, a similar argument yields

$G = \langle a, b \mid a^5 = 1, b^2 = 1, ba = a^4b \rangle$, and this is realized by symmetries of the pentagon where $a = (1\ 2\ 3\ 4\ 5), b = (1\ 4)(2\ 3)$.

10.2 Groups of Order 8

Assume G is nonabelian of order 8. G has no elements of order 8, so the only possibilities are 1, 2, 4.

Assume all elements have order 1, 2. Let $a, b \in G$, consider $(ab)^2 = abab \implies ab = b^{-1}a^{-1} = ba$, so G is abelian. So there must be an element of order 4.

So suppose $a \in G$ has order 4, which is an index 2 subgroup, and so $\langle a \rangle \trianglelefteq G$. But $|G/\langle a \rangle| = 2$ is cyclic, so $G/\langle a \rangle = \langle bH \rangle$.

Note that $b^2 \in H = \langle a \rangle$.

If $b^2 = a, a^3$ then b will have order 8, making G cyclic. So $b^2 = 1, a^2$ (these are both valid options!)

Since $H \trianglelefteq G$, we have $b\langle a \rangle b^{-1} = \langle a \rangle$, and since a has order 4, so does bab^{-1} . So $bab^{-1} = a, a^3$, but a is not an option because this would make G abelian.

So we have two options:

$$\begin{aligned} G_1 &= \langle a, b \mid a^4 = 1, b^2 = 1, bab^{-1} = a^3 \rangle \\ G_2 &= \langle a, b \mid a^4 = 1, b^2 = a^2, bab^{-1} = a^3 \rangle. \end{aligned}$$

Exercise: prove $G_1 \not\cong G_2$.

Now to realize these groups:

- G_1 is the group of symmetries of the square, where $a = (1\ 2\ 3\ 4), b = (1\ 3)$.
- $G_2 \cong Q$, the quaternions, where $Q = \{\pm 1, \pm i, \pm j, \pm k\}$, and there are relations (add picture here).

10.3 Some Nice Facts

- If and $\phi : G \rightarrow G'$, then
 - $N \trianglelefteq G \implies N \trianglelefteq \phi(G)$, although not necessarily in G .
 - $N' \trianglelefteq G' \implies \phi^{-1}(N') \trianglelefteq G$

Definition: A *maximal normal subgroup* is a normal subgroup $M \trianglelefteq G$ that is properly contained in G , and if $M \leq N \trianglelefteq G$ (where N is proper) then $M = N$.

Theorem: M is a maximal normal subgroup of G iff G/M is simple.

10.4 Simple Groups

Definition: A group G is simple iff $N \trianglelefteq G \implies N = \{e\}, G$.

Note that if an abelian group has *any* subgroups, then it is not simple, so $G = \mathbb{Z}_p$ is the only simple abelian group. Another example of a simple group is A_n for $n \geq 5$.

Theorem (Feit-Thompson, 1964): Every finite nonabelian simple group has even order.

Note that this is a consequence of the “odd order theorem”.

10.5 Series of Groups

A composition series is a descending series of pairwise normal subgroups such that each successive quotient is simple:

$$\begin{aligned} G_0 \trianglelefteq G_1 \trianglelefteq G_2 \cdots \trianglelefteq \{e\} \\ G_i/G_{i+1} \text{ simple.} \end{aligned}$$

Example:

$$\begin{aligned} \mathbb{Z}_9 \trianglelefteq \mathbb{Z}_3 \trianglelefteq \{e\} \\ \mathbb{Z}_9/\mathbb{Z}_3 = \mathbb{Z}_3, \\ \mathbb{Z}_3/\{e\} = \mathbb{Z}_3. \end{aligned}$$

Example:

$$\begin{aligned} \mathbb{Z}_6 \trianglelefteq \mathbb{Z}_3 \trianglelefteq \{e\} \\ \mathbb{Z}_6/\mathbb{Z}_3 = \mathbb{Z}_2 \\ \mathbb{Z}_2/\{e\} = \mathbb{Z}_2. \end{aligned}$$

but also

$$\begin{aligned} \mathbb{Z}_6 \trianglelefteq \mathbb{Z}_2 \trianglelefteq \{e\} \\ \mathbb{Z}_6/\mathbb{Z}_2 = \mathbb{Z}_3 \\ \mathbb{Z}_3/\{e\} = \mathbb{Z}_3. \end{aligned}$$

Theorem (Jordan-Holder): Any two composition series are “isomorphic” in the sense that the same quotients appear in both series, up to a permutation.

Definition: A group is *solvable* iff it has a composition series where all factors are abelian.

Exercise: Show that any abelian group is solvable.

Example: S_n is *not* solvable for $n \geq 5$, since

$$\begin{aligned} S_n &\trianglelefteq A_n \trianglelefteq \{e\} \\ S_n/A_n &= \mathbb{Z}_2 \text{ simple} \\ A_n/\{e\} &= A_n \text{ simple} \iff n \geq 5. \end{aligned}$$

Example:

$$\begin{aligned} S_4 &\trianglelefteq A_4 \trianglelefteq G \trianglelefteq \{e\} \quad \text{where } |H| = 4 \\ S_4/A_4 &= \mathbb{Z}_2 \\ A_4/H &= \mathbb{Z}_3 \\ H/\{e\} &= \{a, b\}?. \end{aligned}$$

11 Lecture 8: Series of Groups

Recall that a simple group has no nontrivial normal subgroups.

Example:

$$\begin{aligned} \mathbb{Z}_6 &\trianglelefteq \langle [3] \rangle \trianglelefteq \langle [0] \rangle \\ \mathbb{Z}_6/\langle [3] \rangle &= \mathbb{Z}_3 \\ \langle [3] \rangle/\langle [0] \rangle &= \mathbb{Z}_2. \end{aligned}$$

Definition: A *normal series* (or an *invariant series*) of a group G is a finite sequence $H_i \leq G$ such that $H_i \trianglelefteq H_{i+1}$ and $H_n = G$, so we obtain

$$H_1 \trianglelefteq H_2 \trianglelefteq \cdots \trianglelefteq H_n = G.$$

Definition: A normal series $\{K_i\}$ is a *refinement* of $\{H_i\}$ if $K_i \leq H_i$ for each i .

Definition: We say two normal series of the same group G are isomorphic if there is a bijection from

$$\{H_i/H_{i+1}\} \iff \{K_j/K_{j+1}\}$$

Theorem (Schreier): Two normal series of G has isomorphic refinements.

Definition: A normal series of G is a *composition series* iff all of the successive quotients H_i/H_{i+1} are simple.

Note that every finite group has a composition series, because any group has a maximal normal subgroup.

Theorem (Jordan-Holder): Any two composition series of a group G are isomorphic.

Proof: Apply Schreier's refinement theorem.

Example: Consider $S_n \trianglelefteq A_n \trianglelefteq \{e\}$. This is a composition series, with quotients Z_2, A_n , which are both simple.

Definition: A group G is *solvable* iff it has a composition series in which all of the successive quotients are abelian.

Examples:

- Any abelian group is solvable.
- S_n is not solvable for $n \geq 5$, since A_n is not abelian for $n \geq 5$.(?)

Recall Feit-Thompson: Any nonabelian simple group is of *even* order.

Consequence: Every group of *odd* order is solvable.

11.1 The Commutator Subgroup

Let G be a group, and let $[G, G] \leq G$ be the subgroup of G generated by elements $aba^{-1}b^{-1}$, i.e. every element is a *product* of commutators. So $[G, G]$ is called *the commutator subgroup*.

Theorem: Let G be a group, then $[G, G] \trianglelefteq G$ and $G/[G, G]$ is abelian. Also, $[G, G]$ is the smallest normal subgroup such that the quotient is abelian, i.e. if $H \trianglelefteq G$ and G/H is abelian then $[G, G] \leq H$.

Proof:

1. $[G, G]$ is a subgroup.
 - Closure is clear from definition as generators.
 - The identity is $e = ee^{-1}ee^{-1}$.
 - So it suffices to show that $(aba^{-1}b^{-1})^{-1} \in [G, G]$, but this is given by $bab^{-1}a^{-1}$ which is of the correct form.
2. $[G, G]$ is normal

Let $x_i \in [G, G]$, then we want to show $g \prod x_i g^{-1} \in [G, G]$, but this reduces to just showing $gxg^{-1} \in [G, G]$ for a single $x \in [G, G]$. Then,

$$\begin{aligned} g(aba^{-1}b^{-1})g^{-1} &= (g^{-1}aba^{-1})e(b^{-1}g) \\ &= (g^{-1}aba^{-1})(gb^{-1}bg^{-1})(b^{-1}g) \\ &= [(g^{-1}a)b(g^{-1}a)^{-1}b^{-1}][bg^{-1}b^{-1}g] \\ &\in [G, G]. \end{aligned}$$

3. The quotient is abelian

Let $H = [G, G]$. We have $aHbH = (ab)H$ and $bHaH = (ba)H$. But $abH = baH$ because $(ba)^{-1}(ab) = a^{-1}b^{-1}ab \in [G, G]$.

4. Suppose G/N is abelian. Let $aba^{-1}b^{-1} \in [G, G]$. Then $abN = baN$, so $aba^{-1}b^{-1} \in N$ and thus $[G, G] \subseteq N$.

11.2 Free Abelian Groups

Example: $\mathbb{Z} \times \mathbb{Z}$.

Take $e_1 = (1, 0), e_2 = (0, 1)$. Then $(x, y) \in \mathbb{Z}^2$ can be written $x(1, 0) + y(0, 1)$, so $\{e_i\}$ behaves like a basis for a vector space.

Definition: A group G is *free abelian* if there is a subset $X \subseteq G$ such that every $g \in G$ can be represented as

$$g = \sum_{i=1}^r n_i x_i, \quad x_i \in X, \quad n_i \in \mathbb{Z}.$$

Equivalently, X generates G , so $G = \langle X \rangle$, and if $\sum n_i x_i = 0 \implies n_i = 0 \forall i$.

If this is the case, we say X is a *basis* for G .

Examples:

- \mathbb{Z}^n is free abelian
- \mathbb{Z}_n is not free abelian, since $n[1] = 0$ and $n \neq 0$. In general, you can replace \mathbb{Z}_n by any finite group and replace n with the order of the group.

Theorem: If G is free abelian on X where $|X| = r$, then $G \cong \mathbb{Z}^r$.

Theorem: If $X = \{x_i\}_{i=1}^r$, then a basis for \mathbb{Z}^r is given by

$$\{(1, 0, 0, \dots), (0, 1, 0, \dots), \dots, (0, \dots, 0, 1)\} := \{e_1, e_2, \dots, e_r\}$$

Proof: Use the map $\phi : G \rightarrow \mathbb{Z}^r$ where $x_i \mapsto e_i$, and check that this is an isomorphism of groups.

Theorem: Let G be free abelian with two bases X, X' , then $|X| = |X'|$.

Definition: Let G be free abelian, then if X is a basis then $|X|$ is called the *rank* of G .

12 Another Lecture: On to Rings

Recall the definition of a ring: A ring $(R, +, \times)$ is a set with binary operations such that

1. $(R, +)$ is a group,
2. (R, \times) is a monoid.

Examples: $R = \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$, or the ring of $n \times n$ matrices, or \mathbb{Z}_n .

A ring is *commutative* iff $ab = ba$ for every $a, b \in R$, and a *ring with unity* is a ring such that $\exists 1 \in R$ such that $a1 = 1a = a$. Exercise: show that 1 is unique if it exists.

In a ring with unity, an element $a \in R$ is a *unit* iff $\exists b \in R$ such that $ab = ba = 1$.

A ring with unity is a *division ring* iff every nonzero element is a unit. A division ring is said to be a *field* iff it is commutative.

Suppose that $a, b \neq 0$ with $ab = 0$. Then a, b are said to be *zero divisors*. A commutative ring without zero divisors is an *integral domain*.

In \mathbb{Z}_n , an element a is a zero divisor iff $\gcd(a, n) \neq 1$.

Fact: In a ring with no zero divisors, we have $ab = ac, a \neq 0 \implies b = c$.