

Title

D. Zack Garza

Friday 20th March, 2020

Contents

1 Friday February 21st

1

1 Friday February 21st

Question: how do we define $h_{V,D}$?

Answer: write $D = D_1 - D_2$ which are (very) ample divisors and basepoint free. We then obtain embeddings

$$\begin{aligned}\varphi_1 : V &\hookrightarrow \mathbb{P}_K^{n_1} \\ \varphi_2 : V &\hookrightarrow \mathbb{P}_K^{n_2}.\end{aligned}$$

So write

$$h_{V,D}(p) = h(\varphi_1(p)) - h(\varphi_2(p)) + O(1)$$

Example 1.1.

For E/K an elliptic curve,

- $2[0]$ is an ample divisor
- $3[0]$ is a very ample divisor.

Let K be a local field (i.e. \mathbb{C}, \mathbb{R} , a p -adic field, or $\mathbb{F}_q((t))$ formal Laurent series) and A/K be an abelian variety; we want to understand $A(K)$. We know this has the structure of compact abelian K -analytic Lie group.

- Question 1: What does Lie theory say?
- Question 2: What extra information comes from A/K being a g -dimensional abelian variety?

If $K = \mathbb{C}$, then $A(K) \cong (\mathbb{R}/\mathbb{Z})^{2g}$. If $K = \mathbb{R}$, then $A(K) \cong (\mathbb{R}/\mathbb{Z})^g \oplus \prod_{i=1}^d \mathbb{Z}/2\mathbb{Z}$ where $0 \leq d \leq g$.

Fix d , then

- Let E_1/\mathbb{R} with $\Delta > 0$ (and thus 3 real roots), then $E_1(\mathbb{R})[2] = (\mathbb{Z}/2\mathbb{Z})^2$.
- Let E_2/\mathbb{R} with $\Delta < 0$ (and 1 real root), then $E_2(\mathbb{R})[2] = \mathbb{Z}/2\mathbb{Z}$.

By taking products of E_1 and E_2 , i.e. $A = (E_1)^d \times (E_2)^{g-d}$.

Todo: find reference in Silverman?

Fact $A(K)$ is totally disconnected and homeomorphic to a Cantor set.

Fact (From Lie Theory, Serre p.116) If G is an abelian compact K -analytic Lie group, then there exists a filtration by open finite index subgroups

$$G = G^0 \supset G^1 \supset \dots \supset G^n \supset \dots$$

such that

1. The successive quotients are finite, and each G^i is *standard*, i.e. obtained by evaluating a formal group law on $(\mathfrak{m}^i)^g$.
2. $\bigcap_i G^i = (0)$.
3. G^i/G^{i+1} has exponent p , i.e. it is a finite dimensional $\mathbb{Z}/p\mathbb{Z}$ -vector space.
4. $G'[\text{tors}] = G'[p^\infty]$, all of the prime-to- p torsion is p -primary.

Todo: define p -primary torsion, \mathbb{Q}_p .

What structure theorem does this give?

Theorem 1.1 (C-Lacy).

Let G be a compact, second countable, K -analytic abelian Lie group of dimension $g \geq 1$. Then

- a. If $\text{char } K = 0$ and $d = [K : \mathbb{Q}_p]$, then

$$G \cong_{\text{TopGrp}} \mathbb{Z}_p^{dg} \oplus G[\text{tors}]$$

where $G[\text{tors}]$ is finite.

- b. If $\text{char } K = p$, i.e. $K = \mathbb{F}_q((t))$, and if it is true that $G[\text{tors}]$ is finite iff $G[p]$ finite, then

$$G \cong_{\text{TopGrp}} \prod_{i=1}^{\infty} \mathbb{Z}_p \oplus G[\text{tors}]$$

For any $g \geq 1$, $(T, +)$ a finite discrete abelian group $(R, +) \cong (\mathbb{Z}_p^d, +)$ and $R^g \oplus T$ is a compact abelian K -analytic Lie group isomorphic to $\mathbb{Z}_p^{dg} \oplus T$ (?).

Question: what does this mean for $G = S^1$?