Problem Set 5

D. Zack Garza

October 21, 2019

Contents

1	Problem 1	1
	1.1 Problem 2	2

1 Problem 1

We first make the following definitions:

$$S := \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} a_{jk} = \sup \left\{ \sum_{(j,k) \in B} a_{jk} \ni B \subset \mathbb{N}^2, \ |B| < \infty \right\}$$
$$T := \sum_{k=1}^{\infty} \sum_{j=1}^{\infty} a_{kj} = \sup \left\{ \sum_{(j,k) \in C} a_{kj} \ni C \subset \mathbb{N}^2, \ |B| < \infty \right\}.$$

We will show that S = T by showing that $S \leq T$ and $T \leq S$.

Let $B \subset \mathbb{N}^2$ be finite, so $B \subseteq [0, I] \times [0, J] \subset \mathbb{N}^2$.

Now letting $R > \max(I, J)$, we can define $C = [0, R]^2$, which satisfies $B \subseteq C \subset \mathbb{N}^2$ and $|C| < \infty$.

Moreover, since $a_{jk} \geq 0$ for all pairs (j, k), we have the following inequality:

$$\sum_{(j,k)\in B} a_{jk} < \sum_{(k,j)\in C} a_{jk} \le \sum_{(k,j)\in C} a_{jk} \le T,$$

since T is a supremum over all such sets C, and the terms of any finite sum can be rearranged.

But since this holds for every B, we this inequality also holds for the supremum of the smaller term by order-limit laws, and so

$$S := \sup_{B} \sum_{(k,j) \in B} a_{jk} \le T.$$

An identical argument shows that $T \leq S$, yielding the desired equality. \square

1.1 Problem 2

We want to show the following equality:

$$\int_0^1 g(x) \ dx = \int_0^1 f(x) \ dx.$$

To that end, we can rewrite this using the integral definition of g(x):

$$\int_0^1 \int_x^1 \frac{f(t)}{t} dt dx = \int_0^1 f(x) dx$$

Note that if we can switch the order of integration, we would have

$$\int_{0}^{1} \int_{x}^{1} \frac{f(t)}{t} dt dx = \int_{0}^{1} \int_{0}^{t} \frac{f(t)}{t} dx dt$$

$$= \int_{0}^{1} \frac{f(t)}{t} \int_{0}^{t} dx dt$$

$$= \int_{0}^{1} \frac{f(t)}{t} (t - 0) dt$$

$$= \int_{0}^{1} f(t) dt,$$

which is what we wanted to show, and so we are simply left with the task of showing that this is switch of integrals is justified.