

Assignment 6: The Fourier Transform

D. Zack Garza

October 31, 2019

Contents

1	Problem 1	1
2	Problem 2	2
2.1	Part (a)	2
3	Problem 3	2
4	Problem 4	2
5	Problem 5	2
6	Problem 6	2

1 Problem 1

Assuming the hint, we have

$$\lim_{|\xi| \rightarrow \infty} \hat{f}(\xi) = \lim_{\xi' \rightarrow 0} \frac{1}{2} \int_{\mathbb{R}^n} (f(x) - f(x - \xi')) \exp(-2\pi i x \cdot \xi) \, dx$$

But as an immediate consequence, this yields

$$\begin{aligned} |\hat{f}(\xi)| &= \left| \int_{\mathbb{R}^n} (f(x) - f(x - \xi')) \exp(-2\pi i x \cdot \xi) \, dx \right| \\ &\leq \int_{\mathbb{R}^n} |f(x) - f(x - \xi')| |\exp(-2\pi i x \cdot \xi)| \, dx \\ &\leq \int_{\mathbb{R}^n} |f(x) - f(x - \xi')| \, dx \\ &\rightarrow 0, \end{aligned}$$

which follows from continuity in L^1 since $f(x - \xi') \rightarrow f(x)$ as $\xi' \rightarrow 0$.

It thus only remains to show that the hint holds, and that $\xi' \rightarrow 0$ as $\xi \rightarrow \infty$.

2 Problem 2

2.1 Part (a)

Assuming an interchange of integrals is justified, we have

$$\begin{aligned}\widehat{(f * g)}(\xi) &= \int \int f(x - y)g(y) \exp(-2\pi x \cdot \xi) \, dy \, dx && \stackrel{=?}{=} \int \int f(x - y)g(y) \exp(-2\pi x \cdot \xi) \, dx \, dy \\ &= \int \int f(t) \exp(-2\pi i(x - y) \cdot \xi)g(y) \exp(-2\pi iy \cdot \xi) \, dx \, dy \\ &\quad (t = x - y, \, dt = \, dx) \\ &= \int \int f(t) \exp(-2\pi it \cdot \xi)g(y) \exp(-2\pi iy \cdot \xi) \, dt \, dy \\ &= \int f(t) \exp(-2\pi it \cdot \xi) \left(\int g(y) \exp(-2\pi iy \cdot \xi) \, dy \right) \, dt \\ &= \int f(t) \exp(-2\pi it \cdot \xi) \hat{g}(\xi) \, dt \\ &= \hat{g}(\xi) \int f(t) \exp(-2\pi it \cdot \xi) \, dt \\ &= \hat{g}(\xi) \hat{f}(\xi).\end{aligned}$$

3 Problem 3

4 Problem 4

5 Problem 5

6 Problem 6