# **Title**

## D. Zack Garza

Thursday 3<sup>rd</sup> September, 2020

## **Contents**

1 Wednesday, September 02

1

# 1 Wednesday, September 02

Recommended exercises:

- 0.9
- 0.5 (easy)
- 0.10

Taken:

- 0.11
- 0.3
- 0.4

### Exercise 1.1 (0.9).

Let k be a field and  $d \ge 2$  with  $4 \nmid d$  and  $p \in k[x]$  a polynomial of positive degree.

Factor p in  $\bar{k}[x]$  as  $\prod_{i=1}^{r} (x-a_i)^{e_i}$ , and suppose there is some i such that  $d \nmid e_i$ . Show that

$$f(x,y) \coloneqq y^d - p(x) \in k[x,y]$$

is geometrically irreducible.

Conclude that

$$ff(k[x,y]/\langle f\rangle).$$

is a regular one-variable function field over k.

#### **Solution:**

Recall:

- For L/K
- A polynomial  $f \in k[t_i]$  is geometrically irreducible iff  $f \in \bar{k}[t_i]$  is irreducible as a

polynomial, i.e. if  $f = pq \implies p = 1$  or q = 1.

- A field extension L/k is regular iff any of the following conditions hold:
  - $-\kappa(L)=k$  and L/k is separable, where  $\kappa(L)$  is the field of elements of L algebraic over k

  - $-L \otimes_k \overline{k}$  is a domain or a field. For all L'/k,  $L \otimes_k L'$  is a domain.