# **Complex Analysis Qualifying Exam Review**

D. Zack Garza

# **Table of Contents**

# **Contents**

Table of Contents 2							
1	Use	ful Techniques	4				
	1.1	Notation	4				
	1.2	Greatest Hits	4				
	1.3	Common Tricks	5				
	1.4	Basic but Useful Facts	5				
2	Calc	culus Preliminaries	7				
	2.1	Definitions	7				
	2.2	Theorems	8				
	2.3	Convergence	8				
	2.4	Series and Sequences	9				
	2.5	Exercises	10				
3	Preliminaries 11						
	3.1	Complex Arithmetic	11				
	3.2	Complex Log	12				
	3.3	Complex Calculus	13				
		3.3.1 Holomorphy and Cauchy-Riemann	13				
		3.3.2 Delbar and the Laplacian	16				
		3.3.3 Harmonic Functions and the Laplacian	16				
		3.3.4 Exercises	17				
	3.4	Power Series	19				
		3.4.1 Exercises: Series	22				
4	Cauchy's Theorem 24						
	4.1	Complex Integrals	24				
	4.2	Applications of Cauchy's Theorem	24				
		4.2.1 Integral Formulas and Estimates	24				
		4.2.2 Liouville	27				
		4.2.3 Continuation Principle	28				
	4.3	Exercises	28				
	4.4	Morera's Theorem	30				
		4.4.1 Symmetric Regions	31				
5	Zero	os and Singularities	31				
6	Residues 34						
	6.1	Basics	34				
	6.2	Estimates	35				

## Contents

	6.3	Residue Formulas	37				
		6.3.1 Exercises	38				
	6.4	Argument Principle	42				
		6.4.1 Exercises	43				
7	Rou		43				
	7.1	Counting Zeros	43				
3	Con	formal Maps	44				
	8.1	Linear Fractional Transformations	44				
	8.2	Schwarz	45				
	8.3	By Type	46				
		8.3.1 Plane to Disc	46				
		8.3.2 Sector to Disc	47				
		8.3.3 Strip to Disc	47				
9	Sch	warz Reflection	reflection 47				
10	Sch	warz Lemma	48				
11	Line	ear Fractional Transformations	48				
12	Mor	ntel's Theorem	48				
13	Uns	orted Theorems	48				
1 /	Ann	pendix: Proofs of the Fundamental Theorem of Algebra	49				
. 4	App	14.0.1 Fundamental Theorem of Algebra: Argument Principle	49				
		14.0.2 Fundamental Theorem of Algebra: Rouche's Theorem	49				
		14.0.3 Fundamental Theorem of Algebra: Liouville's Theorem	50				
		14.0.4 Fundamental Theorem of Algebra: Open Mapping Theorem	50				
15	Арр	pendix	51				
	15.1	Misc Basic Algebra	51				
16	Draf	ft of Problem Book	5/1				

Contents 3

A great deal of content borrowed from the following: https://web.stanford.edu/~chriseur/notes\_pdf/Eur\_ComplexAnalysis\_Notes.pdf

# 1 | Useful Techniques

## 1.1 Notation

Notation	Definition
$\mathbb{D} \coloneqq \left\{ z \mid  z  \le 1 \right\}$	The unit disc
$\mathbb{H} := \left\{ x + iy \mid y > 0 \right\}$	The upper half-plane
$X_{rac{1}{2}}$	A "half version of $X$ ", see examples
$\mathbb{H}_{\frac{1}{2}}$ $\mathbb{D}_{\frac{1}{2}}$	The first quadrant  The portion of the first quadrant inside the unit disc
$S := \left\{ x + iy \mid x \in \mathbb{R}, \ 0 < y < \pi \right\}$	The horizontal strip

## 1.2 Greatest Hits



Things to know well:

- Estimates for derivatives, mean value theorem
- ??CauchyTheorem|Cauchy's Theorem
- ??CauchyIntegral|Cauchy's Integral Formula
- ??CauchyInequality]Cauchy's Inequality
- ??Morera]Morera's Theorem
- ??Liouville]Liouville's Theorem
- ??MaximumModulus|Maximum Modulus Principle
- ??Rouche]Rouché's Theorem

1.2 Greatest Hits

- ??SchwarzReflection|The Schwarz Reflection Principle
- ??SchwarzLemma]The Schwarz Lemma
- ??Casorati|Casorati-Weierstrass Theorem
- Properties of linear fractional transformations
- Automorphisms of  $\mathbb{D}, \mathbb{C}, \mathbb{CP}^1$ .

## 1.3 Common Tricks

• Virtually any time: consider 1/f(z) and f(1/z).

Remark 1.3.1 (Showing a function is constant): If you want to show that a function f is constant, try one of the following:

- Write f = u + iv and use Cauchy-Riemann to show  $u_x, u_y = 0$ , etc.
- Show that f is entire and bounded.
  - If you additionally want to show f is zero, show  $\lim_{z\to\infty} f(z) = 0$ .

## 1.4 Basic but Useful Facts

Fact 1.4.1 (Some useful facts about basic complex algebra)

 $\bullet \quad z\bar{z} = |z|^2$ 

$$\Re(z) = \frac{z + \bar{z}}{2}$$

$$\Im(z) = \frac{z - \bar{z}}{2i}.$$

- $\operatorname{Arg}(z/w) = \operatorname{Arg}(z) \operatorname{Arg}(w)$ .
- Exponential forms of cosine and sine, where it's sometimes useful to set  $w \coloneqq e^{iz}$ :

$$\cos(z) = \frac{1}{2} \left( e^{iz} + e^{-iz} \right) = \frac{1}{2} (w + w^{-1})$$

$$\sin(z) = \frac{1}{2i} \left( e^{iz} - e^{-iz} \right) = \frac{1}{2i} (w - w^{-1})$$

$$\cosh(z) = \cos(iz) = \frac{1}{2} \left( e^z + e^{-z} \right)$$

$$\sinh(z) = \sin(iz) = \frac{1}{2} (e^z - e^{-z}).$$

- Setting  $w = e^z$  is useful:
- Various differentials:

$$dz = dx + i \, dy$$
$$d\bar{z} = dx - i \, dy$$

$$f_z = f_x = f_y/i$$
.

• Integral of a complex exponential:

$$\int_0^{2\pi} e^{i\ell x} dx = \begin{cases} 2\pi & \ell = 0\\ 0 & \text{else} \end{cases}.$$

Fact 1.4.2 (Generalized Binomial Theorem)

Define  $(n)_k$  to be the falling factorial

$$\prod_{j=0}^{k-1} (n-k) = n(n-1)\cdots(n-k+1)$$

and set  $\binom{n}{k} := (n)_k/k!$ , then

$$(x+y)^n = \sum_{k>0} \binom{n}{k} x^k y^{n-k}.$$

Fact 1.4.3 (Some useful series)

$$\sum_{k=1}^{n} k = \frac{n(n+1)}{2}$$

$$\sum_{k=1}^{n} k^{2} = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{k=1}^{n} k^{3} = \frac{n^{2}(n+1)^{2}}{4}$$

$$\log(1-x) = \sum_{n\geq 0} \frac{x^{n}}{n}$$

$$x \in ($$

$$\frac{\partial}{\partial z} \sum_{j=0}^{\infty} a_{j}z^{j} = \sum_{j=0}^{\infty} a_{j+1}z^{j}$$

$$\sqrt{1+x} = (1+x)^{1/2} = 1 + (1/2)x + \frac{(1/2)(-1/2)}{2!}x^{2} + \frac{(1/2)(-1/2)(-3/2)}{3!}x^{3} + \cdots$$

$$= 1 + \frac{1}{2}x - \frac{1}{8}x^{2} + \frac{1}{16}x^{3} - \cdots$$

1.4 Basic but Useful Facts

#### Fact 1.4.4

Useful trick for expanding square roots:

$$\sqrt{z} = \sqrt{z_0 + z - z_0} = \sqrt{z_0 \left(1 + \frac{z - z_0}{z}\right)} = \sqrt{z_0} \sqrt{1 + u}, \quad u := \frac{z - z_0}{z}$$

$$\implies \sqrt{z} = \sqrt{z_0} \sum_{k \ge 0} \binom{1/2}{k} \left(\frac{z - z_0}{z}\right)^k.$$

# 2 | Calculus Preliminaries

#### 2.1 Definitions

**Definition 2.1.1** (Locally uniform convergence)

A sequence of functions  $f_n$  is said to converge **locally uniformly** on  $\Omega \subseteq \mathbb{C}$  iff  $f_n \to f$  uniformly on every compact subset  $K \subseteq \Omega$ .

**Definition 2.1.2** (Equicontinuous Family)

A family of functions  $f_n$  is **equicontinuous** iff for every  $\varepsilon$  there exists a  $\delta = \delta(\varepsilon)$  (not depending on n or  $f_n$ ) such that  $|x - y| < \varepsilon \implies |f_n(x) - f_n(y)| < \varepsilon$  for all n.

Remark 2.1.3: Recall Arzelà-Ascoli, an analog of Heine-Borel: for X compact Hausdorff, consider the Banach space  $C(X;\mathbb{R})$  equipped with the uniform norm  $\|f\|_{\infty,X} := \sup_{x \in X} |f(x)|$ . Then a subset  $A \subseteq X$  is compact iff A is closed, uniformly bounded, and equicontinuous. As a consequence, if A is a sequence, it contains a subsequence converging uniformly to a continuous function. The proof is an  $\varepsilon/3$  argument.

**Definition 2.1.4** (Normal Family)

Remark 2.1.5: A continuous function on a compact set is uniformly continuous.

**Definition 2.1.6** (Univalent functions)

A function  $f \in \text{Hol}(U; \mathbb{C})$  is called **univalent** if f is injective.

**Remark 2.1.7:** If  $f: \Omega \to \Omega'$  is a univalent surjection, f is invertible on  $\Omega$  and  $f^{-1}$  is holomorphic. Compare to real functions:  $f(x) = x^3$  is injective on (-c, c) for any c but f'(0) = 0 and  $f^{-1}(x) := x^{1/3}$  is not differentiable at zero.

#### 2.2 Theorems



#### Theorem 2.2.1 (Implicit Function Theorem).

#### Theorem 2.2.2(Inverse Function Theorem).

For  $f \in C^1(\mathbb{R}; \mathbb{R})$  with  $f'(a) \neq 0$ , then f is invertible in a neighborhood  $U \ni a, g := f^{-1} \in C^1(U; \mathbb{R})$ , and at b := f(a) the derivative of g is given by

$$g'(b) = \frac{1}{f'(a)}.$$

For  $F \in C^1(\mathbb{R}^n, \mathbb{R}^n)$  with  $D_f$  invertible in a neighborhood of a, so  $\det(J_f) \neq 0$ , then setting b := F(a),

$$J_{F^{-1}}(q) = (J_F(p))^{-1}$$
.

The version for holomorphic functions: if  $f \in \operatorname{Hol}(\mathbb{C};\mathbb{C})$  with  $f'(p) \neq 0$  then there is a neighborhood  $V \ni p$  with that  $f \in \operatorname{BiHol}(V, f(V))$ .

#### Theorem 2.2.3 (Green's Theorem).

If  $\Omega \subseteq \mathbb{C}$  is bounded with  $\partial \Omega$  piecewise smooth and  $f, g \in C^1(\overline{\Omega})$ , then

$$\int_{\partial\Omega} f \, dx + g \, dy = \iint_{\Omega} \left( \frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \right) \, dA.$$

## 2.3 Convergence

Remark 2.3.1: Recall that absolutely convergent implies convergent, but not conversely:  $\sum k^{-1} = \infty$  but  $\sum (-1)^k k^{-1} < \infty$ . This converges because the even (odd) partial sums are monotone increasing/decreasing respectively and in (0,1), so they converge to a finite number. Their difference converges to 0, and their common limit is the limit of the sum.

## Proposition 2.3.2 (Uniform Convergence of Series).

A series of functions  $\sum_{n=1}^{\infty} f_n(x)$  converges uniformly iff

$$\lim_{n \to \infty} \left\| \sum_{k \ge n} f_k \right\|_{\infty} = 0.$$

## Theorem 2.3.3 (Weierstrass M-Test).

If  $\{f_n\}$  with  $f_n: \Omega \to \mathbb{C}$  and there exists a sequence  $\{M_n\}$  with  $\|f_n\|_{\infty} \leq M_n$  and  $\sum_{n \in \mathbb{N}} M_n < \infty$ ,

2.2 Theorems

then  $f(x) := \sum_{n \in \mathbb{N}} f_n(x)$  converges absolutely and uniformly on  $\Omega$ . Moreover, if the  $f_n$  are continuous, by the uniform limit theorem, f is again continuous.

## 2.4 Series and Sequences

Remark 2.4.1: Note that if a power series converges uniformly, then summing commutes with integrating or differentiating.

## Proposition 2.4.2 (Ratio Test).

Consider  $\sum c_k z^k$ , set  $R = \lim \left| \frac{c_{k+1}}{c_k} \right|$ , and recall the **ratio test**:

- $R \in (0,1) \implies$  convergence.
- $R \in (1, \infty] \implies$  divergence.
- R = 1 yields no information.

**Theorem 2.2 (Root Test).** Suppose  $\sum a_n(z-z_0)^n$  is a formal power series. Let

$$R = \liminf_{n \to \infty} |a_n|^{-\frac{1}{n}} = \frac{1}{\limsup_{n \to \infty} |a_n|^{\frac{1}{n}}} \in [0, +\infty].$$

Then  $\sum_{n=0}^{\infty} a_n (z-z_0)^n$ 

- (a) converges absolutely in  $\{z : |z z_0| < R\}$ ,
- (b) converges uniformly in  $\{z : |z z_0| \le r\}$  for all r < R, and
- (c) diverges in  $\{z : |z z_0| > R\}$ .

Figure 1: image 2021-05-27-15-40-58

Proposition 2.4.3 (Root Test).

## Proposition 2.4.4 (Radius of Convergence by the Root Test).

For 
$$f(z) = \sum_{k \in \mathbb{N}} c_k z^k$$
, defining

$$\frac{1}{R} \coloneqq \limsup_{k} |a_k|^{\frac{1}{k}},$$

then f converges absolutely and uniformly for  $D_R := |z| < R$  and diverges for |z| > R.

Moreover f is holomorphic in  $D_R$ , can be differentiated term-by-term, and  $f' = \sum_{k \in \mathbb{N}} nc_k z^k$ .

#### Fact 2.4.5

Recall the *p*-test:

$$\sum n^{-p} < \infty \iff p \in (1, \infty).$$

#### Fact 2.4.6

The product of two sequences is given by the Cauchy product

$$\sum a_k z^k \cdot \sum b_k z^k = \sum c_k z^k, \quad c_k := \sum_{j \le k} a_k b_{k-j}.$$

#### Fact 2.4.7

Recall how to carry out polynomial long division:

Polynomial long division

#### Fact 2.4.8 (Partial Fraction Decomposition)

- For every root  $r_i$  of multiplicity 1, include a term  $A/(x-r_i)$ .
- For any factors g(x) of multiplicity k, include terms A<sub>1</sub>/g(x), A<sub>2</sub>/g(x)<sup>2</sup>, ···, A<sub>k</sub>/g(x)<sup>k</sup>.
  For irreducible quadratic factors h<sub>i</sub>(x), include terms of the form Ax + B/h<sub>i</sub>(x).

#### 2.5 Exercises

## Exercise 2.5.1 (?)

Find the radius of convergences for the power series expansion of  $\sqrt{z}$  about  $z_0 = 4 + 3i$ .

# **3** | Preliminaries

#### **Definition 3.0.1** (Toy contour)

A closed Jordan curve that separates  $\mathbb{C}$  into an exterior and interior region is referred to as a **toy contour**.

## 3.1 Complex Arithmetic

~

## Fact 3.1.1 (Complex roots of a number)

The complex nth roots of  $z := re^{i\theta}$  are given by

$$\left\{ \omega_k := r^{1/n} e^{i\left(\frac{\theta + 2k\pi}{n}\right)} \mid 0 \le k \le n - 1 \right\}.$$

Note that one root is  $r^{1/n} \in \mathbb{R}$ , and the rest are separated by angles of  $2\pi/n$ . Mnemonic:

$$z = re^{i\theta} = re^{i(\theta + 2k\pi)} \implies z^{1/n} = \cdots$$

#### Fact 3.1.2

Common trick:

$$f^{1/n} = e^{\frac{1}{n}\log(f)},$$

taking (say) a principal branch of log given by  $\mathbb{C} \setminus (-\infty, 0] \times 0$ .

#### Fact 3.1.3

Some computations that come up frequently:

$$|z \pm w|^2 = |z|^2 + |w|^z + 2\Re(\overline{w}z)$$
$$(a+bi)(c+di) = (ac-bd) + (ad+bc).$$

#### Fact 3.1.4

Some useful facts:

$$|e^z| = e^{\Re(z)}.$$

Preliminaries 11

On  $S^1$ ,

$$1/z = e^{-i\theta}$$

$$\cos(\theta) = \frac{e^{i\theta} + e^{-i\theta}}{2} = \frac{z + 1/z}{2}$$

$$\sin(\theta) = \frac{e^{i\theta} - e^{-i\theta}}{2i} = \frac{z - 1/z}{2i}.$$

## 3.2 Complex Log

Fact 3.2.1 (Complex Log)

For  $z = re^{i\theta} \neq 0$ ,  $\theta$  is of the form  $\Theta + 2k\pi$  where  $\Theta = \operatorname{Arg} z$  We define

$$\log(z) = \ln(|z|) + i\operatorname{Arg}(z)$$

and  $z^c := e^{c \log(z)}$ .

Proposition 3.2.2 (Existence of complex log).

Suppose  $\Omega$  is a simply-connected region such that  $1 \in \Omega, 0 \notin \Omega$ . Then there exists a branch of F(z) := Log(z) such that

- F is holomorphic on  $\Omega$ ,
- $e^{F(z)} = z$  for all  $z \in \Omega$
- $F(x) = \log(x)$  for  $x \in \mathbb{R}$  in a neighborhood of 1.

**Definition 3.2.3** (Principal branch and exponential)

Take  $\mathbb{C}$  and delete  $\mathbb{R}^{\leq 0}$  to obtain the **principal branch** of the logarithm, defined as

$$Log(z) := log(r) + i\theta$$
  $|\theta| < \pi$ .

Similarly define

$$z^{\alpha} := e^{\alpha \operatorname{Log}(z)}$$
.

Theorem 3.2.4 (Existence of log).

If f is holomorphic and nonvanishing on a simply-connected region  $\Omega$ , then there exists a holomorphic G on  $\Omega$  such that

$$f(z) = e^{G(z)}.$$

## 3.3 Complex Calculus

3.2 Complex Log

**Remark 3.3.1:** When parameterizing integrals  $\int_{\gamma} f(z) dz$ , parameterize  $\gamma$  by  $\theta$  and write  $z = re^{i\theta}$  so  $dz = ire^{i\theta} d\theta$ .

## **⚠** Warning 3.3.2

 $f(z) = \sin(z), \cos(z)$  are unbounded on  $\mathbb{C}$ ! An easy way to see this: they are nonconstant and entire, thus unbounded by Liouville.

**Example 3.3.3(?):** You can show  $f(z) = \sqrt{z}$  is not holomorphic by showing its integral over  $S^1$  is nonzero. This is a direct computation:

$$\int_{S^1} z^{1/2} dz = \int_0^{2\pi} (e^{i\theta})^{1/2} i e^{i\theta} d\theta$$

$$= i \int_0^{2\pi} e^{\frac{i3\theta}{2}} d\theta$$

$$= i \left(\frac{2}{3i}\right) e^{\frac{i3\theta}{2}} \Big|_0^{2\pi}$$

$$= \frac{2}{3} \left(e^{3\pi i - 1}\right)$$

$$= -\frac{4}{3}.$$

Note an issue: a different parameterization yields a different (still nonzero) number

$$\dots = \int_{-\pi}^{\pi} (e^{i\theta})^{1/2} i e^{i\theta} d\theta$$
$$= \frac{2}{3} \left( e^{\frac{3\pi i}{2}} - e^{\frac{-3\pi i}{2}} \right)$$
$$= -\frac{4i}{3}.$$

This is these are paths that don't lift to closed loops on the Riemann surface defined by  $z \mapsto z^2$ .

#### 3.3.1 Holomorphy and Cauchy-Riemann

#### **Definition 3.3.4** (Analytic)

A function  $f: \Omega \to \mathbb{C}$  is analytic at  $z_0 \in \Omega$  iff there exists a power series  $g(z) = \sum a_n (z - z_0)^n$  with radius of convergence R > 0 and a neighborhood  $U \ni z_0$  such that f(z) = g(z) on U.

**Definition 3.3.5** (Complex differentiable / holomorphic /entire)

A function  $f: \mathbb{C} \to \mathbb{C}$  is **complex differentiable** or **holomorphic** at  $z_0$  iff the following limit exists:

$$\lim_{h \to 0} \frac{f(z_0 + h) - f(h)}{h}.$$

A function that is holomorphic on  $\mathbb{C}$  is said to be **entire**. Equivalently, there exists an  $\alpha \in \mathbb{C}$  such that

$$f(z_0 + h) - f(z_0) = \alpha h + R(h)$$
  $R(h) \xrightarrow{h \to 0} 0.$ 

In this case,  $\alpha = f'(z_0)$ .

## Example 3.3.6 (Holomorphic vs non-holomorphic):

- $f(z) = \frac{1}{z}$  is holomorphic on  $\mathbb{C} \setminus \{0\}$ .
- $f(z) = \bar{z}$  is not holomorphic, since  $\frac{\bar{h}}{h}$  does not converge (but is real differentiable).

## Definition 3.3.7 (Real (multivariate) differentiable)

A function  $F: \mathbb{R}^n \to \mathbb{R}^m$  is **real-differentiable** at **p** iff there exists a linear transformation A such that

$$\frac{\|F(\mathbf{p} + \mathbf{h}) - F(\mathbf{p}) - A(\mathbf{h})\|}{\|\mathbf{h}\|} \stackrel{\|\mathbf{h}\| \to 0}{\longrightarrow} 0.$$

Rewriting,

$$||F(\mathbf{p} + \mathbf{h}) - F(\mathbf{p}) - A(\mathbf{h})|| = ||\mathbf{h}|| ||R(\mathbf{h})||$$
  $||R(\mathbf{h})|| \stackrel{||\mathbf{h}|| \to 0}{\longrightarrow} 0.$ 

Equivalently,

$$F(\mathbf{p} + \mathbf{h}) - F(\mathbf{p}) = A(\mathbf{h}) + \|\mathbf{h}\| R(\mathbf{h}) \qquad \|R(\mathbf{h})\| \stackrel{\|\mathbf{h}\| \to 0}{\longrightarrow} 0.$$

Or in a slightly more useful form,

$$F(\mathbf{p} + \mathbf{h}) = F(\mathbf{p}) + A(\mathbf{h}) + R(\mathbf{h}) \qquad \qquad R \in o(\|\mathbf{h}\|), \text{ i.e. } \frac{\|R(\mathbf{h})\|}{\|\mathbf{h}\|} \xrightarrow{\mathbf{h} \to 0} 0.$$

#### Proposition 3.3.8 (Complex differentiable implies Cauchy-Riemann).

If f is differentiable at  $z_0$ , then the limit defining  $f'(z_0)$  must exist when approaching from any direction. Identify f(z) = f(x, y) and write  $z_0 = x + iy$ , then first consider  $h \in RR$ , so  $h = h_1 + ih_2$  with  $h_2 = 0$ . Then

$$f'(z_0) = \lim_{h_1 \to 0} \frac{f(x+h_1, y) - f(x, y)}{h_1} := \frac{\partial f}{\partial x}(x, y).$$

Taking  $h \in i\mathbb{R}$  purely imaginary, so  $h = ih_2$ ,

$$f'(z_0) = \lim_{ih_2 \to 0} \frac{f(x, y + h_2) - f(x, y)}{ih_2} := \frac{1}{i} \frac{\partial f}{\partial y}(x, y).$$

Equating,

$$\frac{\partial f}{\partial x} = \frac{1}{i} \frac{\partial f}{\partial y},$$

and writing f = u + iv and 1/i = -i yields

$$\begin{split} \frac{\partial f}{\partial x} &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \\ \frac{1}{i} \frac{\partial f}{\partial y} &= \frac{1}{i} \left( \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y} \,. \end{split}$$

Thus

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \qquad \qquad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

## Proposition 3.3.9 (Polar Cauchy-Riemann equations).

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}$$
 and  $\frac{1}{r} \frac{\partial u}{\partial \theta} = -\frac{\partial v}{\partial r}$ .

Proof .

Setting

$$z = re^{i\theta} = r(\cos(\theta) + i\sin(\theta)) = x + iy$$

yields  $x = r\cos(\theta), y = r\sin(\theta)$ , one can identify

$$x_r = \cos(\theta), x_\theta = -r\sin(\theta)$$
  
 $y_r = \sin(\theta), y_\theta = r\cos(\theta).$ 

Now apply the chain rule:

$$u_r = u_x x_r + u_y y_r$$

$$= v_y x_r - v_x y_r$$

$$= v_y \cos(\theta) - v_x \sin(\theta)$$

$$= \frac{1}{r} (v_y r \cos(\theta) - v_x r \sin(\theta))$$

$$= \frac{1}{r} (v_y y_\theta + v_x x_\theta)$$

$$= \frac{1}{r} v_\theta.$$
CR

Similarly,

$$v_r = v_x x_r + v_y y_r$$

$$= v_x \cos(\theta) + v_y \sin(\theta)$$

$$= -u_y \cos(\theta) + u_x \sin(\theta)$$

$$= \frac{1}{r} (-u_y r \cos(\theta) + u_x r \sin(\theta))$$

$$= \frac{1}{r} (-u_y y_\theta - u_x x_0)$$

$$= -\frac{1}{r} u_\theta.$$
CR

Thus

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}$$
 and  $\frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}$ 

Proposition 3.3.10 (Holomorphic functions are continuous.).

f is holomorphic at  $z_0$  iff there exists an  $a \in \mathbb{C}$  such that

$$f(z_0 + h) - f(z_0) - ah = h\psi(h), \quad \psi(h) \stackrel{h \to 0}{\to} 0.$$

In this case,  $a = f'(z_0)$ .

Prove

## 3.3.2 Delbar and the Laplacian

**Definition 3.3.11** (del and delbar operators)

$$\partial := \partial_z := \frac{1}{2} (\partial_x - i \partial_y)$$
 and  $\bar{\partial} := \partial_{\bar{z}} = \frac{1}{2} (\partial_x + i \partial_y)$ .

Moreover,  $f' = \partial f + \bar{\partial} f$ .

Proposition 3.3.12 (Holomorphic iff delbar vanishes).

f is holomorphic at  $z_0$  iff  $\bar{\partial} f(z_0) = 0$ :

$$2\overline{\partial}f := (\partial_x + i\partial_y)(u + iv)$$

$$= u_x + iv_x + iu_y - v_y$$

$$= (u_x - v_y) + i(u_y + v_x)$$

$$= 0$$

by Cauchy-Riemann.

#### 3.3.3 Harmonic Functions and the Laplacian

**Definition 3.3.13** (Laplacian and Harmonic Functions)

A real function of two variables u(x,y) is **harmonic** iff it is in the kernel of the Laplacian operator:

$$\Delta u \coloneqq \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right) u = 0.$$

## Proposition 3.3.14 (Cauchy-Riemann implies holomorphic).

If f = u + iv with  $u, v \in C^1(\mathbb{R})$  satisfying the Cauchy-Riemann equations on  $\Omega$ , then f is holomorphic on  $\Omega$  and

$$f'(z) = \partial f = \frac{1}{2} (u_x + iv_x).$$

## Proposition 3.3.15 (Holomorphic functions have harmonic components).

If f(z) = u(x, y) + iv(x, y) is holomorphic, then u, v are harmonic.

Proof (?).

• By CR,

$$u_x = v_y$$

$$u_y = -v_x$$
.

• Differentiate with respect to x:

$$u_{xx} = v_{yx}$$

$$u_{yx} = -v_{xx}$$
.

• Differentiate with respect to y:

$$u_{xy} = v_{yy}$$

$$u_{yy} = -v_{xy}.$$

• Clairaut's theorem: partials are equal, so

$$u_{xx} - v_{yx} = 0 \implies u_{xx} + u_{yy} = 0$$

$$v_{xx} + u_{yx} = 0 \implies v_{xx} + v_{yy} = 0$$

.

3.3.4 Exercises

## Proposition 3.3.16 (Injectivity Relates to Derivatives).

If  $z_0$  is a zero of f' of order n, then f is (n+1)-to-one in a neighborhood of  $z_0$ .

Proof.

proof

**Exercise 3.3.17** (Zero derivative implies constant) Show that if f' = 0 on a domain  $\Omega$ , then f is constant on  $\Omega$ 

#### **Solution:**

Write f = u + iv, then  $0 = 2f' = u_x + iv_x = u_y - iu_y$ , so grad u = grad v = 0. Show f is constant along every straight line segment L by computing the directional derivative  $\text{grad } u \cdot \mathbf{v} = 0$  along L connecting p, q. Then u(p) = u(q) = a some constant, and v(p) = v(q) = b, so f(z) = a + bi everywhere.

Exercise 3.3.18 (f and fbar holomorphic implies constant)

Show that if f and  $\bar{f}$  are both holomorphic on a domain  $\Omega$ , then f is constant on  $\Omega$ .

#### Solution:

- Strategy: show f' = 0.
- Write f = u + iv. Since f is analytic, it satisfies CR, so

$$u_x = v_y u_y = -v_x.$$

• Similarly write  $\bar{f} = U + iV$  where U = u and V = -v. Since  $\bar{f}$  is analytic, it also satisfies CR , so

$$U_x = V_y U_y = -V_x$$

$$\implies u_x = -v_y \qquad \qquad u_y = v_x.$$

- Add the LHS of these two equations to get  $2u_x = 0 \implies u_x = 0$ . Subtract the right-hand side to get  $-2v_x = 0 \implies v_x = 0$
- Since f is analytic, it is holomorphic, so f' exists and satisfies  $f' = u_x + iv_x$ . But by above, this is zero.
- By the previous exercise,  $f' = 0 \implies f$  is constant.

Exercise 3.3.19 (SS 1.13: Constant real/imaginary/magnitude implies constant) If f is holomorphic on  $\Omega$  and any of the following hold, then f is constant:

- 1.  $\Re(f)$  is constant.
- 2.  $\Im(f)$  is constant.
- 3. |f| is constant.

#### Solution:

#### Part 3:

- Write  $|f| = c \in \mathbb{R}$ .
- If c = 0, done, so suppose c > 0.

- Use  $f\overline{f} = |f|^2 = c^2$  to write  $\overline{f} = c^2/f$ .
- Since  $|f(z)| = 0 \iff f(z) = 0$ , we have  $f \neq 0$  on  $\Omega$ , so  $\overline{f}$  is analytic.
- Similarly f is analytic, and  $f, \bar{f}$  analytic implies f' = 0 implies f is constant.

Finish

## 3.4 Power Series

### Theorem 3.4.1 (Improved Taylor's Theorem).

If f is holomorphic on a region  $\Omega$  with  $\overline{D_R(z_0)} \subseteq \Omega$ , and for every  $z \in D_r(z_0)$ , f has a power series expansion of the following form:

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$$
 where  $a_n = \frac{f^{(n)}(z_0)}{n!} = \frac{1}{2\pi r^n} \int_0^{2\pi} f(z_0 + re^{i\theta}) e^{-in\theta} d\theta$ .

## Proposition 3.4.2 (Power Series are Smooth).

Any power series is smooth (and thus holomorphic) on its disc of convergence, and its derivatives can be obtained using term-by-term differentiation:

$$\frac{\partial}{\partial z} f(z) = \frac{\partial}{\partial z} \sum_{k \ge 0} c_k (z - z_0)^k = \sum_{k \ge 1} k c_k (z - z_0)^k.$$

Moreover, the coefficients are given by

$$c_k = \frac{f^{(n)}(z_0)}{n!}.$$

**Remark 3.4.3:** By an application of the Cauchy integral formula (see S&S 7.1) if f is holomorphic on  $D_R(z_0)$  there is a formula for all  $k \ge 0$  and all 0 < r < R:

$$c_k = \frac{1}{2\pi r^k} \int_0^{2\pi} f(z_0 + re^{i\theta}) e^{-in\theta} d\theta.$$

#### Proposition 3.4.4 (Exponential is uniformly convergent in discs).

 $f(z) = e^z$  is uniformly convergent in any disc in  $\mathbb{C}$ .

Proof.

Apply the estimate

$$|e^z| \le \sum \frac{|z|^n}{n!} = e^{|z|}.$$

Now by the M-test,

$$|z| \le R < \infty \implies \left| \sum \frac{z^n}{n!} \right| \le e^R < \infty.$$

#### Lemma 3.4.5 (Dirichlet's Test).

Given two sequences of real numbers  $\{a_k\}, \{b_k\}$  which satisfy

1. The sequence of partial sums  $\{A_n\}$  is bounded,

$$2. \ b_k \searrow 0.$$

then

$$\sum_{k>1} a_k b_k < \infty.$$

Proof (?).

See http://www.math.uwaterloo.ca/~krdavids/Comp/Abel.pdf

Use summation by parts. For a fixed  $\sum a_k b_k$ , write

$$\sum_{n=1}^{m} x_n Y_n + \sum_{n=1}^{m} X_n y_{n+1} = X_m Y_{m+1}.$$

Set  $x_n := a_n, y_N := b_n - b_{n-1}$ , so  $X_n = A_n$  and  $Y_n = b_n$  as a telescoping sum. Importantly, all  $y_n$  are negative, so  $|y_n| = |b_n - b_{n-1}| = b_{n-1} - b_n$ , and moreover  $a_n b_n = x_n Y_n$  for all n. We have

$$\sum_{n\geq 1} a_n b_n = \lim_{N\to\infty} \sum_{n\leq N} x_n Y_n$$

$$= \lim_{N\to\infty} \sum_{n\leq N} X_N Y_N - \sum_{n\leq N} X_n y_{n+1}$$

$$= -\sum_{n\geq 1} X_n y_{n+1},$$

where in the last step we've used that

$$|X_N| = |A_N| \le M \implies |X_N Y_N| = |X_N| |b_{n+1}| \le M b_{n+1} \to 0.$$

So it suffices to bound the latter sum:

$$\sum_{k \ge n} |X_k y_{k+1}| \le M \sum_{k \ge 1} |y_{k+1}|$$

$$\le M \sum_{k \ge 1} b_k - b_{k+1}$$

$$\le 2M(b_1 - b_{n+1})$$

$$\le 2Mb_1.$$

## Theorem 3.4.6 (Abel's Theorem).

If  $\sum_{k=1}^{\infty} c_k z^j$  converges on |z| < 1 then

$$\lim_{z \to 1^{-}} \sum_{k \in \mathbb{N}} c_k z^k = \sum_{k \in \mathbb{N}} c_k.$$

## Lemma 3.4.7 (Abel's Test).

If  $f(z) := \sum_{k} c_k z^k$  is a power series with  $c_k \in \mathbb{R}^{\geq 0}$  and  $c_k \searrow 0$ , then f converges on  $S^1$  except possibly at z = 1.

**Example 3.4.8** (application of Abel's theorem): What is the value of the alternating harmonic series? Integrate a geometric series to obtain

$$\sum \frac{(-1)^k z^k}{n} = \log(z+1) \qquad |z| < 1.$$

Since  $c_k := (-1)^k/k \searrow 0$ , this converges at z = 1, and by Abel's theorem  $f(1) = \log(2)$ .

**Remark 3.4.9:** The converse to Abel's theorem is false: take  $f(z) = \sum (-z)^n = 1/(1+z)$ . Then  $f(1) = 1 - 1 + 1 - \cdots$  diverges at 1, but 1/1 + 1 = 1/2. So the limit  $s := \lim_{x \to 1^-} f(x)1/2$ , but  $\sum a_n$  doesn't converge to s.

## Proposition 3.4.10 (Summation by Parts).

Setting  $A_n := \sum_{k=1}^n b_k$  and  $B_0 := 0$ ,

$$\sum_{k=m}^{n} a_k b_k = A_n b_n - A_{m-1} b_m - \sum_{k=m}^{n-1} A_k (b_{k+1} - b_k).$$

Compare this to integrating by parts:

$$\int_{a}^{b} fg = F(b)g(b) - F(a)g(a) - \int_{a}^{b} Fg'.$$

Note there is a useful form for taking the product of sums:

$$A_n B_n = \sum_{k=1}^n A_k b_k + \sum_{k=1}^n a_k B_{k-1}.$$

#### Proof (?).

An inelegant proof: define  $A_n := \sum_{k \le n} a_k$ , use that  $a_k = A_k - A_{k-1}$ , reindex, and peel a top/bottom term off of each sum to pattern-match.

Behold:

$$\begin{split} \sum_{m \leq k \leq n} a_k b_k &= \sum_{m \leq k \leq n} (A_k - A_{k-1}) b_k \\ &= \sum_{m \leq k \leq n} A_k b_k - \sum_{m \leq k \leq n} A_{k-1} b_k \\ &= \sum_{m \leq k \leq n} A_k b_k - \sum_{m-1 \leq k \leq n-1} A_k b_{k+1} \\ &= A_n b_n + \sum_{m \leq k \leq n-1} A_k b_k - \sum_{m-1 \leq k \leq n-1} A_k b_{k+1} \\ &= A_n b_n - A_{m-1} b_m + \sum_{m \leq k \leq n-1} A_k b_k - \sum_{m \leq k \leq n-1} A_k b_{k+1} \\ &= A_n b_n - A_{m-1} b_m + \sum_{m \leq k \leq n-1} A_k (b_k - b_{k+1}) \\ &= A_n b_n - A_{m-1} b_m - \sum_{m \leq k \leq n-1} A_k (b_{k+1} - b_k). \end{split}$$

#### Proposition 3.4.11(?).

If f is non-constant, then f' is analytic and the zeros of f' are isolated. If f, g are analytic with f' = g', then f - g is constant.

#### 3.4.1 Exercises: Series

Exercise 3.4.12 (Application of summation by parts)

Use summation by parts to show that  $\sin(n)/n$  converges.

Exercise 3.4.13 (1.20: Series convergence on the circle) Show that

- ∑ kz<sup>k</sup> diverges on S<sup>1</sup>.
   ∑ k<sup>-2</sup>z<sup>k</sup> converges on S<sup>1</sup>.
   ∑ k<sup>-1</sup>z<sup>k</sup> converges on S<sup>1</sup> \ {1} and diverges at 1.

1. Use that  $|z^k| = 1$  and  $\sum c_k z^k < \infty \implies |c_k| \to 0$ , but  $|kz^k| = |k| \to \infty$  here.

- 2. Use that absolutely convergent implies convergent, and  $\sum |k^{-2}z^k| = \sum |k^{-2}|$  converges by the p-test.
- 3. If z=1, this is the harmonic series. Otherwise take  $a_k=1/k, b_k=e^{ik\theta}$  where  $\theta\in(0,2\pi)$ is some constant, and apply Dirichlet's test. It suffices to bound the partial sums of the

 $b_k$ . Recalling that  $\sum_{k \leq N} r^k = (1 - r^{N+1})/(1 - r)$ ,

$$\left\| \sum_{k \le m} e^{ik\theta} \right\| = \left\| \frac{1 - e^{i(m+1)\theta}}{1 - e^{i\theta}} \right\| \le \frac{2}{\|1 - e^{i\theta}\|} := M,$$

which is a constant. Here we've used that two points on  $S^1$  are at most distance 2 from each other.

Exercise 3.4.14 (Laurent expansions inside and outside of a disc)

Expand  $f(z) = \frac{1}{z(z-1)}$  in both

- |z| < 1
- |z| > 1

Solution:

$$\frac{1}{z(z-1)} = -\frac{1}{z}\frac{1}{1-z} = -\frac{1}{z}\sum z^k.$$

and

$$\frac{1}{z(z-1)} = \frac{1}{z^2(1-\frac{1}{z})} = \frac{1}{z^2} \sum \left(\frac{1}{z}\right)^k.$$

Exercise 3.4.15 (Laurent expansions about different points)

Find the Laurent expansion about z = 0 and z = 1 respectively of the following function:

$$f(z) \coloneqq \frac{z+1}{z(z-1)}.$$

Solution:

Note: once you see that everything is in terms of powers of  $(z - z_0)$ , you're essentially done. For z = 0:

$$\frac{z+1}{z(z-1)} = \frac{1}{z} \frac{z+1}{z-1}$$

$$= -\frac{z+1}{z} \frac{1}{1-z}$$

$$= -\left(1 + \frac{1}{z}\right) \sum_{k>0} z^k.$$

For z = 1:

$$\frac{z+1}{z(z-1)} = \frac{1}{z-1} \left( 1 + \frac{1}{z} \right)$$

$$= \frac{1}{z-1} \left( 1 + \frac{1}{1-(1-z)} \right)$$

$$= \frac{1}{z-1} \left( 1 + \sum_{k \ge 0} (1-z)^k \right)$$

$$= \frac{1}{z-1} \left( 1 + \sum_{k \ge 0} (-1)^k (z-1)^k \right).$$

# 4 Cauchy's Theorem

## 4.1 Complex Integrals

**Definition 4.1.1** (Complex Integral)

$$\int_{\gamma} f dz := \int_{I} f(\gamma(t)) \gamma'(t) dt = \int_{\gamma} (u + iv) dx \wedge (-v + iu) dy.$$

Theorem 4.1.2 (Cauchy-Goursat Theorem).

If f is holomorphic on a region  $\Omega$  with  $\pi_1\Omega=1$ , then for any closed path  $\gamma\subseteq\Omega$ ,

$$\int_{\gamma} f(z) \, dz = 0.$$

#### Slogan 4.1.3

Closed path integrals of holomorphic functions vanish.

## 4.2 Applications of Cauchy's Theorem

#### 4.2.1 Integral Formulas and Estimates

See reference

Cauchy's Theorem 24

## Theorem 4.2.1 (Cauchy Integral Formula).

Suppose f is holomorphic on  $\Omega$ , then for any  $z_0 \in \Omega$  and any open disc  $\overline{D_R(z_0)}$  such that  $\gamma := \partial \overline{D_R(z_0)} \subseteq \Omega$ ,

$$f(z_0) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\xi)}{\xi - z_0} d\xi$$

and

$$\frac{\partial^n f}{\partial z^n}(z_0) = \frac{n!}{2\pi i} \int_{\gamma} \frac{f(\xi)}{(\xi - z_0)^{n+1}} d\xi.$$

*Proof.* It follows from a consequence of Cauchy's theorem (see above) that if  $C(z_0, r)$  denotes the circle of radius r around  $z_0$  for a sufficiently small r > 0 then

$$\left| \frac{1}{2\pi i} \int_{C} \frac{f(z)}{z - z_{0}} dz - f(z_{0}) \right| = \left| \frac{1}{2\pi i} \int_{C(z_{0}, r)} \frac{f(z) - f(z_{0})}{z - z_{0}} dz \right| 
= \left| \frac{1}{2\pi i} \int_{0}^{2\pi} \frac{f(z_{0} + re^{i\theta}) - f(z_{0})}{re^{i\theta}} ire^{i\theta} d\theta \right| 
\leq \frac{1}{2\pi} 2\pi \times \sup_{\theta \in [0, 2\pi)} |f(z_{0} + re^{i\theta}) - f(z_{0})| 
\text{(by } ML \text{ inequality)}.$$

As f is continuous it follows that the righthand side goes to zero as r tends to zero. This completes the proof.

Figure 2: image 2021-05-27-16-54-06

Proof (?).

Proof (?).

*Proof.* (\*) Using Cauchy's integral formula we can write that

$$f'(z_0) = \lim_{h \to 0} \frac{f(z_0 + h) - f(z_0)}{h} = \lim_{h \to 0} \frac{1}{2\pi i h} \int_C (\frac{f(z)}{z - z_0 - h} - \frac{f(z)}{z - z_0}) dz$$

$$(C \text{ is so chosen that the point } z_0 + h \text{ is enclosed by } C)$$

$$= \lim_{h \to 0} \frac{1}{2\pi i h} \int_C \frac{f(z)h}{(z - z_0 - h)(z - z_0)} dz.$$

So we need to prove that

$$\left| \int_{C} \frac{f(z)}{(z - z_0 - h)(z - z_0)} dz - \int_{C} \frac{f(z)}{(z - z_0)^2} dz \right|$$

$$= \left| \int_{C} \frac{f(z)h}{(z - z_0 - h)(z - z_0)^2} dz \right| \to 0, \text{ as } h \to 0.$$

We will basically use ML inequality to prove this. Note that, as f is continuous it is bounded on C by M (say). Let  $\alpha = \min\{|z - z_0| : z \in C\}$ . Then  $|z - z_0|^2 \ge \alpha^2$  and  $\alpha \le |z - z_0| = |z - z_0 - h + h| \le |z - z_0 - h| + |h|$  and hence for  $|h| \le \frac{\alpha}{2}$  (after all h is going to be small) we get  $|z - z_0 - h| \ge \alpha - |h| \ge \frac{\alpha}{2}$ . Therefore

$$\left| \int_C \frac{f(z)h}{(z-z_0-h)(z-z_0)^2} dz \right| \le \frac{M|h|l}{\frac{\alpha}{2}\alpha^2} = \frac{2M|h|l}{\alpha^3} \to 0,$$

as  $h \to 0$ . By repeating exactly the same technique we get  $f^2(z_0) = \frac{2!}{2\pi i} \int_C \frac{f(z)}{(z-z_0)^3} dz$  and so on.

Theorem 4.2.2 (Cauchy's Inequality / Cauchy's Estimate).

For  $z_0 \in D_R(z_0) \subset \Omega$ , setting  $M := \sup_{z \in \gamma} |f(z)|$  so  $|f(z)| \leq M$  on  $\gamma$ 

$$\left| f^{(n)}(z_0) \right| \le \frac{n!}{2\pi} \int_0^{2\pi} \frac{M}{R^{n+1}} R \, d\theta = \frac{Mn!}{R^n}.$$

Proof (of Cauchy's inequality).

- Given  $z_0 \in \Omega$ , pick the largest disc  $D_R(z_0) \subset \Omega$  and let  $C = \partial D_R$ .
- Then apply the integral formula.

$$\left| f^{(n)}(z_0) \right| = \left| \frac{n!}{2\pi i} \int_C \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta \right|$$

$$= \left| \frac{n!}{2\pi i} \int_0^{2\pi} \frac{f\left(z_0 + re^{i\theta}\right) rie^{i\theta}}{(re^{i\theta})^{n+1}} d\theta \right|$$

$$\leq \frac{n!}{2\pi} \int_0^{2\pi} \left| \frac{f\left(z_0 + re^{i\theta}\right) rie^{i\theta}}{(re^{i\theta})^{n+1}} \right| d\theta$$

$$= \frac{n!}{2\pi} \int_0^{2\pi} \frac{\left| f\left(z_0 + re^{i\theta}\right) \right|}{r^n} d\theta$$

$$\leq \frac{n!}{2\pi} \int_0^{2\pi} \frac{M}{r^n} d\theta$$

$$= \frac{Mn!}{r^n}.$$

#### **Slogan 4.2.3**

The *n*th Taylor coefficient of an analytic function is at most  $\sup_{|z|=R} |f|/R^n$ .

## Theorem 4.2.4 (Mean Value Property for Holomorphic Functions).

If f is holomorphic on  $D_r(z_0)$ 

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{i\theta}) d\theta = \frac{1}{\pi r^2} \iint_{D_r(z_0)} f(z) dA.$$

Taking the real part of both sides, one can replace f = u + iv with u.

#### 4.2.2 Liouville

#### Theorem 4.2.5 (Liouville's Theorem).

If f is entire and bounded, f is constant.

Proof (of Liouville).

- Since f is bounded,  $f(z) \leq M$  uniformly on  $\mathbb{C}$ .
- Apply Cauchy's estimate for the 1st derivative:

$$|f'(z)| \le \frac{1! ||f||_{C_R}}{R} \le \frac{M}{R} \stackrel{R \to \infty}{\longrightarrow} 0,$$

so f'(z) = 0 for all z.

**Exercise 2.E.** [SSh03, 2.15] Suppose f is continuous and non-zero on  $\overline{\mathbb{D}}$  and holomorphic on  $\mathbb{D}$  such that |f(z)| = 1 for all |z| = 1. Show that f is then constant.

Figure 3: image 2021-05-17-11-54-14

## Exercise 4.2.6 (?)

#### 4.2.3 Continuation Principle

## Theorem 4.2.7 (Continuation Principle / Identity Theorem).

If f is holomorphic on a bounded connected domain  $\Omega$  and there exists a sequence  $\{z_i\}$  with a limit point in  $\Omega$  such that  $f(z_i) = 0$ , then  $f \equiv 0$  on  $\Omega$ .

#### **Slogan 4.2.8**

Two functions agreeing on a set with a limit point are equal on a domain.

Proof (?).

Apply Improved Taylor Theorem?

todo

**Exercise 2.D.** [SSh03, 2.13] If f is holomorphic on a region  $\Omega$  and for each  $z_0 \in \Omega$  at least one coefficient in the power series expansion  $f(z) = \sum_{n=0}^{\infty} c_n (z-z_0)^n$  is zero. Then show that f is a polynomial.

Figure 4: image\_2021-05-17-11-53-33

Exercise 4.2.9 (?)

## 4.3 Exercises

Exercise 4.3.1 (Primitives imply vanishing integral)

Show that if f has a primitive F on  $\Omega$  then  $\int_{\gamma} f = 0$  for every closed curve  $\gamma \subseteq \Omega$ .

#### Exercise 4.3.2 (?)

Prove the uniform limit theorem for holomorphic functions: if  $f_n \to f$  locally uniformly and each  $f_n$  is holomorphic then f is holomorphic.

#### Solution:

This is S&S Theorem 5.2. Statement: if  $f_n \to f$  uniformly locally uniformly on  $\Omega$  then f is holomorphic on  $\Omega$ .

• Let  $D \subset \Omega$  with  $\overline{\mathbb{D}} \subset \Omega$  and  $\Delta \subset D$  be a triangle.

4.3 Exercises 28

- Apply Goursat: ∫<sub>Δ</sub> f<sub>n</sub> = 0.
   f<sub>n</sub> → f uniformly on Δ since it is closed and bounded and thus compact by Heine-Borel, so f is continuous and

$$\lim_{n} \int_{\Delta} f_n = \int_{\Delta} \lim_{n} f_n := \int_{\Delta} f.$$

• Apply Morera's theorem:  $\int_{\Lambda} f$  vanishes on every triangle in  $\Omega$ , so f is holomorphic on

## Exercise 4.3.3 (?)

Prove that if  $f_n \to f$  locally uniformly with  $f_n$  holomorphic, then  $f'_n \to f'$  locally uniformly and f' is holomorphic.

#### Solution:

- Simplifying step: for some reason, it suffices to assume  $f_n \to f$  uniformly on all of  $\Omega$ ?
- Take  $\Omega_R$  to be  $\Omega$  with a buffer of R, so  $d(z,\partial\Omega) > R$  for every  $z \in \overline{\Omega_R}$ .
- It suffices to show the following bound for F any holomorphic function on  $\Omega$ :

$$\sup_{z \in \Omega_R} |F'(z)| \le \frac{1}{R} \sup_{\zeta \in \Omega} |F(\zeta)| \qquad \forall R,$$

where on the right we take the sup over all  $\Omega$ .

- Then take  $F := f_n f$  and  $R \to 0$  to conclude, since the right-hand side is a constant not depending on  $\Omega_R$ .
- For any  $z \in \Omega_R$ , we have  $\overline{D_R(z)} \subseteq \Omega_R$ , so Cauchy's integral formula can be applied:

 $|F'(z)| = \left| \frac{1}{2\pi i} \int_{\partial D_R(z)} \frac{F(\xi)}{(\xi - z)^2} d\xi \right|$   $\leq \frac{1}{2\pi} \int_{\partial D_R(z)} \frac{|F(\xi)|}{|\xi - z|^2} d\xi$   $\leq \frac{1}{2\pi} \int_{\partial D_R(z)} \frac{\sup_{\zeta \in \Omega} |F(\zeta)|}{|\xi - z|^2} d\xi$   $= \frac{1}{2\pi} \sup_{\zeta \in \Omega} |F(\zeta)| \int_{\partial D_R(z)} \frac{1}{R^2} d\xi$   $= \frac{1}{2\pi} \sup_{\zeta \in \Omega} |F(\zeta)| \frac{1}{R^2} \int_{\partial D_R(z)} d\xi$   $= \frac{1}{2\pi} \sup_{\zeta \in \Omega} |F(\zeta)| \frac{1}{R^2} 2\pi R$   $\leq \frac{1}{2\pi} \sup_{\zeta \in \Omega} |F(\zeta)| \frac{1}{R^2} (2\pi R)$   $= \frac{1}{R} \sup_{\zeta \in \Omega} |F(\zeta)|.$ 

Now

$$||f'_n - f'||_{\infty,\Omega_R} \le \frac{1}{R} ||f_n - f||_{\infty,\Omega},$$

where if R is fixed then by uniform convergence of  $f_n \to f$ , for n large enough  $||f_n - f|| < \varepsilon/R$ .

#### 4.4 Morera's Theorem

Theorem 4.4.1 (Morera's Theorem).

If f is continuous on a domain  $\Omega$  and  $\int_T f = 0$  for every triangle  $T \subset \Omega$ , then f is holomorphic.

#### Slogan 4.4.2

If every integral along a triangle vanishes, implies holomorphic.

Corollary 4.4.3 (Sufficient condition for a sequence to converge to a holomorphic function).

If  $\{f_n\}_{n\in\mathbb{N}}$  is a holomorphic sequence on a region  $\Omega$  which uniformly converges to f on every compact subset  $K\subseteq\Omega$ , then f is holomorphic, and  $f'_n\to f'$  uniformly on every such compact subset K.

4.4 Morera's Theorem 30

Proof (?).

Commute limit with integral and apply Morera's theorem.

**Remark 4.4.4:** This can be applied to series of the form  $\sum_{k} f_k(z)$ .

#### 4.4.1 Symmetric Regions

In this section, take  $\Omega$  to be a region symmetric about the real axis, so  $z \in \Omega \iff \bar{z} \in \Omega$ . Partition this set as  $\Omega^+ \subseteq \mathbb{H}, I \subseteq \mathbb{R}, \Omega^- \subseteq \overline{\mathbb{H}}$ .

#### Theorem 4.4.5 (Symmetry Principle).

Suppose that  $f^+$  is holomorphic on  $\Omega^+$  and  $f^-$  is holomorphic on  $\Omega^-$ , and f extends continuously to I with  $f^+(x) = f^-(x)$  for  $x \in I$ . Then the following piecewise-defined function is holomorphic on  $\Omega$ :

$$f(z) := \begin{cases} f^{+}(z) & z \in \Omega^{+} \\ f^{-}(z) & z \in \Omega^{-} \\ f^{+}(z) = f^{-}(z) & z \in I. \end{cases}$$

Proof (?).

Apply Morera?

#### Theorem 4.4.6 (Schwarz Reflection).

If f is continuous and holomorphic on  $\mathbb{H}^+$  and real-valued on  $\mathbb{R}$ , then the extension defined by  $F^-(z) = \overline{f(\overline{z})}$  for  $z \in \mathbb{H}^-$  is a well-defined holomorphic function on  $\mathbb{C}$ .

Proof (?).

Apply the symmetry principle.

**Remark 4.4.7:**  $\mathbb{H}^+$ ,  $\mathbb{H}^-$  can be replaced with any region symmetric about a line segment  $L \subseteq \mathbb{R}$ .

# **5** Zeros and Singularities

#### **Definition 5.0.1** (Singularity)

A point  $z_0$  is an **isolated singularity** if  $f(z_0)$  is undefined but f(z) is defined in a punctured neighborhood  $D(z_0) \setminus \{z_0\}$  of  $z_0$ .

There are three types of isolated singularities:

Zeros and Singularities 31

- Removable singularities
- Poles
- Essential singularities

**Example 5.0.2**(?): The singularities of a rational function are always isolated, since there are finitely many zeros of any polynomial. The function F(z) := Log(z) has a singularity at z = 0 that is **not** isolated, since every neighborhood intersects the branch cut  $(-\infty, 0) \times \{0\}$ , where F is not even defined. The function  $G(z) := 1/\sin(\pi/z)$  has a non-isolated singularity at 0 and isolated singularities at 1/n for all n.

#### **Definition 5.0.3** (Removable Singularities)

If  $z_0$  is a singularity of f, then  $z_0$  is a **removable singularity** iff there exists a holomorphic function g such that f(z) = g(z) in a punctured neighborhood of  $z_0$ . Equivalently,

$$\lim_{z \to z_0} (z - z_0) f(z) = 0.$$

Equivalently, f is bounded on a neighborhood of  $z_0$ .

#### Theorem 5.0.4 (Extension over removable singularities).

If f is holomorphic on  $\Omega \setminus \{z_0\}$  where  $z_0$  is a removable singularity, then there is a unique holomorphic extension of f to all of  $\Omega$ .

Proof (?).

Take  $\gamma$  to be a circle centered at  $z_0$  and use

$$f(z) \coloneqq \int_{\gamma} \frac{f(\xi)}{\xi - z} dx.$$

This is valid for  $z \neq z_0$ , but the right-hand side is analytic. (?)

Revisit

#### Theorem 5.0.5 (Improved Taylor Remainder Theorem).

If f is analytic on a region  $\Omega$  containing  $z_0$ , then f can be written as

$$f(z) = \left(\sum_{k=0}^{n-1} \frac{f^{(k)}(z_0)}{k!} (z - z_0)^k\right) + R_n(z) (z - z_0)^n,$$

where  $R_n$  is analytic.

#### **Definition 5.0.6** (Zeros)

If f is analytic and not identically zero on  $\Omega$  with  $f(z_0) = 0$ , then there exists a nonvanishing holomorphic function g such that

$$f(z) = (z - z_0)^n q(z).$$

We refer to  $z_0$  as a **zero of order** n.

Zeros and Singularities 32

## **Definition 5.0.7** (Poles (and associated terminology))

A pole  $z_0$  of a function f(z) is a zero of  $g(z) := \frac{1}{f(z)}$ . Equivalently,  $\lim_{z \to z_0} f(z) = \infty$ . In this case there exists a minimal n and a holomorphic h such that

$$f(z) = (z - z_0)^{-n} h(z).$$

Such an n is the order of the pole. A pole of order 1 is said to be a simple pole.

## Definition 5.0.8 (Principal Part and Residue)

If f has a pole of order n at  $z_0$ , then there exist a holomorphic G in a neighborhood of  $z_0$  such that

$$f(z) = \frac{a_{-n}}{(z - z_0)^n} + \dots + \frac{a_{-1}}{z - z_0} + G(z) := P(z) + G(z).$$

The term P(z) is referred to as the *principal part of f at z*<sub>0</sub> consists of terms with negative degree, and the *residue* of f at  $z_0$  is the coefficient  $a_{-1}$ .

#### **Definition 5.0.9** (Essential Singularity)

A singularity  $z_0$  is essential iff it is neither removable nor a pole. Equivalently, a Laurent series expansion about  $z_0$  has a principal part with infinitely many terms.

#### Theorem 5.0.10 (Casorati-Weierstrass).

If f is holomorphic on  $\Omega \setminus \{z_0\}$  where  $z_0$  is an essential singularity, then for every  $V \subset \Omega \setminus \{z_0\}$ , f(V) is dense in  $\mathbb{C}$ .

#### Slogan 5.0.11

The image of a punctured disc at an essential singularity is dense in  $\mathbb{C}$ .

#### Proof (of Casorati-Weierstrass).

Pick  $w \in \mathbb{C}$  and suppose toward a contradiction that  $D_R(w) \cap f(V)$  is empty. Consider

$$g(z) \coloneqq \frac{1}{f(z) - w},$$

and use that it's bounded to conclude that  $z_0$  is either removable or a pole for f.

### **Definition 5.0.12** (Singularities at infinity)

For any f holomorphic on an unbounded region, we say  $z = \infty$  is a singularity (of any of the above types) of f if g(z) := f(1/z) has a corresponding singularity at z = 0.

#### **Definition 5.0.13** (Meromorphic)

A function  $f:\Omega\to\mathbb{C}$  is meromorphic iff there exists a sequence  $\{z_n\}$  such that

•  $\{z_n\}$  has no limit points in  $\Omega$ .

Zeros and Singularities 33

6 Residues

- f is holomorphic in  $\Omega \setminus \{z_n\}$ .
- f has poles at the points  $\{z_n\}$ .

Equivalently, f is holomorphic on  $\Omega$  with a discrete set of points delete which are all poles of f.

## Theorem 5.0.14 (Meromorphic implies rational).

Meromorphic functions on  $\mathbb C$  are rational functions.

Proof (?).

Consider f(z) - P(z), subtracting off the principal part at each pole  $z_0$ , to get a bounded entire function and apply Liouville.

#### Theorem 5.0.15 (Riemann Extension Theorem).

A singularity of a holomorphic function is removable if and only if the function is bounded in some punctured neighborhood of the singular point.

# 6 Residues

## 6.1 Basics

Remark 6.1.1: Check: do you need residues at all?? You may be able to just compute an integral!

• Directly by parameterization:

$$\int_{\gamma} f \, dz = \int_{a}^{b} f(z(t)) \, z'(t) \, dt \qquad \text{for } z(t) \text{ a parameterization of } \gamma,$$

• Finding a primitive F, then

$$\int_{\gamma} f = F(b) - F(a).$$

- Note: you can parameterize a circle around  $z_0$  using

$$z = z_0 + re^{i\theta}.$$

Fact 6.1.2 (Integrating  $z^k$  around  $S^1$  powers residues)

The major fact that reduces integrals to residues:

$$\int_{\gamma} z^k\,dz = \int_0^{2\pi} e^{ik\theta}ie^{i\theta}\,d\theta = \int_0^{2\pi} e^{i(k+1)\theta\,d\theta} = \begin{cases} 2\pi i & k=-1\\ 0 & \text{else.} \end{cases}.$$

Residues 34

6 Residues

Thus

$$\int \sum_{k \ge -M} c_k z^k = \sum_{k \ge -M} \int c_k z^k = 2\pi i c_{-1},$$

i.e. the integral picks out the  $c_{-1}$  coefficient in a Laurent series expansion.

Example 6.1.3(?): Consider

$$f(z) \coloneqq \frac{e^{iz}}{1 + z^2}$$

where  $z \neq \pm i$ , and attempt to integrate

$$\int_{\mathbb{R}} f(z) \, dz.$$

Use a semicircular contour  $\gamma_R$  where  $z=Re^{it}$  and check

$$\sup_{z \in \gamma_R} |f(z)| = \max_{t \in [0, \pi} \frac{1}{1 + (Re^{it})^2}$$

$$= \max_{t \in [0, \pi} \frac{1}{1 + R^2 e^{2it}}$$

$$= \frac{1}{R^2 - 1}.$$

## 6.2 Estimates

Proposition 6.2.1 (Length bound / ML Estimate).

$$\left| \int_{\gamma} f \right| \leq ML := \sup_{z \in \gamma} |f| \cdot \operatorname{length}(\gamma).$$

Proof (?).

$$\left| \int_{\gamma} f(z) dz \right| \leq \sup_{t \in [a,b]} \left| f(z(t)) \right| \int_{a}^{b} \left| z'(t) \right| dt \leq \sup_{z \in \gamma} \left| f(z) \right| \cdot \operatorname{length}(\gamma).$$

Proposition 6.2.2 (Jordan's Lemma).

6.2 Estimates 35

6 Residues

Suppose that  $f(z) = e^{iaz}g(z)$  for some g, and let  $C_R := \{z = Re^{it} \mid t \in [0, \pi]\}$ . Then

$$\left| \int_{C_R} f(z) \, dz \right| \le \frac{\pi M_R}{a}$$

where  $M_R := \sup_{t \in [0,\pi]} |g(Re^{it})|$ .

Proof(?).

$$\left| \int_{C_R} f(z) \, dz \right| = \left| \int_{C_R} e^{iaz} g(z) \, dz \right|$$

$$= \left| \int_{[0,\pi]} e^{ia(Re^{it})} g(Re^{it}) i Re^{it} \, dt \right|$$

$$\leq \int_{[0,\pi]} \left| e^{ia(Re^{it})} g(Re^{it}) i Re^{it} \right| \, dt$$

$$= R \int_{[0,\pi]} \left| e^{ia(Re^{it})} g(Re^{it}) \right| \, dt$$

$$\leq R M_R \int_{[0,\pi]} \left| e^{ia(Re^{it})} \right| \, dt$$

$$= R M_R \int_{[0,\pi]} e^{\Re(iaRe^{it})} \, dt$$

$$= R M_R \int_{[0,\pi]} e^{\Re(iaR(\cos(t)+i\sin(t)))} \, dt$$

$$= R M_R \int_{[0,\pi]} e^{-aR\sin(t)} \, dt$$

$$= 2R M_R \int_{[0,\pi/2]} e^{-aR\sin(t)} \, dt$$

$$\leq 2R M_R \int_{[0,\pi/2]} e^{-aR(\frac{2t}{\pi})} \, dt$$

$$= 2R M_R \left(\frac{\pi}{2aR}\right) \left(1 - e^{-aR}\right)$$

$$= \frac{\pi M_R}{a}.$$

where we've used that on  $[0, \pi/2]$ , there is an inequality  $2t/\pi \le \sin(t)$ . This is obvious from a picture, since  $\sin(t)$  is a height on  $S^1$  and  $2t/\pi$  is a height on a diagonal line:

6.2 Estimates 36




Figure 5:  $image_2021-06-09-01-29-22$ 

# 6.3 Residue Formulas

# Theorem 6.3.1 (The Residue Theorem).

Let f be meromorphic on a region  $\Omega$  with poles  $\{z_1, z_2, \dots, z_N\}$ . Then for any  $\gamma \in \Omega \setminus \{z_1, z_2, \dots, z_N\}$ ,

$$\frac{1}{2\pi i} \int_{\gamma} f(z) dz = \sum_{j=1}^{N} n_{\gamma}(z_j) \operatorname{Res}_{z=z_j} f.$$

If  $\gamma$  is a toy contour, then

$$\frac{1}{2\pi i} \int_{\gamma} f \, dz = \sum_{j=1}^{N} \operatorname{Res}_{z=z_{j}} f.$$

Proposition 6.3.2 (Residue formula for higher order poles).

If f has a pole  $z_0$  of order n, then

$$\operatorname{Res}_{z=z_0} f = \lim_{z \to z_0} \frac{1}{(n-1)!} \left( \frac{\partial}{\partial z} \right)^{n-1} (z - z_0)^n f(z).$$

Proposition 6.3.3 (Residue formula for simple poles).

As a special case, if  $z_0$  is a simple pole of f, then

$$\operatorname{Res}_{z=z_0} f = \lim_{z \to z_0} (z - z_0) f(z).$$

Corollary 6.3.4(Better derivative formula that sometimes works for simple poles). If additionally f = g/h where  $h(z_0) = 0$  and  $h'(z_0) \neq 0$ ,

Res 
$$_{z=z_0} \frac{g(z)}{h(z)} = \frac{g(z_0)}{h'(z_0)}.$$

Proof (?).

Apply L'Hopital:

$$(z - z_0) \frac{g(z)}{h(z)} = \frac{(z - z_0)g(z)}{h(z)} \stackrel{LH}{=} \frac{g(z) + (z - z_0)g'(z)}{h'(z)} \stackrel{z \to z_0}{\longrightarrow} \frac{g(z_0)}{h'(z_0)}.$$

Example 6.3.5 (Residue of a simple pole (order 1)): Let  $f(z) = \frac{1}{1+z^2}$ , then  $g(z) = 1, h(z) = 1+z^2$ , and h'(z) = 2z so that  $h'(i) = 2i \neq 0$ . Thus

$$\operatorname{Res}_{z=i} \frac{1}{1+z^2} = \frac{1}{2i}.$$

#### 6.3.1 Exercises

Some good computations here.

#### Exercise 6.3.6

Show that the complex zeros of  $f(z) := \sin(\pi z)$  are exactly  $\mathbb{Z}$ , and each is order 1. Calculate the residue of  $1/\sin(\pi x)$  at  $z = n \in \mathbb{Z}$ .

Exerci

**Exercise 3.A.** [SSh03, 3.1] Show that the complex zeros of  $\sin \pi z$  are exactly at the integers, and are each of order 1. Calculate the residue of  $1/\sin \pi x$  are  $z=n\in\mathbb{Z}$ .

Exercise 3.C. [SSh03, 3.8] Prove that

$$\int_0^{2\pi} \frac{d\theta}{a + b\cos\theta} = \frac{2\pi}{\sqrt{a^2 - b^2}}$$

Exercise 6.3.7 (?)

$$\int_{\mathbb{R}} \frac{1}{(1+x^2)^2} \, dx.$$

Solution:

• Factor  $(1+z^2)^2 = ((z-i)(z+i))^2$ , so f has poles at  $\pm i$  of order 2.

• Take a semicircular contour  $\gamma := I_R \cup D_R$ , then  $f(z) \approx 1/z^4 \to 0$  for large R and  $\int_{D_R} f \to 0$ .

• Note  $\int_{I_R} f \to \int_{\mathbb{R}} f$ , so  $\int_{\gamma} f \to \int_{\mathbb{R}} f$ .

•  $\int_{\gamma} f = 2\pi i \sum_{z_0} \underset{z=z_0}{\text{Res }} f$ , and  $z_0 = i$  is the only pole in this region.

• Compute

$$\operatorname{Res}_{z=i} f = \lim_{z \to i} \frac{1}{(2-1)!} \frac{\partial}{\partial z} (z-i)^2 f(z)$$

$$= \lim_{z \to i} \frac{\partial}{\partial z} \frac{1}{(z+i)^2}$$

$$= \lim_{z \to i} \frac{-2}{(z+i)^3}$$

$$= -\frac{2}{(2i)^3}$$

$$= \frac{1}{4i}$$

$$\implies \int_{\gamma} f = \frac{2\pi i}{4i} = \pi/2,$$

Exercise 6.3.8 (?)

Use a direct Laurent expansion to show

$$\operatorname{Res}_{z=0} \frac{1}{z - \sin(z)} = \frac{3!}{5 \cdot 4}.$$

Note the necessity: one doesn't know the order of the pole at zero, so it's unclear how many derivatives to take.

6.3 Residue Formulas 39

 $\mathbf{E}\mathbf{x}\mathbf{e}$ 

#### Solution:

Expand:

$$\frac{1}{z - \sin(z)} = z^{-1} \left( 1 - z^{-1} \sin(z) \right)^{-1}$$

$$= z^{-1} \left( 1 - z^{-1} \left( z - \frac{1}{3!} z^3 + \frac{1}{5!} z^5 - \cdots \right) \right)^{-1}$$

$$= z^{-1} \left( 1 - \left( 1 - \frac{1}{3!} z^2 + \frac{1}{5!} z^4 - \cdots \right) \right)^{-1}$$

$$= z^{-1} \left( \frac{1}{3!} z^2 - \frac{1}{5!} z^4 + \cdots \right)^{-1}$$

$$= z^{-1} \cdot 3! z^{-2} \left( 1 - \frac{1}{5!/3!} z^2 + \cdots \right)^{-1}$$

$$= \frac{3!}{z^3} \left( \frac{1}{1 - \left( \frac{1}{5 \cdot 4} z^2 + \cdots \right)} \right)$$

$$= \frac{3!}{z^3} \left( 1 + \left( \frac{1}{5 \cdot 4} z^2 \right) + \left( \frac{1}{5 \cdot 4} z^2 \right)^2 + \cdots \right)$$

$$= 3! z^{-3} + \frac{3!}{5 \cdot 4} z^{-1} + O(z)$$

Exercise 6.3.9 (?)

Compute

$$\operatorname{Res}_{z=0} \frac{1}{z^2 \sin(z)}.$$

Solution:

First expand  $(\sin(z))^{-1}$ :

$$\frac{1}{\sin(z)} = \left(z - \frac{1}{3!}z^3 + \frac{1}{5!}z^5 - \cdots\right)^{-1}$$

$$= z^{-1} \left(1 - \frac{1}{3!}z^2 + \frac{1}{5!}z^4 - \cdots\right)^{-1}$$

$$= z^{-1} \left(1 + \left(\frac{1}{3!}z^2 - \frac{1}{5!}z^4 + \cdots\right) + \left(\frac{1}{3!}z^2 - \cdots\right)^2 + \cdots\right)$$

$$= z^{-1} \left(1 + \frac{1}{3!}z^2 \pm O(z^4)\right),$$

using that  $(1-x)^{-1} = 1 + x + x^2 + \cdots$ 

Thus

$$z^{-2} \left( \sin(z) \right)^{-1} = z^{-2} \cdot z^{-1} \left( 1 + \frac{1}{3!} z^2 \pm O(z^4) \right)$$
$$= z^{-3} + \frac{1}{3!} z^{-1} + O(z).$$

**Exercise 6.3.10** (Keyhole contour and ML estimate) Compute

$$\int_{[0,\infty]} \frac{\log(x)}{(1+x^2)^2} \, dx.$$

#### Solution:

Factor  $(1+z^2)^2 = (z+i^2(z-i)^2)$ . Take a keyhole contour similar to the following:




Figure 6: image\_2021-06-09-02-11-59

Show that outer radius R and inner radius  $\rho$  circles contribute zero in the limit by the ML estimate? Compute the residues by just applying the formula and manually computing derivatives:

$$\begin{aligned} \operatorname*{Res}_{z=\pm i} f(z) &= \lim_{z \to \pm i} \frac{\partial}{\partial z} \frac{\log^2(z)}{(z \pm i)^2} \\ &= \lim_{z \to \pm i} \frac{2 \log(z) (z \pm i)^2 - 2 (z \pm i)^2 \log^2(z)}{((z \pm i)^2)^2} \\ &= \frac{2 \log(\pm i) (\pm 2i)^2 - 2 (\pm 2i)^2 \log^2(\pm i)}{(\pm 2i)^4} \\ &=_? \frac{\pi}{4} \pm \frac{i\pi^2}{16}. \end{aligned}$$

See p.4: http://www.math.toronto.edu/mnica/complex1.pdf

Exercise 6.3.11 (Sinc Function)

Show

$$\int_{(0,\infty)} \frac{\sin(x)}{x} \, dx = \frac{\pi}{2}.$$

Solution

Take an indented semicircle. Let I be the original integral, then

$$I = \frac{1}{2i} \int_{\mathbb{R}} \frac{e^{iz} - 1}{z} \, dz.$$

# 6.4 Argument Principle

**Definition 6.4.1** (Winding Number)

For  $\gamma \subseteq \Omega$  a closed curve not passing through a point  $z_0$ , the winding number of  $\gamma$  about  $z_0$  is defined as

$$n_{\gamma}(z_0) \coloneqq \frac{1}{2\pi i} \int_{\gamma} \frac{1}{\xi - z_0} d\xi.$$

Theorem 6.4.2 (Argument Principle).

For f meromorphic in  $\gamma^{\circ}$  with zeros  $\{z_j\}$  and poles  $\{p_k\}$  repeated with multiplicity where  $\gamma$  does not intersect any zeros or poles, then

$$\Delta_{\gamma} \arg f(z) := \frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = \sum_{i} n_{\gamma}(z_{i}) - \sum_{k} n_{\gamma}(p_{k}) = Z_{f} - P_{f},$$

where  $Z_f$  and  $P_f$  are the number of zeros and poles respectively enclosed by  $\gamma$ , counted with multiplicity.

Proof (?).

Residue formula applied to  $\frac{f'}{f}$ ?

Corollary 6.4.3 (Rouché's Theorem).

If f, g are analytic on a domain  $\Omega$  with finitely many zeros in  $\Omega$  and  $\gamma \subset \Omega$  is a closed curve surrounding each point exactly once, where |g| < |f| on  $\gamma$ , then f and f + g have the same number of zeros.

Alternatively:

Suppose f = g + h with  $g \neq 0, \infty$  on  $\gamma$  with |g| > |h| on  $\gamma$ . Then

$$\Delta_{\gamma} \arg(f) = \Delta_{\gamma} \arg(h)$$
 and  $Z_f - P_f = Z_g - P_g$ .

6.4 Argument Principle 42

Rouche

Prove

## Corollary 6.4.4 (Open Mapping).

Any holomorphic non-constant map is an open map.

## Corollary 6.4.5 (Maximum Modulus).

If f is holomorphic and nonconstant on an open connected region  $\Omega$ , then |f| can not attain a maximum on  $\Omega$ . If  $\Omega$  is bounded and f is continuous on  $\overline{\Omega}$ , then  $\max |f|$  occurs on  $\partial\Omega$ .

Conversely, if f attains a local supremum at  $z_0 \in \Omega$ , then f is constant on  $\Omega$ .

# Corollary 6.4.6(?).

If f is nonzero on  $\Omega$ , then f attains a minimum on  $\partial\Omega$ . This follows from applying the MMP to 1/f.

#### 6.4.1 Exercises

Exercise 3.E. [SSh03, 3.14] Prove that all entire functions that are also injective take the form f(z) = az + b with  $a, b \in \mathbb{C}$  and  $a \neq 0$ .

Figure 7: image 2021-05-17-13-33-55

# Rouche

# 7.1 Counting Zeros

Example 7.1.1:

- Take  $P(z) = z^4 + 6z + 3$ .
- On |z| < 2:
  - Set  $f(z) = z^4$  and g(z) = 6z + 3, then  $|g(z)| \le 6|z| + 3 = 15 < 16 = |f(z)|$ .
  - So P has 4 zeros here.
- On |z| < 1:

  - Set f(z) = 6z and  $g(z) = z^4 + 3$ . Check  $|g(z)| \le |z|^4 + 3 = 4 < 6 = |f(z)|$ .

Rouche 43 - So P has 1 zero here.

## Example 7.1.2:

- Claim: the equation  $\alpha z e^z = 1$  where  $|\alpha| > e$  has exactly one solution in  $\mathbb{D}$ .
- Set  $f(z) = \alpha z$  and  $g(z) = e^{-z}$ .
- Estimate at |z| = 1 we have  $|g| = |e^{-z}| = e^{-\Re(z)} \le e^1 < |\alpha| = |f(z)|$
- f has one zero at  $z_0 = 0$ , thus so does f + g.

# 8 | Conformal Maps

#### 8.1 Linear Fractional Transformations

## **Definition 8.1.1** (Conformal Map / Biholomorphism)

A map f is **conformal** on  $\Omega$  iff f is complex-differentiable,  $f'(z) \neq 0$  for  $z \in \Omega$ , and f preserves signed angles (so f is orientation-preserving). Conformal implies holomorphic, and a bijective conformal map has a holomorphic inverse. A bijective conformal map  $f: U \to V$  is called a **biholomorphism**, and we say U and V are **biholomorphic**. Self-biholomorphisms of a domain  $\Omega$  form a group  $\mathrm{Aut}_{\mathbb{C}}(\Omega)$ .

**Remark 8.1.2:** There is an oft-used weaker condition that  $f'(z) \neq 0$  for any point. Note that that this condition alone doesn't necessarily imply f is holomorphic, since anti-holomorphic maps may be nonzero derivative. For example, take  $f(z) = \bar{z}$ , so f(x+iy) = x - iy – this does not satisfy the Cauchy-Riemann equations.

**Remark 8.1.3:** A bijective holomorphic map automatically has a holomorphic inverse. This can be weakened: an injective holomorphic map satisfies  $f'(z) \neq 0$  and  $f^{-1}$  is well-defined on its range and holomorphic.

**Definition 8.1.4** (Linear fractional transformation / Mobius transformation) A map of the following form is a *linear fractional transformation*:

$$T(z) = \frac{az+b}{cz+d},$$

where the denominator is assumed to not be a multiple of the numerator. These have inverses given by

$$T^{-1}(w) = \frac{dw - b}{-cw + a}.$$

#### Proposition 8.1.5(?).

Given any three points  $z_1, z_2, z_3$ , the following Mobius transformation sends them to  $1, 0, \infty$ 

Conformal Maps 44

respectively:

$$f(z) := \frac{(z - z_2)(z_1 - z_3)}{(z - z_3)(z_1 - z_2)}.$$

Such a map is sometimes denoted  $(z, z_1, z_2, z_3)$ .

# Example 8.1.6(?):

- $(z, i, 1, -1) : \mathbb{D} \to \mathbb{H}$
- $(z,0,-1,1): \mathbb{D} \cap \mathbb{H} \to Q_1.$

# Theorem 8.1.7 (Cayley Transform).

The fractional linear transformation given by  $F(z) = \frac{i-z}{i+z}$  maps  $\mathbb{D} \to \mathbb{H}$  with inverse  $G(w) = i\frac{1-w}{1+w}$ .

# Theorem 8.1.8 (Classification of Conformal Maps).

There are 8 major types of conformal maps:

Type/Domains	Formula
Translation/Dilation/Rotation	$z \mapsto e^{i\theta}(cz+h)$
Sectors to sectors	$z \mapsto z^n$
$\mathbb{D}_{\frac{1}{2}} \to \mathbb{H}_{\frac{1}{2}}$ , the first quadrant	$z \mapsto \frac{1+z}{1-z}$
$\mathbb{H}  o S$	$z\mapsto \log(z)$
$\mathbb{D}_{\frac{1}{2}} \to S_{\frac{1}{2}}$	$z \mapsto \log(z)$
$egin{aligned} \mathbb{D}_{rac{1}{2}} & ightarrow S_{rac{1}{2}} \ S_{rac{1}{2}} & ightarrow \mathbb{D}_{rac{1}{2}} \end{aligned}$	$z \mapsto e^{iz}$
$\mathbb{D}_{rac{1}{2}} o\mathbb{H}$	$z \mapsto \frac{1}{2} \left( z + \frac{1}{z} \right)$
$S_{rac{1}{2}}  o \mathbb{H}$	$z\mapsto\sin(z)$

#### Theorem 8.1.9 (Characterization of conformal maps).

Conformal maps  $\mathbb{D} \to \mathbb{D}$  have the form

$$g(z) = \lambda \frac{1-a}{1-\bar{a}z}, \quad |a| < 1, \quad |\lambda| = 1.$$

# 8.2 Schwarz

#### Theorem 8.2.1 (Schwarz Lemma).

If  $f: \mathbb{D} \to \mathbb{D}$  is holomorphic with f(0) = 0, then

1. 
$$|f(z)| \leq |z|$$
 for all  $z \in \mathbb{D}$ 

8.2 Schwarz 45

2. 
$$|f'(0)| \le 1$$
.

Moreover, if

- $|f(z_0)| = |z_0|$  for any  $z_0 \in \mathbb{D}$ , or |f'(0)| = 1,

then f is a rotation.

Proof (?).

Apply the maximum modulus principle to f(z)/z.

Exercise 8.2.2 (?)

Show that  $\operatorname{Aut}_{\mathbb{C}}(\mathbb{C}) = \{ z \mapsto az + b \mid a \in \mathbb{C}^{\times}, b \in \mathbb{C} \}.$ 

Theorem 8.2.3 (Biholomorphisms of the disc).

$$\operatorname{Aut}_{\mathbb{C}}(\mathbb{D}) = \left\{ z \mapsto e^{i\theta} \left( \frac{\alpha - z}{1 - \overline{\alpha}z} \right) \right\}.$$

Proof (?).

Schwarz lemma.

Theorem 8.2.4(?).

$$\operatorname{Aut}_{\mathbb{C}}(\mathbb{H}) = \left\{ z \mapsto \frac{az+b}{cz+d} \mid a, b, c, d \in \mathbb{C}, ad-bc = 1 \right\} \cong \operatorname{PSL}_{2}(\mathbb{R}).$$

# 8.3 By Type

#### 8.3.1 Plane to Disc

$$\varphi: \mathbb{H} \to \mathbb{D}$$
 
$$\varphi(z) = \frac{z-i}{z+i} \qquad f^{-1}(z) = i\left(\frac{1+w}{1-w}\right).$$

8.3 By Type 46


#### 8.3.2 Sector to Disc

For  $S_{\alpha} := \{z \in \mathbb{C} \mid 0 < \arg(z) < \alpha\}$  an open sector for  $\alpha$  some angle, first map the sector to the half-plane:

$$g: S_{\alpha} \to \mathbb{H}$$
$$g(z) = z^{\frac{\pi}{\alpha}}.$$

Then compose with a map  $\mathbb{H} \to \mathbb{D}$ :

$$f: S_{\alpha} \to \mathbb{D}$$
 
$$f(z) = (\varphi \circ g)(z) = \frac{z^{\frac{\pi}{\alpha}} - i}{z^{\frac{\pi}{\alpha}} + i}.$$




#### 8.3.3 Strip to Disc

- Map to horizontal strip by rotation  $z \mapsto \lambda z$ .
- Map horizontal strip to sector by  $z\mapsto e^z$
- Map sector to  $\mathbb{H}$  by  $z \mapsto z^{\frac{\pi}{\alpha}}$ .
- Map  $\mathbb{H} \to \mathbb{D}$ .

$$e^z: \mathbb{R} \times (0,\pi) \to \mathbb{R} \times (0,\infty).$$

# 9 | Schwarz Reflection

# 10 | Schwarz Lemma



- 11 | Linear Fractional Transformations
- 12 | Montel's Theorem
- 13 Unsorted Theorems

#### Theorem 13.0.1 (Riemann Mapping).

If  $\Omega$  is simply connected, nonempty, and not  $\mathbb{C}$ , then for every  $z_0 \in \Omega$  there exists a unique conformal map  $F: \Omega \to \mathbb{D}$  such that  $F(z_0) = 0$  and  $F'(z_0) > 0$ .

Thus any two such sets  $\Omega_1, \Omega_2$  are conformally equivalent.

#### Theorem 13.0.2 (Riemann's Removable Singularity Theorem).

If f is holomorphic on  $\Omega$  except possibly at  $z_0$  and f is bounded on  $\Omega \setminus \{z_0\}$ , then  $z_0$  is a removable singularity.

#### Theorem 13.0.3 (Little Picard).

If  $f: \mathbb{C} \to \mathbb{C}$  is entire and nonconstant, then  $\operatorname{im}(f)$  is either  $\mathbb{C}$  or  $\mathbb{C} \setminus \{z_0\}$  for some point  $z_0$ .

#### Corollary 13.0.4.

The ring of holomorphic functions on a domain in  $\mathbb{C}$  has no zero divisors.

*Proof* . ???

Find the proof!

Schwarz Lemma 48

#### Morera

Proposition 13.0.5 (Bounded Complex Analytic Functions form a Banach Space). For  $\Omega \subseteq \mathbb{C}$ , show that  $A(\mathbb{C}) := \{ f : \Omega \to \mathbb{C} \mid f \text{ is bounded} \}$  is a Banach space.

Proof.

Apply Morera's Theorem and Cauchy's Theorem

# 14 | Appendix: Proofs of the Fundamental Theorem of Algebra

#### 14.0.1 Fundamental Theorem of Algebra: Argument Principle

- Let  $P(z) = a_n z^n + \cdots + a_0$  and g(z) = P'(z)/P(z), note P is holomorphic
- Since  $\lim_{|z|\to\infty} P(z) = \infty$ , there exist an R > 0 such that P has no roots in  $\{|z| \ge R\}$ .
- Apply the argument principle:

$$N(0) = \frac{1}{2\pi i} \oint_{|\xi|=R} g(\xi) d\xi.$$

- Check that  $\lim_{|z \to \infty|} zg(z) = n$ , so g has a simple pole at  $\infty$
- Then g has a Laurent series  $\frac{n}{z} + \frac{c_2}{z^2} + \cdots$  Integrate term-by-term to get N(0) = n.

#### 14.0.2 Fundamental Theorem of Algebra: Rouche's Theorem

- Let  $P(z) = a_n z^n + \dots + a_0$  Set  $f(z) = a_n z^n$  and  $g(z) = P(z) f(z) = a_{n-1} z^{n-1} + \dots + a_0$ , so f + g = P. Choose  $R > \max\left(\frac{|a_{n-1}| + \dots + |a_0|}{|a_n|}, 1\right)$ , then

$$\begin{split} |g(z)| &\coloneqq |a_{n-1}z^{n-1} + \dots + a_1z + a_0| \\ &\le |a_{n-1}z^{n-1}| + \dots + |a_1z| + |a_0| \quad \text{by the triangle inequality} \\ &= |a_{n-1}| \cdot |z^{n-1}| + \dots + |a_1| \cdot |z| + |a_0| \\ &= |a_{n-1}| \cdot R^{n-1} + \dots + |a_1|R + |a_0| \\ &\le |a_{n-1}| \cdot R^{n-1} + |a_{n-2}| \cdot R^{n-1} + \dots + |a_1| \cdot R^{n-1} + |a_0| \cdot R^{n-1} \quad \text{since } R > 1 \implies R^{a+b} \ge R^a \\ &= R^{n-1} \left( |a_{n-1}| + |a_{n-2}| + \dots + |a_1| + |a_0| \right) \\ &\le R^{n-1} \left( |a_n| \cdot R \right) \quad \text{by choice of } R \\ &= R^n |a_n| \\ &= |a_n z^n| \\ &\coloneqq |f(z)| \end{split}$$

• Then  $a_n z^n$  has n zeros in |z| < R, so f + g also has n zeros.

#### 14.0.3 Fundamental Theorem of Algebra: Liouville's Theorem

- Suppose p is nonconstant and has no roots, then  $\frac{1}{p}$  is entire. We will show it is also bounded and thus constant, a contradiction.
- Write  $p(z) = z^n \left( a_n + \frac{a_{n-1}}{z} + \dots + \frac{a_0}{z^n} \right)$
- Outside a disc
  - Note that  $p(z) \stackrel{z \to \infty}{\to} \infty$ . so there exists an R large enough such that  $|p(z)| \ge \frac{1}{A}$  for any fixed chosen constant A.
  - Then  $|1/p(z)| \leq A$  outside of |z| > R, i.e. 1/p(z) is bounded there.
- Inside a disc:
  - p is continuous with no roots and thus must be bounded below on |z| < R.
  - p is entire and thus continuous, and since  $\overline{D}_r(0)$  is a compact set, p achieves a min A there
  - Set  $C := \min(A, B)$ , then  $|p(z)| \ge C$  on all of  $\mathbb C$  and thus  $|1/p(z)| \le C$  everywhere.
  - So 1/p(z) is bounded an entire and thus constant by Liouville's theorem but this forces p to be constant.  $\mathcal{I}$


#### 14.0.4 Fundamental Theorem of Algebra: Open Mapping Theorem


- p induces a continuous map  $\mathbb{CP}^1 \to \mathbb{CP}^1$
- The continuous image of compact space is compact;
- Since the codomain is Hausdorff space, the image is closed.
- p is holomorphic and non-constant, so by the Open Mapping Theorem, the image is open.
- Thus the image is clopen in  $\mathbb{CP}^1$ .

Appendix

- The image is nonempty, since  $p(1) = \sum a_i \in \mathbb{C}$
- $\mathbb{CP}^1$  is connected
- But the only nonempty clopen subset of a connected space is the entire space.
- So p is surjective, and  $p^{-1}(0)$  is nonempty.
- So p has a root.

# **Appendix**





• Circle:  $x^2 + y^2 = r^2$ • Ellipse:  $\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1$ 

• Hyperbola:  $\left(\frac{x}{a}\right)^2 - \left(\frac{y}{b}\right)^2 = 1$ 

- Rectangular Hyperbola:  $xy = \frac{c^2}{2}$ .

• Parabola:  $-4ax + y^2 = 0$ .

Mnemonic: Write  $f(x,y) = Ax^2 + Bxy + Cy^2 + \cdots$ , then consider the discriminant  $\Delta = B^2 - 4AC$ :

•  $\Delta < 0 \iff \text{ellipse}$ 

 $-\Delta < 0$  and  $A = C, B = 0 \iff$  circle

•  $\Delta = 0 \iff \text{parabola}$ 

•  $\Delta > 0 \iff \text{hyperbola}$ 

#### Fact 15.1.2 (Completing the square)

$$x^{2} - bx = (x - s)^{2} - s^{2}$$
 where  $s = \frac{b}{2}$   
 $x^{2} + bx = (x + s)^{2} - s^{2}$  where  $s = \frac{b}{2}$ .

#### Fact 15.1.3

The sum of the interior angles of an *n*-gon is  $(n-2)\pi$ , where each angle is  $\frac{n-2}{n}\pi$ .

Appendix 51 15 Appendix

# **Definition 15.1.4** (The Dirichlet Problem)

Given a bounded piecewise continuous function  $u: S^1 \to \mathbb{R}$ , is there a unique extension to a continuous harmonic function  $\tilde{u}: \mathbb{D} \to \mathbb{R}$ ?

**Remark 15.1.5:** More generally, this is a boundary value problem for a region where the *values* of the function on the boundary are given. Compare to prescribing conditions on the normal vector on the boundary, which would be a Neumann BVP. Why these show up: a harmonic function on a simply connected region has a harmonic conjugate, and solutions of BVPs are always analytic functions with harmonic real/imaginary parts.

**Example 15.1.6** (Dirichlet problem on the strip): See section 27, example 1 in Brown and Churchill. On the strip  $(x, y) \in (0, \pi) \times (0, \infty)$ , set up the BVP for temperature on a thin plate with no sinks/sources:

$$\Delta T = 0 \qquad T(0, y) = 0, T(\pi, y) = 0 \ \forall y$$

$$T(x,0) = \sin(x)$$
 
$$T(x,y) \stackrel{y \to \infty}{\longrightarrow} 0.$$

Then the following function is harmonic on  $\mathbb{R}^2$  and satisfies that Dirichlet problem:

$$T(x,y) = e^{-y}\sin(x) = \Re(-ie^{iz}) = \Im(e^{iz}).$$

#### **Definition 15.1.7** (Logarithmic Derivative)

The **logarithmic derivative** of f is  $(\ln f)' = f'/f$ .

**Remark 15.1.8:** Why this is useful: deriving the argument principle. If f has a pole of order n at  $z_0$ , then write  $f(z) = (z - z_0)^{-n} g(z)$  with g analytic in a neighborhood of  $z_0$ . Then a direct computation of the derivatives will show

$$(\ln f)' := \frac{f'(z)}{f(z)} = -\frac{n}{z - z_0} + \frac{g'(z)}{g(z)}.$$

Thus

$$\frac{1}{2\pi i} \int_{\gamma} (\ln f)' = -n,$$

for  $\gamma$  a small circle about  $z_0$ . A similar argument for  $z_0$  a **zero** of f yields

$$\frac{1}{2\pi i} \int_{\gamma} h = +n.$$

## Exercise 15.1.9 (?)

Show that there is no continuous square root function defined on all of  $\mathbb{C}$ .

15.1 Misc Basic Algebra 52

#### Solution:

Suppose  $f(z)^2 = z$ . Then f is a section to the covering map

$$p: \mathbb{C}^{\times} \to \mathbb{C}^{\times}$$
$$z \mapsto z^2,$$

so  $p \circ f = \text{id}$ . Using  $\pi_1(\mathbb{C}^\times) = \mathbb{Z}$ , the induced maps are  $p_*(1) = 2$  and  $f_*(1) = n$  for some  $n \in \mathbb{Z}$ . But then  $p_* \circ f_*$  is multiplication by 2n, contradicting  $p_* \circ f_* = \text{id}$  by functoriality.

#### Theorem 15.1.10 (Uniformization).

Every Riemann surface S is the quotient of a free proper holomorphic action of a group G on the universal cover  $\tilde{S}$  of S, so  $S \cong \tilde{S}/G$  is a biholomorphism. Moreover,  $\tilde{S}$  is biholomorphic to either

- $\mathbb{CP}^1$
- C
- D

Basics

- Show that  $\frac{1}{z}\sum_{k=1}^{\infty}\frac{z^k}{k}$  converges on  $S^1\setminus\{1\}$  using summation by parts.
- Show that any power series is continuous on its domain of convergence.
- Show that a uniform limit of continuous functions is continuous.

??

- Show that if f is holomorphic on  $\mathbb D$  then f has a power series expansion that converges uniformly on every compact  $K \subset \mathbb D$ .
- Show that any holomorphic function f can be uniformly approximated by polynomials.
- Show that if f is holomorphic on a connected region  $\Omega$  and  $f' \equiv 0$  on  $\Omega$ , then f is constant on  $\Omega$
- Show that if |f| = 0 on  $\partial\Omega$  then either f is constant or f has a zero in  $\Omega$ .
- Show that if  $\{f_n\}$  is a sequence of holomorphic functions converging uniformly to a function f on every compact subset of  $\Omega$ , then f is holomorphic on  $\Omega$  and  $\{f'_n\}$  converges uniformly to f' on every such compact subset.
- Show that if each  $f_n$  is holomorphic on  $\Omega$  and  $F := \sum f_n$  converges uniformly on every compact subset of  $\Omega$ , then F is holomorphic.
- Show that if f is once complex differentiable at each point of  $\Omega$ , then f is holomorphic.

# ${f 16}\,|$ Draft of Problem Book

- Prove the triangle inequality
- Prove the reverse triangle inequality
- Show that  $\sum z^{k-1}/k$  converges for all  $z \in S^1$  except z = 1.
- What is an example of a noncontinuous limit of continuous functions?
- Show that the uniform limit of continuous functions is continuous.
- Show that f is holomorphic if and only if  $\bar{\partial} f = 0$ .
- Show  $n^{\frac{1}{n}} \stackrel{n \to \infty}{\to} 1$ .
- Show that if f is holomorphic with f' = 0 on  $\Omega$  then f is constant.
- Show that holomorphic implies analytic.
- Use Cauchy's inequality to prove Liouville's theorem

Problem 16.0.1 (?)

What is a pair of conformal equivalences between  $\mathbb{H}$  and  $\mathbb{D}$ ?

**Solution:** 

$$F: HH \to \mathbb{D}$$
 
$$z \mapsto \frac{i-z}{i+z}$$

$$G: \mathbb{D} \to \mathbb{H}$$
 
$$w \mapsto i\frac{1-w}{1+w}.$$

Mnemonic: any point in  $\mathbb{H}$  is closer to i than -i, so |F(z)| < 1.

• Maps  $\mathbb{R} \to S^1 \setminus \{-1\}$ .

Problem 16.0.2 (?)

What is conformal equivalence  $\mathbb{H} \rightleftharpoons S \coloneqq \{ w \in \mathbb{C} \mid 0 < \arg(w) < \alpha \pi \}$ ?

Solution:

$$f(z) = z^{\alpha}$$
.