Algebra Qualifying Exam Review

D. Zack Garza

Table of Contents

Contents

Table of Contents 2

1 Topics and Remarks 2 12
- 1.1 General References . . . 12
- 1.2 Group Theory . . . 12
  - 1.2.1 Topics . . . 12
- 1.3 Linear Algebra . . . 14
  - 1.3.1 Topics . . . 14
- 1.4 Rings . . . 15
  - 1.4.1 Topics . . . 15
- 1.5 Modules . . . 17
  - 1.5.1 Topics . . . 17
- 1.6 Field Theory . . . 17
  - 1.6.1 Topics . . . 17

2 Group Theory 18
- 2.1 Big List of Notation . . . 18
- 2.2 Definitions . . . 19
- 2.3 Subgroups, Cosets, Quotients . . . 21
- 2.4 Cosets . . . 22
- 2.5 Special Groups . . . 23
  - 2.5.1 Cyclic Groups . . . 24
  - 2.5.2 The Symmetric Group . . . 25
- 2.6 Exercises . . . 26
- 2.7 Counting Theorems . . . 27
- 2.8 Group Actions . . . 28
- 2.9 Examples of Orbit-Stabilizer and the Class Equation . . . 30
  - 2.9.1 The Class Equation and Burnside's Lemma . . . 30
  - 2.9.2 Conjugation on Subgroups . . . 32
  - 2.9.3 Left Translation on Cosets . . . 33

3 Sylow Theorems 34
- 3.1 Sylow 1 (Cauchy for Prime Powers) . . . 34
- 3.2 Sylow 2 (Sylows are Conjugate) . . . 34
- 3.3 Sylow 3 (Numerical Constraints) . . . 35
- 3.4 Corollaries and Applications . . . 35
- 3.5 Proof of Sylow Theorems . . . 35
- 3.6 Exercises . . . 36
- 3.7 Isomorphism Theorems . . . 36
- 3.8 Products . . . 38
- 3.9 Automorphism Groups . . . 40

3.10 Special Classes of Groups . . . 40
3.11 Classification of Groups . . . 40
3.11.1 Finitely Generated Abelian Groups . . . 41
3.11.2 Classifying Abelian Groups of a Given Order . . . 42
3.12 Series of Groups . . . 43

4 Ring Theory **44**
4.1 Definitions . . . 44
4.1.1 Undergrad Review . . . 44
4.1.2 Types of Rings . . . 46
4.1.3 Commutative Algebra . . . 47
4.2 Structure Theorems . . . 47
4.3 Zorn's Lemma . . . 49
4.4 Unsorted . . . 50

5 Ring Theory **50**

6 Field Theory **50**
6.1 Facts and Definitions . . . 50
6.2 Extensions . . . 52
6.3 Finite Fields . . . 55
6.4 Cyclotomic Polynomials . . . 56
6.5 Splitting Fields . . . 58
6.6 Unsorted . . . 58

7 Galois Theory **58**
7.0.1 Lemmas About Towers . . . 59
7.0.2 Fundamental Theorem of Galois Theory . . . 61
7.0.3 Examples . . . 62
7.0.4 Counterexamples . . . 63

8 Modules **63**
8.1 General Theory . . . 64
8.2 Free and Projective Modules . . . 64
8.3 Exact Sequences . . . 66
8.4 Classification of Modules over a PID . . . 67
8.5 Algebraic Properties . . . 68

9 Linear Algebra **69**
9.1 Definitions . . . 69
9.2 Minimal / Characteristic Polynomials . . . 70
9.3 Finding Minimal Polynomials . . . 71
9.4 Canonical Forms . . . 71
9.4.1 Rational Canonical Form . . . 72
9.4.2 Jordan Canonical Form . . . 72
9.4.3 Finding Possible Canonical Forms . . . 73
9.4.4 Using Canonical Forms . . . 73
9.5 Diagonalizability . . . 74
9.6 Matrix Counterexamples . . . 75

9.7 Matrix Groups . . . 76

10 Representation Theory **77**

11 Extra Problems **78**
11.1 Commutative Algebra . . . 78
11.2 Group Theory . . . 78
11.2.1 Basic Structure . . . 78
11.2.2 Centralizing and Normalizing . . . 79
11.2.3 Primes in Group Theory . . . 80
11.2.4 p-Groups . . . 81
11.2.5 Symmetric Groups . . . 81
11.2.6 Alternating Groups . . . 83
11.2.7 Dihedral Groups . . . 83
11.2.8 Other Groups . . . 83
11.2.9 Classification . . . 83
11.2.10 Group Actions . . . 84
11.2.11 Series of Groups . . . 84
11.2.12 Misc . . . 84
11.2.13 Nonstandard Topics . . . 85
12.1 Ring Theory . . . 85
12.2 Field Theory . . . 87
12.3 Galois Theory . . . 88
12.3.1 Theory . . . 88
12.3.2 Computations . . . 89
12.4 Modules and Linear Algebra . . . 89
12.5 Linear Algebra . . . 90

13 Even More Algebra Questions **90**
13.1 Groups . . . 90
13.1.1 Question 1.1 . . . 90
13.1.2 Question 1.2 . . . 90
13.1.3 Question 1.3 . . . 90
13.1.4 Question 1.4 . . . 91
13.1.5 Question 1.5 . . . 91
13.1.6 Question 1.6. . . . 91
13.1.7 Question 1.7 . . . 91
13.1.8 Question 1.8 . . . 91
13.1.9 Question 1.9 . . . 91
13.1.10 Question 1.10 . . . 91
13.1.11 Question 1.11 . . . 92
13.1.12 Question 1.12 . . . 92
13.1.13 Question 1.13 . . . 92
13.1.14 Question 1.14 . . . 92
13.1.15 Question 1.15 . . . 92
13.1.16 Question 1.16 . . . 92
13.1.17 Question 1.17 . . . 92
13.1.18 Question 1.18 . . . 92

13.1.19 Question 1.19 . . . 93
13.1.20 Question 1.20 . . . 93
13.1.21 Question 1.21 . . . 93
13.1.22 Question 1.22 . . . 93
13.1.23 Question 1.23 . . . 93
13.1.24 Question 1.24 . . . 93
13.1.25 Question 1.25 . . . 93
13.1.26 Question 1.26. . . . 93
13.1.27 Question 1.27 . . . 94
13.1.28 Question 1.28 . . . 94
13.1.29 Question 1.29 . . . 94
13.1.30 Question 1.30 . . . 94
13.1.31 Question 1.31 . . . 94
13.1.32 Question 1.32 . . . 94
13.1.33 Question 1.33 . . . 94
13.1.34 Question 1.34 . . . 94
13.1.35 Question 1.35 . . . 95
13.1.36 Question 1.36 . . . 95
13.1.37 Question 1.37 . . . 95
13.1.38 Question 1.38 . . . 95
13.1.39 Question 1.39 . . . 95
13.1.40 Question 1.40 . . . 95
13.1.41 Question 1.41. . . . 96
13.1.42 Question 1.42 . . . 96
13.1.43 Question 1.43 . . . 96
13.1.44 Question 1.44 . . . 96
13.2 Classification of Finite groups . . . 96
13.2.1 Question 2.1 . . . 96
13.2.2 Question 2.2 . . . 96
13.2.3 Question 2.3 . . . 97
13.2.4 Question 2.4 . . . 97
13.2.5 Question 2.5 . . . 97
13.2.6 Question 2.6 . . . 97
13.2.7 Question 2.7 . . . 97
13.2.8 Question 2.8 . . . 97
13.2.9 Question 2.9 . . . 97
13.2.10 Question 2.10 . . . 97
13.2.11 Question 2.11 . . . 98
13.2.12 Question 2.12 . . . 98
13.2.13 Question 2.13 . . . 98
13.2.14 Question 2.14 . . . 98
13.2.15 Question 2.15 . . . 98
13.2.16 Question 2.16. . . . 98
13.2.17 Question 2.17 . . . 98
13.2.18 Question 2.18 . . . 98
13.2.19 Question 2.19 . . . 99
13.2.20 Question 2.20 . . . 99

13.3 Fields and Galois Theory . . . 99
13.3.1 Question 3.1 . . . 99
13.3.2 Question 3.2 . . . 99
13.3.3 Question 3.3 . . . 99
13.3.4 Question 3.4 . . . 99
13.3.5 Question 3.5 . . . 99
13.3.6 Question 3.6 . . . 100
13.3.7 Question 3.7 . . . 100
13.3.8 Question 3.8 . . . 100
13.3.9 Question 3.9 . . . 100
13.3.10 Question 3.10 . . . 100
13.3.11 Question 3.11 . . . 100
13.3.12 Question 3.12 . . . 100
13.3.13 Question 3.13. . . . 101
13.3.14 Question 3.14 . . . 101
13.3.15 Question 3.15 . . . 101
13.3.16 Question 3.16 . . . 101
13.3.17 Question 3.17 . . . 101
13.3.18 Question 3.18 . . . 101
13.3.19 Question 3.19 . . . 101
13.3.20 Question 3.20 . . . 101
13.3.21 Question 3.21 . . . 102
13.3.22 Question 3.22 . . . 102
13.3.23 Question 3.23 . . . 102
13.3.24 Question 3.24 . . . 102
13.3.25 Question 3.25 . . . 102
13.3.26 Question 3.26 . . . 102
13.3.27 Question 3.27 . . . 102
13.3.28 Question 3.28 . . . 102
13.3.29 Question 3.29 . . . 103
13.3.30 Question 3.30 . . . 103
13.3.31 Question 3.31 . . . 103
13.3.32 Question 3.32 . . . 103
13.3.33 Question 3.33 . . . 103
13.3.34 Question 3.34. . . . 103
13.3.35 Question 3.35 . . . 103
13.3.36 Question 3.36 . . . 103
13.3.37 Question 3.37 . . . 104
13.3.38 Question 3.38 . . . 104
13.3.39 Question 3.39 . . . 104
13.3.40 Question 3.40 . . . 104
13.3.41 Question 3.41 . . . 104
13.3.42 Question 3.42 . . . 104
13.3.43 Question 3.43 . . . 104
13.3.44 Question 3.44 . . . 104
13.3.45 Question 3.45 . . . 105
13.3.46 Question 3.46 . . . 105
13.3.47 Question 3.47 . . . 105

13.3.48 Question 3.48 . . . 105
13.3.49 Question 3.49 . . . 105
13.3.50 Question 3.50 . . . 105
13.3.51 Question 3.51 . . . 105
13.3.52 Question 3.52 . . . 105
13.3.53 Question 3.53 . . . 106
13.3.54 Question 3.54 . . . 106
13.3.55 Question 3.55 . . . 106
13.3.56 Question 3.56. . . . 106
13.3.57 Question 3.57 . . . 106
13.3.58 Question 3.58 . . . 106
13.3.59 Question 3.59 . . . 106
13.3.60 Question 3.60 . . . 107
13.3.61 Question 3.61 . . . 107
13.3.62 Question 3.62 . . . 107
13.3.63 Question 3.63 . . . 107
13.3.64 Question 3.64 . . . 107
13.3.65 Question 3.65 . . . 107
13.3.66 Question 3.66 . . . 107
13.3.67 Question 3.67 . . . 107
13.3.68 Question 3.68 . . . 108
13.3.69 Question 3.69 . . . 108
13.3.70 Question 3.70 . . . 108
13.3.71 Question 3.71 . . . 108
13.3.72 Question 3.72 . . . 108
13.3.73 Question 3.73 . . . 108
13.3.74 Question 3.74 . . . 108
13.3.75 Question 3.75 . . . 108
13.3.76 Question 3.76 . . . 109
13.3.77 Question 3.77 . . . 109
13.3.78 Question 3.78 . . . 109
13.4 Normal Forms . . . 109
13.4.1 Question 4.1 . . . 109
13.4.2 Question 4.2 . . . 109
13.4.3 Question 4.3 . . . 109
13.4.4 Question 4.4 . . . 109
13.4.5 Question 4.5 . . . 110
13.4.6 Question 4.6 . . . 110
13.4.7 Question 4.7 . . . 110
13.4.8 Question 4.8 . . . 110
13.4.9 Question 4.9 . . . 110
13.4.10 Question 4.10 . . . 110
13.4.11 Question 4.11 . . . 110
13.4.12 Question 4.12 . . . 110
13.4.13 Question 4.13 . . . 111
13.4.14 Question 4.14 . . . 111
13.4.15 Question 4.15 . . . 111
13.4.16 Question 4.16 . . . 111

13.4.17 Question 4.17 . . . 111
13.4.18 Question 4.18 . . . 111
13.4.19 Question 4.19 . . . 111
13.4.20 Question 4.20 . . . 111
13.4.21 Question 4.21 . . . 112
13.5 Matrices and Linear Algebra . . . 112
13.5.1 Question 5.1 . . . 112
13.5.2 Question 5.2 . . . 112
13.5.3 Question 5.3 . . . 112
13.5.4 Question 5.4 . . . 112
13.5.5 Question 5.5 . . . 112
13.5.6 Question 5.6 . . . 112
13.5.7 Question 5.7 . . . 113
13.5.8 Question 5.8 . . . 113
13.5.9 Question 5.9 . . . 113
13.5.10 Question 5.10 . . . 113
13.5.11 Question 5.11 . . . 113
13.5.12 Question 5.12 . . . 113
13.5.13 Question 5.13 . . . 113
13.5.14 Question 5.14 . . . 113
13.5.15 Question 5.15 . . . 114
13.5.16 Question 5.16 . . . 114
13.5.17 Question 5.17 . . . 114
13.5.18 Question 5.18 . . . 114
13.5.19 Question 5.19 . . . 114
13.5.20 Question 5.20 . . . 114
13.6 Rings . . . 114
13.6.1 Question 6.1 . . . 114
13.6.2 Question 6.2 . . . 115
13.6.3 Question 6.3 . . . 115
13.6.4 Question 6.4 . . . 115
13.6.5 Question 6.5 . . . 115
13.6.6 Question 6.6 . . . 115
13.6.7 Question 6.7 . . . 115
13.6.8 Question 6.8 . . . 115
13.6.9 Question 6.9 . . . 115
13.6.10 Question 6.10 . . . 116
13.6.11 Question 6.11 . . . 116
13.6.12 Question 6.12 . . . 116
13.6.13 Question 6.13 . . . 116
13.6.14 Question 6.14 . . . 116
13.6.15 Question 6.15 . . . 116
13.6.16 Question 6.16 . . . 116
13.6.17 Question 6.17 . . . 116
13.6.18 Question 6.18 . . . 117
13.6.19 Question 6.19 . . . 117
13.6.20 Question 6.20 . . . 117
13.6.21 Question 6.21 . . . 117

13.6.22 Question 6.22 . . . 117
13.6.23 Question 6.23. . . . 117
13.6.24 Question 6.24 . . . 117
13.6.25 Question 6.25 . . . 118
13.6.26 Question 6.26 . . . 118
13.6.27 Question 6.27 . . . 118
13.6.28 Question 6.28 . . . 118
13.6.29 Question 6.29 . . . 118
13.6.30 Question 6.30 . . . 118
13.6.31 Question 6.31 . . . 118
13.6.32 Question 6.32 . . . 119
13.6.33 Question 6.33 . . . 119
13.6.34 Question 6.34 . . . 119
13.6.35 Question 6.35 . . . 119
13.6.36 Question 6.36 . . . 119
13.6.37 Question 6.37 . . . 119
13.6.38 Question 6.38 . . . 119
13.6.39 Question 6.39 . . . 119
13.6.40 Question 6.40 . . . 120
13.6.41 Question 6.41 . . . 120
13.6.42 Question 6.42. . . . 120
13.6.43 Question 6.43 . . . 120
13.6.44 Question 6.44 . . . 120
13.6.45 Question 6.45 . . . 120
13.6.46 Question 6.46 . . . 120
13.6.47 Question 6.47 . . . 121
13.6.48 Question 6.48 . . . 121
13.6.49 Question 6.49 . . . 121
13.6.50 Question 6.50 . . . 121
13.6.51 Question 6.51 . . . 121
13.6.52 Question 6.52 . . . 121
13.7 Modules . . . 121
13.7.1 Question 7.1 . . . 121
13.7.2 Question 7.2 . . . 122
13.7.3 Question 7.3 . . . 122
13.7.4 Question 7.4 . . . 122
13.7.5 Question 7.5 . . . 122
13.7.6 Question 7.6. . . . 122
13.7.7 Question 7.7 . . . 122
13.7.8 Question 7.8 . . . 122
13.7.9 Question 7.9 . . . 123
13.7.10 Question 7.10 . . . 123
13.8 Representation Theory . . . 123
13.8.1 Question 8.1 . . . 123
13.8.2 Question 8.2 . . . 123
13.8.3 Question 8.3 . . . 123
13.8.4 Question 8.4 . . . 123
13.8.5 Question 8.5 . . . 123

13.8.6 Question 8.6 . . . 124
13.8.7 Question 8.7 . . . 124
13.8.8 Question 8.8 . . . 124
13.8.9 Question 8.9 . . . 124
13.8.10 Question 8.10 . . . 124
13.8.11 Question 8.11 . . . 124
13.8.12 Question 8.12 . . . 124
13.8.13 Question 8.13 . . . 124
13.8.14 Question 8.14. . . . 125
13.8.15 Question 8.15 . . . 125
13.8.16 Question 8.16 . . . 125
13.8.17 Question 8.17 . . . 125
13.8.18 Question 8.18 . . . 125
13.8.19 Question 8.19 . . . 125
13.8.20 Question 8.20 . . . 126
13.8.21 Question 8.21 . . . 126
13.8.22 Question 8.22 . . . 126
13.8.23 Question 8.23 . . . 126
13.8.24 Question 8.24 . . . 126
13.8.25 Question 8.25 . . . 126
13.8.26 Question 8.26 . . . 126
13.8.27 Question 8.27 . . . 126
13.8.28 Question 8.28 . . . 127
13.8.29 Question 8.29 . . . 127
13.8.30 Question 8.30 . . . 127
13.8.31 Question 8.31. . . . 127
13.8.32 Question 8.32 . . . 127
13.8.33 Question 8.33 . . . 127
13.8.34 Question 8.34 . . . 128
13.8.35 Question 8.35 . . . 128
13.8.36 Question 8.36 . . . 128
13.8.37 Question 8.37 . . . 128
13.9 Categories and Functors . . . 128
13.9.1 Question 9.1 . . . 128
13.9.2 Question 9.2 . . . 128
13.9.3 Question 9.3 . . . 128

14 Appendix: Extra Topics **129**
14.1 Characteristic Subgroups . . . 129
14.2 Normal Closures and Cores . . . 130
14.2.1 Exercises . . . 130
14.3 Nilpotent Groups . . . 131

15 UGA Fall 2019 Problem Sets **132**
15.1 Problem Set One . . . 132
15.1.1 Exercises . . . 132
15.1.2 Qual Problems . . . 133

15.2 Problem Set Two . . . 138
15.2.1 Exercises . . . 138
15.2.2 Qual Problems . . . 139
15.3 Problem Set Three . . . 140
15.3.1 Exercises . . . 140
15.3.2 Qual Problems . . . 140
15.4 Problem Set Four . . . 141
15.4.1 Exercises . . . 141
15.4.2 Qual Problems . . . 142
15.5 Problem Set Five . . . 143
15.5.1 Exercises . . . 143
15.5.2 Qual Problems . . . 144
15.6 Problem Set Six . . . 145
15.6.1 Exercises . . . 145
15.6.2 Qual Problems . . . 145
15.7 Problem Set Seven . . . 146
15.7.1 Exercises . . . 146
15.7.2 Qual Problems . . . 148
15.8 Problem Set Eight . . . 149
15.8.1 Exercises . . . 149
15.8.2 Qual Problems . . . 150
15.9 Problem Set Nine . . . 151
15.9.1 Exercises . . . 151
15.9.2 Qual Problems . . . 152
15.10 Problem Set Ten . . . 153
15.10.1 Exercises . . . 153
15.10.2 Qual Problems . . . 154

Bibliography **155**

1 Topics and Remarks 2

Remark 1.0.1: Adapted from remark written by Roy Smith, August 2006:

As a general rule, students are responsible for knowing both the theory (proofs) and practical applications (e.g. **how to find the Jordan or rational canonical form** of a given matrix, **or the Galois group of a given polynomial**) of the topics mentioned.

1.1 General References

- David Dummit and Richard Foote, Abstract Algebra, Wiley, 2003. [1]
- Kenneth Hoffman and Ray Kunze, Linear Algebra, Prentice-Hall, 1971. [2]
- Thomas W. Hungerford, Algebra, Springer, 1974. [3]
- Roy Smith, Algebra Course Notes (843-1 through 845-3). [4]
  - Note: scroll down the page to find links to his course notes.

1.2 Group Theory

References: [1], [3], [4] "The first 6 chapters (220 pages) of Dummit and Foote are excellent. All the definitions and proofs of these theorems on groups are given in Smith's web based lecture notes for math 843 part 1."

1.2.1 Topics

Chapters 1-9 of Dummit and Foote

- **The first isomorphism theorem**,
- Fundamental theorem of finite abelian groups
- Left and right cosets
- Normalizer

- Lagrange's theorem
- Isomorphism theorems
- Lagrange's Theorem
- Group generated by a subset
- Subgroups and quotient groups
- Fundamental homomorphism theorems
- Direct and semi-direct products
  - Recognition of internal direct product
  - Recognition of semi-direct product
- Composite groups
- Structures of special types of groups such as:
  - p-groups
  - Dihedral,
    - ♢ Cyclic groups
    - ♢ Free groups
      - ♢ Generators and relations
  - Symmetric and Alternating groups
    - ♢ Cycle decompositions
- Group actions with applications to the structure of groups such as
  - **The Sylow Theorems**
    - ♢ Proof of Sylow theorems
  - Orbit stabilizer theorem
  - Orbits act on left cosets of subgroups
  - Action of G on itself by conjugation
  - Class equation
  - Cayley's theorem
- The simple groups of order between 60 and 168 have prime order
- The simplicity of A_n, for $n \geq 5$
- Solvable groups
- Subgroups of index p, the smallest prime dividing $\#G$, are normal

- p-groups
- p^2 groups are abelian
- Automorphisms
  - Inner automorphisms
- A_n is simple for $n \geq 5$
- Classification of groups of order pq
- Commutator subgroup
- Nilpotent groups
- Upper central series
- Lower central series
- Derived series
- Solvable groups
- Fratini's argument
- The Jordan Holder theorem

*The proof of Jordan-Holder is seldom tested on the qual**, but proofs are always of interest.*

1.3 Linear Algebra

References: [1],[2],[4]

1.3.1 Topics

- Determinants
- Eigenvalues and eigenvectors
- Cayley-Hamilton Theorem
- Canonical forms for matrices
- Linear groups ($\mathrm{GL}_n, \mathrm{SL}_n, \mathrm{O}_n, \mathrm{U}_n$)
- Duality

  - Dual spaces,
  - Dual bases,
  - Induced dual map,
  - Double duals

- Finite-dimensional spectral theorem

1.4 Rings

References: [1],[3],[4]

- DF chapters 13,14 (about 145 pages).

- Smith:

  - 843-2, sections 11,12, and 16-21 (39 pages)
  - 844-1, sections 7-9 (20 pages)
  - 844-2, sections 10-16, (37 pages)

- DF Chapters 7, 8, 9.

1.4.1 Topics

- Properties of ideals and quotient rings

- The fundamental isomorphism theorems for rings

- I maximal iff R/I is a field

- Zorn's lemma

  - Every vector space has a basis
  - Maximal ideals exist
  - Construct algebraic field closures
  - Why it is unnecessary in countable or noetherian rings.

Smith discusses extensively in 844-1.

- Chinese Remainder Theorem

- Euclidean algorithm

- Primes and irreducibles

- Gaussian integers
- Localization of a domain
- Field of fractions
- Factorization in domains
- Factorization in $Z[i]$
- Characterizations and properties of special rings such as:
  - Euclidean $\implies$ PID $\implies$ UFD
  - Domains
    - ♢ Primes are irreducible
  - UFDs
    - ♢ Have GCDs
    - ♢ Sometimes PIDs
  - PIDs
    - ♢ Noetherian
    - ♢ Irreducibles are prime
    - ♢ Are UFDs
    - ♢ Have GCDs
    - ♢ Results about PIDs *(DF Section 8.2)*
      - ♢ Example of a PID that is not a Euclidean domain *(DF p.277)*
      - ♢ Proof that a Euclidean domain is a PID and hence a UFD
      - ♢ Proof that $\mathbb{Z}$ and $k[x]$ are UFDs *(p.289 Smith, p.300 DF)*
      - ♢ A polynomial ring in infinitely many variables over a UFD is still a UFD *(Easy, DF, p.305)*
  - Euclidean domains
    - ♢ Are PIDs
- Gauss's important theorem on unique factorization of polynomials:
  - $\mathbb{Z}[x]$ is a UFD
  - $R[x]$ is a UFD when R is a UFD
- Polynomial rings
- Polynomials
  - Gauss' lemma
  - Remainder and factor theorem
  - Eisenstein's criterion *(DF p.309)* > Stated only for monic polynomials – proof of general case identical. > See Smith's notes for the full version.

  - Reducibility
  - Rational root test

- Cyclic product structure of $(\mathbb{Z}/n\mathbb{Z})^\times$

Exercise in DF, Smith 844-2, section 18

- Gröbner bases and division algorithms for polynomials in several variables *(DF 9.6.)*

1.5 Modules

References: [1],[3],[4]

1.5.1 Topics

- Fundamental homomorphism theorems for rings and modules
- Applications to the structure of:
  - Finitely generated abelian groups
  - Canonical forms of matrices
- Classification of finitely generated modules over PIDs *(with emphasis on Euclidean Domains)*
- Modules over PIDs and canonical forms of matrices. *DF sections 10.1, 10.2, 10.3, and 12.1, 12.2, 12.3.*
  - Constructive proof of decomposition: DF Exercises 12.1.16-19

Smith 845-1 and 845-2: Detailed discussion of the constructive proof.

1.6 Field Theory

1.6.1 Topics

References: [1],[3],[4]

- Algebraic extensions of fields

- Properties of finite fields
- Separable extensions
- Fundamental theorem of Galois theory
- Computations of Galois groups
  - of polynomials of small degree
  - of cyclotomic polynomials
- Solvability of polynomials by radicals

2 | Group Theory

2.1 Big List of Notation

Notation	Definition
$C_G(x)$	Centralizer of an element $\coloneqq \left\{ g \in \Gamma \;\middle|\; [g,x] = 1 \right\} \subseteq \Gamma$
$C_G(H)$	Centralizer of an subgroup $\coloneqq \left\{ g \in \Gamma \;\middle|\; [g,x] = 1 \ \forall h \in H \right\} = \bigcap_{h \in H} C_H(h) \subseteq G$
$C(H)$	Conjugacy Class $\coloneqq \left\{ ghg^{-1} \;\middle|\; g \in G \right\} \leq G \subseteq G$
$Z(G)$	Center $\coloneqq \left\{ x \in G \;\middle|\; \forall g \in G,\, gxg^{-1} = x \right\} \subseteq G$
$N_G(H)$	Normalizer $\coloneqq \left\{ g \in G \;\middle|\; gHg^{-1} = H \right\} \subseteq G$
$\mathrm{Inn}(G)$	Inner Automorphisms $\coloneqq \left\{ \varphi_g(x) \coloneqq gxg^{-1} \right\} \subseteq \mathrm{Aut}(G)$
$\mathrm{Out}(G)$	Outer Automorphisms $\mathrm{Aut}(G)/\mathrm{Inn}(G) \hookleftarrow \mathrm{Aut}(G)$
$[gh]$	Commutator of Elements $\coloneqq ghg^{-1} \in G$
$[GH]$	Commutator of Subgroups $\coloneqq \left\langle \left\{ [gh] \;\middle|\; g \in G,\, h \in H \right\} \right\rangle \leq G$
$\mathcal{O}_x$, Gx	Orbit of an Element $\coloneqq \left\{ gx \;\middle|\; x \in X \right\}$

Notation	Definition
$\mathrm{Stab}_G(x)$, G_x	Stabilizer of an Element $:= \left\{ g \in G \;\middle|\; gx = x \right\} \subseteq G$
X/G	Set of Orbits $:= \left\{ G_x \;\middle|\; x \in X \right\} \subseteq 2^X$
X^g	Fixed Points $\left\{ x \in X \;\middle|\; \forall g \in G,\, gx = x \right\} \subseteq X$
2^X	The powerset of X $:= \{U \subseteq X\}$

- For any p dividing the order of G, $\mathrm{Syl}_p(G)$ denotes the *set* of Sylow-p subgroups of G.

2.2 Definitions

Fact 2.2.1
An set morphism that is *either* injective or surjective between sets of the same size is automatically a bijection. Consequently, a group morphism between groups of the same size that is either injective or surjective is automatically an isomorphism.

Fact 2.2.2 (The division algorithm)
If $a, b \in \mathbb{Z}$ with $\gcd(a, b) = d$, then there exist $s, t \in \mathbb{Z}$ such that

$$as + bt = d.$$

Remark 2.2.3: Useful context clue! In particular, this works when a, b are coprime and $d = 1$. If you see "coprime" in a finite group question, try the division algorithm.

Definition 2.2.4 (Subgroup Generated by a Subset)
If $H \subset G$, then $\langle H \rangle$ is the smallest subgroup containing H:

$$\langle H \rangle = \cap \left\{ H \;\middle|\; H \subseteq M \leq G \right\} M = \left\{ h_1^{\pm 1} \cdots h_n^{\pm 1} \;\middle|\; n \geq 0, h_i \in H \right\}$$

where adjacent h_i are distinct.

Definition 2.2.5 (Conjugacy class)
The **conjugacy class** of h is defined as

$$C(h) := \left\{ ghg^{-1} \;\middle|\; g \in G \right\}.$$

Remark 2.2.6: $[e] = \{e\}$ is always in a conjugacy class of size one – this is useful for counting and divisibility arguments. Conjugacy classes are **not** subgroups in general, since they don't generally contain e. However, by orbit-stabilizer and the conjugation action, their sizes always divide the order of G.

Also note that$[x] = \{x\} \iff x \in Z(G)$, i.e. having a trivial conjugacy class is the same as being central.

Definition 2.2.7 (Conjugate subgroups)
Two subgroups $H, K \leq G$ are **conjugate** iff there exists some $g \in G$ such that $gHg^{-1} = K$. Note that all conjugates have the same cardinality.

Definition 2.2.8 (Normal subgroup)
A subgroup $N \leq G$ is **normal** iff $gH = Hg$ for every $g \in G$, or equivalently $gHg^{-1} = H$ for all g, so H has only itself as a conjugate. We denote this by $N \trianglelefteq G$. Equivalently, for every inner automorphism $\psi \in \text{Inn}(G)$, $\psi(N) = N$.

Proposition 2.2.9 ***(Normal iff disjoint union of conjugacy classes).***
$N \trianglelefteq G \iff N = \coprod'[h_i]$ is a disjoint union of conjugacy classes, where the index set for this union is one h_i from each conjugacy class.

Proof (?).
Note that $C(h_i) = \left\{ gh_ig^{-1} \;\middle|\; g \in G \right\}$, and $gh_ig^{-1} \in H$ since H is normal, so $C(h_i) \subseteq G$ for all i. Conversely, if $C(h_i) \subseteq H$ for all $h_i \in H$, then $gh_ig^{-1} \in H$ for all i and H is normal. ■

Definition 2.2.10 (Centralizer)

$$C_G(H) = \left\{ g \in G \;\middle|\; ghg^{-1} = h \;\forall h \in H \right\}$$

These are elements that fix H pointwise under the conjugation action.

Definition 2.2.11 (Normalizer)

$$N_G(H) = \left\{ g \in G \;\middle|\; gHg^{-1} = H \right\} = \cup \left\{ H \;\middle|\; H \trianglelefteq M \leq G \right\} M$$

Contrast to the centralizer: these don't have to fix H pointwise.

Remark 2.2.12: $C_G(S) \trianglelefteq N_G(H)$ for any H.

Definition 2.2.13 (Commutator)
The **commutator subgroup** of G is denoted $[G, G] \leq G$. It is the subgroup generated by all

elementary commutators:

$$[G,G] \coloneqq \left\langle aba^{-1}b^{-1} \;\middle|\; a,b \in G \right\rangle.$$

It is the smallest normal subgroup $N \trianglelefteq G$ such that G/N is abelian, so if $H \leq G$ and G/H is abelian, $H \subseteq [G,G]$.

Definition 2.2.14 (Group Presentation)
An expression of the form $G = \left\langle S \;\middle|\; R \right\rangle$ where S is a set of elements and R a set of words defining relations means that $G \coloneqq F[S]/\mathrm{cl}_n(R)$ where $F[S]$ is the free group on the set S and $\mathrm{cl}_n(R)$ is the normal closure, the smallest normal subgroup of $F[S]$ containing R.

Remark 2.2.15: Finding morphisms between presentations: if G is presented with generators g_i with relations r_i and H is any group containing elements h_i also satisfying r_i, there is a group morphism

$$\begin{aligned} \varphi : G &\to H \\ g_i &\mapsto h_i \quad \forall i. \end{aligned}$$

Why this exists: the presentation yields a morphism $\pi : F(g_i) \to G$ with $G \cong F(g_i)/\ker\pi$. Define a map $\psi : F(g_i) \to H$ where $g_i \mapsto h_i$, then since the h_i satisfy the relations r_i, $\ker\pi \subseteq \ker\psi$. So ψ factors through $\ker\pi$ yielding a morphism $F/\ker\pi \to H$.

Proposition 2.2.16 ***(NC Theorem).***
$N_G(H)/C_G(H)$ is isomorphic to a subgroup of $\mathrm{Aut}(H)$.

2.3 Subgroups, Cosets, Quotients

Definition 2.3.1 (Subgroup)
A subset $H \subseteq G$ is a **subgroup** iff

1. Closure: $HH \subset H$
2. Identity: $e \in H$
3. Inverses: $g \in H \iff g^{-1} \in H$.

Proposition 2.3.2 ***(One-step subgroup test).***
If $H \subseteq G$ and $a,b \in H \implies ab^{-1} \in H$, then $H \leq G$.

Proof (of the one-step subgroup test).

- Identity: $a = b = x \implies xx^{-1} = e \in H$
- Inverses: $a = e, b = x \implies x^{-1} \in H$.
- Closure: let $x,y \in H$, then $y^{-1} \in H$ by above, so $xy = x(y^{-1})^{-1} \in H$.

■

Fact 2.3.3
Coprime order subgroups are disjoint, or more generally $\mathbb{Z}_p, \mathbb{Z}_q \subset G \implies \mathbb{Z}_p \cap \mathbb{Z}_q = \mathbb{Z}_{(p,q)}$.

Proposition 2.3.4 ***(Intersection of subgroups is a subgroup).***
If $H, K \leq G$ then $J \coloneqq H \cap K \leq G$ is a subgroup. Moreover $J \leq H$ and $J \leq K$.

Proof (?).
One-step subgroup test.

■

Proposition 2.3.5 ***(Tower law for subgroups).***

$$K \leq H \leq G \implies [G : K] = [G : H][H : K].$$

Proposition 2.3.6 ***(Indices grow).***
If $H, K \leq G$, then

$$[H : H \cap K] \leq [G : K].$$

Proof (?).
Write $G/H \cap K \coloneqq G/J = \{h_1 J, \cdots, h_m J\}$ as distinct cosets. Then $i \neq j \implies h_i h_j^{-1} \notin H \cap K$, but $h_i h_j^{-1} \in H$ since $H \leq G$, which forces $h_i h_j^{-1} \notin K$. So $h_i K \neq h_j K$, meaning there are at least m cosets in G/K.

■

2.4 Cosets

Proposition 2.4.1 ***(Cosets are identical or disjoint).***
Any two cosets xH, yH are either identical or disjoint.

Proof (?).
Note $x \in xH$, since $e \in H$ because H is a subgroup and we can take $h = e$ to get $x = xe \coloneqq xh \in xH$. The reverse containment is clear, so $G = \cup_{x \in G} xH$ is a union of its cosets. Suppose toward a contradiction that $\ell \in xH \cap yH$ we'll show $xH = yH$. Write $\ell = xh_1 = yh_2$ for some h_i, then

$$\begin{aligned} xh_1 = yh_2 &\implies x = yh_2 h_1^{-1} \\ xh_3 \in xH &\implies xh_3 = (yh_2 h_1^{-1})h_3 \in yH, \end{aligned}$$

so $xH \subseteq yH$. A symmetric argument shows $y_H \subseteq xH$. [a]

■

[a] See full argument: D&F p.80.

Theorem 2.4.2 ***(The Fundamental Theorem of Cosets).***

$$aH = bH \iff a^{-1}b \in H \iff b^{-1}a \in H.$$

Proof (?).
[a]

$$aH = bH \iff a \in bH \iff a = bh \text{ for some } h \iff b^{-1}a = h \iff ba^{-1} \in H.$$

■

[a] See full argument: D&F p.80.

Definition 2.4.3 (Index of a subgroup)
The **index** $[G : H]$ of a subgroup $H \leq G$ is the number of left (or right) cosets gH.

Remark 2.4.4 ***(Common coset trick):*** If you can reduce a problem to showing $X \subseteq H$, it suffices to show $xH = H$ for all $x \in X$.

Remark 2.4.5: Cosets form an equivalence relation and thus partition a group. Nice trick: write $G/H = \{g_1H, g_2H, \cdots, g_nH\}$, then $G = \coprod_{i \leq n} g_iH$.

Theorem 2.4.6 ***(Counting Cosets).***
If $H \trianglelefteq G$, then

$$[G : H] = |G/H| = \frac{|G|}{|H|}.$$

2.5 Special Groups

Definition 2.5.1 (The Dihedral Group)
A **dihedral group** of order $2n$ is given by

$$D_n = \left\langle r, s \;\middle|\; r^n, s^2, rsr^{-1} = s^{-1} \right\rangle = \left\langle r, s \;\middle|\; r^n, s^2, (rs)^2 \right\rangle$$

Definition 2.5.2 (The Quaternion Group)

The **Quaternion group** of order 8 is given by

$$\begin{aligned} Q &= \left\langle x, y, z \;\middle|\; x^2 = y^2 = z^2 = xyz = -1 \right\rangle \\ &= \left\langle x, y \;\middle|\; x^4 = y^4, x^2 = y^2, yxy^{-1} = x^{-1} \right\rangle \end{aligned}$$

Mnemonic: multiply clockwise to preserve sign, counter-clockwise to negate sign. Everything squares to -1, and the triple product is -1:



Link to Diagram

Definition 2.5.3 (Transitive Subgroup)
A subgroup of S_n is **transitive** iff its action on $\{1, 2, \cdots, n\}$ is transitive.

2.5.1 Cyclic Groups

Theorem 2.5.4 ***(Subgroups of Cyclic Groups).***
G is cyclic of order $n := \#G$ iff G has a unique subgroup of order d for each d dividing n.

Proof (?).
$\Longleftarrow$: Use that $\sum_{d|n} \varphi(d) = n$, and that there are at most $\varphi(d)$ elements of order d, forcing equality.
$\Longrightarrow$: If $G = \langle a \rangle$ with $a^n = e$, then for each $d \mid n$ take $H_d := \left\langle a^{\frac{n}{d}} \right\rangle$ for existence.
■

2.5.2 The Symmetric Group

Definition 2.5.5 (The symmetric group)
The transposition presentation:

$$S_n \coloneqq \left\langle \sigma_1, \cdots, \sigma_{n-1} \;\middle|\; \sigma_i^2, [\sigma_i, \sigma_j]\,(j \neq i+1), \sigma_i\sigma_{i+1}\sigma_i = \sigma_{i+1}\sigma_i\sigma_{i+1} \right\rangle.$$

Definition 2.5.6 (The sign homomorphism)
Defined by the map

$$\begin{aligned} \operatorname{sgn} : S_n &\to (\mathbb{Z}/2, +) \\ \prod_{i \leq 2k} (a_i b_i) &\mapsto 0 \\ \prod_{i \leq 2k+1} (a_i b_i) &\mapsto 1. \end{aligned}$$

- The kernel is the alternating group, cycles that
  - **Even** cycles
  - For a single cycle: has **odd** length
  - Have an **even** number of even length cycles.
  - Can be written as an **even** number of transpositions
- The fiber over 1 is everything else:
  - **Odd** cycles
  - For a single cycle: has **even** length
  - Have an **odd** number of even length cycles.
  - Can be written as an **odd** number of transpositions

Mnemonic: the cycle parity of a k-cycle is the integer parity of $k-1$.

Definition 2.5.7 (Alternating Group)
The **alternating group** is the subgroup of **even** permutations, i.e.

$$A_n \coloneqq \left\{ \sigma \in S_n \;\middle|\; \operatorname{sgn}(\sigma) = 0 \right\}$$

Proposition 2.5.8 ***(A_n is generated by 3-cycles).***
For $n \geq 3$, A_n is generated by 3-cycles.

Proof (?).
Every 3-cycle (abc) is even, and thus in A_n. Given an arbitrary even permutation $(t_1 \dots t_{2k})$, it decomposes into a product of an odd number of transpositions $(t_{2j-1}t_{2j})$. So it suffices to write every such transposition as a 3-cycle. There are only 3 cases the occur:

- $(ab)(ab) = ()$

- $(ab)(ac) = (abc)$
- $(ab)(cd) = (abc)(adc)$.

■

Example 2.5.9 ***(Of an explicit alternating group):***

$$\begin{aligned} A_4 = \{&\mathrm{id}, \\ &(1,3)(2,4), (1,2)(3,4), (1,4)(2,3), \\ &(1,2,3), (1,3,2), \\ &(1,2,4), (1,4,2), \\ &(1,3,4), (1,4,3), \\ &(2,3,4), (2,4,3)\} \end{aligned}$$

Fact 2.5.10 (Some useful facts)

- $\sigma \circ (a_1 \cdots a_k) \circ \sigma^{-1} = (\sigma(a_1), \cdots \sigma(a_k))$
- Conjugacy classes are determined by cycle type
- The order of a cycle is its length.
- The order of an element is the least common multiple of the sizes of its disjoint cycles.
- Disjoint cycles commute.
- $A_{n \geq 5}$ is *simple*.

2.6 Exercises

- Show that normal groups absorb conjugacy classes: if $N \trianglelefteq G$ and $[g_i]$ is a conjugacy class in g, either $[g_i] \subseteq N$ or $[g_i] \cap N = \emptyset$.

- Prove that the size of a conjugacy class of g_i is the index of its centralizer, $[G : Z(g_i)] \coloneqq [G : C_G(g_i)]$.

- Prove that if G is a p-group, every subgroup $N \trianglelefteq G$ intersects the center $Z(G)$.

- Show that if G is a finite group acting transitively on a set X with at least two elements, then there exists $g \in G$ which fixes no point of X.

- Let p be prime. For each abelian group K of order p^2, how many subgroups $H \leq \mathbb{Z}^{\times 3}$ are there with $\mathbb{Z}^3/H \cong K$?

- Let $\#G = pq$, with p, q distinct primes. Show that G has a nontrivial proper normal subgroup, and if $p \not\equiv 1 \pmod{q}$ and $q \not\equiv 1 \pmod{p}$ then G is abelian.

- Let G be a finite group and let p be the smallest prime dividing $\#G$, and assume G has a normal subgroup of order p. Show that $H \subset Z(G)$.

- Let G be finite and P a Sylow 2-subgroup. Assume P is cyclic and generated by x. Show that the sign of the permutation of G corresponding to $x \mapsto gx$ is 1, and deduce that G has a nontrivial quotient of order 2.

2.7 Counting Theorems

Theorem 2.7.1 ***(Lagrange's Theorem).***

$$H \leq G \implies \#H \mid \#G.$$

Moreover, there is an equality $[G : H] = \#G/\#H$ when G is finite.

Proof (of Lagrange's theorem).
Write $G/H = \{g_0 H, g_1 H, \cdots, g_N H\}$ for some $N \coloneqq [G : H]$. Since cosets are equal or disjoint and have equal cardinality,

$$G = \coprod_{k \leq N} g_k H \implies \#G = \sum_{k \leq N} \#(g_k H) = \sum_{k \leq N} \#H = N\#H,$$

so $\#G = N\#H$, $\#H$ divides $\#G$ and $N = [G : H]$ divides $\#G$.

■

Corollary 2.7.2 ***(?).***

$$\#G = \#(G/H)\#H \coloneqq [G : H]\,\#H,$$

or written another way,

$$\#(G/H) = \#G/\#H.$$

Corollary 2.7.3.
The order of every element divides the size of G, i.e.

$$g \in G \implies o(g) \mid o(G) \implies g^{|G|} = e.$$

⚠ **Warning 2.7.4**
There do **not** necessarily exist $H \leq G$ with $|H| = n$ for every $n \mid |G|$. Counterexample: take $G = A_5$, then $\#G = 5!/2 = 60$ but G has no subgroup of order 30. If it did, this would be index 2 and thus normal, but $A_{n \geq 5}$ is simple.

Another direct counterexample: $|A_4| = 12$ but has no subgroup of order 6. If such an H existed, it can't contain every 3-cycle, since A_4 is generated by 3-cycles. For x any 3-cycle *not* in H, use that $\#A_4/H = 2$ and consider H, xH, x^2H. $x \notin H$, so $H \neq xH$, but two must be equal:

- $x^2H = H$: use $x^2 = x^{-1}$ since $x^3 = e$, but $x \in H \implies x^{-1} \in H$, ↯
- $xH = x^2H$: the fundamental theorem of cosets forces $x^{-1}x^2 \in H$, so $x \in H$. ↯

Theorem 2.7.5 ***(Cauchy's Theorem).***
For every prime p dividing $|G|$. there is an element (and thus a subgroup) of order p.

This is a partial converse to Lagrange's theorem, and strengthened by Sylow's theorem.

Proof (?).
See https://kconrad.math.uconn.edu/blurbs/grouptheory/cauchypf.pdf.
■

2.8 Group Actions

Definition 2.8.1 (Group Action)
An action of G on X is a group morphism

$$\begin{aligned} \varphi : G \times X &\to X \\ (g, x) &\mapsto gx \end{aligned}$$

or equivalently

$$\begin{aligned} \varphi : G &\to \mathrm{Aut}(X) \\ g &\mapsto (x \mapsto \varphi_g(x) \coloneqq g \cdot x) \end{aligned}$$

satisfying

1. $e \cdot x = x$
2. $g \cdot (h \cdot x) = (gh) \cdot x$

Fact 2.8.2
For any group action, the kernel is the intersection of all stabilizers:

$$\ker \psi = \bigcap_{x \in X} G_x.$$

Definition 2.8.3 (Transitive Group Action)
A group action $G \curvearrowright X$ is **transitive** iff for all $x, y \in X$ there exists a $g \in G$ such that $g \cdot x = x$.
Equivalently, the action has a single orbit.

Proposition 2.8.4 ***(Orbit Stabilizer Isomorphism).***
If $G \curvearrowright X$ transitively, then for any choice of $x \in X$ there is an isomorphism of sets given by

$$\begin{aligned} \Phi : G/G_x &\xrightarrow{\sim} X \\ gG_x &\mapsto g \curvearrowright x. \end{aligned}$$

Proof (?).

- Injectivity: $\Phi(gG_x) = \Phi(hG_x) \iff g \curvearrowright x = h \curvearrowright x \iff gh^{-1} \curvearrowright x = x \iff gh^{-1} \in G_x \iff gG_x = hG_x$.
- Well-definedness: use $gG_x = hG_x \iff gh^{-1} \in G_x \iff g^{-1}h \curvearrowright x = x$. Then $g(g^{-1}h) \curvearrowright x = g \curvearrowright x$ on one hand, and on the other $(gg^{-1})h \curvearrowright x = h \curvearrowright x$, so

$$\Phi(hG_x) \coloneqq h \curvearrowright x = (gg^{-1})h \curvearrowright x = g(g^{-1}h) \curvearrowright x = g \curvearrowright x = \Phi(gG_x).$$

- Surjectivity: equivalent to the action being transitive.

■

Proposition 2.8.5 ***(?).***
If $X \in G\text{-}\mathsf{Set}$ where $G \curvearrowright X$ transitively, then for any points $x_i \in X$, the stabilizers G_{x_0} and G_{x_1} are conjugate.

Prove

Proof (?).

- Injectivity: check that $\varphi(\bar{g}) = \varphi(\bar{h}) \iff g \curvearrowright x_0 = h \curvearrowright x_0 \iff gh^{-1} \in G_{x_0}$.
- Surjectivity: follows from transitivity.

■

Remark 2.8.6 ***(Reminder of notation):*** For a group G acting on a set X,

Notation	Definition
$\mathcal{O}(x) = Gx = \left\{ g \cdot x \;\middle|\; g \in G \right\} \subseteq X$	Orbit
$\mathrm{Stab}(x) = G_x = \left\{ g \in G \;\middle|\; g \cdot x = x \right\} \leq G$	Stabilizer
$X/G \subseteq 2^X$	Set of Orbits
$\mathrm{Fix}(G) = X^G = \left\{ x \in X \;\middle|\; g \cdot x = x \, \forall g \in G \right\} \subseteq X$	Set of Fixed Points

Note that being in the same orbit is an equivalence relation which partitions X, and G acts transitively if restricted to any single orbit.

Theorem 2.8.7 ***(Orbit-Stabilizer).***

$$\#Gx = [G : G_x] = \#G/\#G_x \quad \text{if } G \text{ is finite.}$$

Mnemonic: $G/G_x \cong Gx$.

2.9 Examples of Orbit-Stabilizer and the Class Equation

Example 2.9.1 ***(Trivial):*** Let G act on itself by left translation, where $g \mapsto (h \mapsto gh)$.

- The orbit $\mathcal{O}(x) = Gx = G$ is the entire group.
  - This action is transitive.
- The set of fixed points $\text{Fix}(G) = \left\{g \in G \;\middle|\; gx = x \,\forall x \in G\right\} = \{e\}$ is just the identity.
- The stabilizer $G_x = \left\{g \in G \;\middle|\; gx = x\right\} = \{e\}$ is just the identity.
- The kernel is the identity.

2.9.1 The Class Equation and Burnside's Lemma

Example 2.9.2 ***(Conjugation yields centers/centralizers):*** Let G act on *itself* by conjugation.

- The orbit $\mathcal{O}(g) = C(g)$ is the **conjugacy class** of x.
  - Thus the action is transitive iff there is one conjugacy class.
- The set of fixed points $\text{Fix}(G) = Z(G)$ is the **center**.
- The stabilizer is $\text{Stab}(g) = Z(g) = C_G(g)$, the **centralizer** of g.
- The kernel is the intersection of all centralizers, i.e. again the **center** $Z(G)$.

Remark 2.9.3: Note that $[G : C_G(x_i)]$ is the number of elements in the conjugacy class of x_i, and each $x_i \in Z(G)$ has a singleton conjugacy class.

Corollary 2.9.4.
Directly interpreting this using the orbit-stabilizer formula, the size of a conjugacy class $C(x)$ is the index of its centralizer, $[G : C_G(x)]$, i.e.

$$\#C(x) = [G : Z(x)].$$

Corollary 2.9.5 ***(The Class Equation).***

$$|G| = |Z(G)| + \sum_{\substack{\text{One } g \text{ from} \\ \text{each nontrivial} \\ \text{conj. class}}} [G : Z(g)]$$

Proof (of the class equation).

G is a disjoint union of its conjugacy classes, so $G = \coprod_{g\in G}{}' C(g)$ where $\coprod'$ denotes taking one representative from each conjugacy class. Thus

$$\#G = \sum_{g\in G}{}' \#C(g) = \sum_{g\in G}{}' [G : Z(g)].$$

Elements $g \in Z(g)$ in the center satisfy $Z(g) = \{e\}$ and $[G : Z(g)] = 1$ since $Z(g) = G$, so pulling these out of the sum yields

$$\begin{aligned}
\#G &= \sum_{\substack{g\in G' \\ \#[G:Z(g)]=1}}{}' [G : Z(g)] + \sum_{\substack{g\in G \\ [G:Z(g)]>1}}{}' [G : Z(g)] \\
&= \sum_{\substack{g\in G' \\ \#[G:Z(g)]=1}}{}' 1 + \sum_{\substack{g\in G \\ [G:Z(g)]>1}}{}' [G : Z(g)] \\
&= \#Z(G) + \sum_{\substack{g\in G \\ [G:Z(g)]>1}}{}' [G : Z(g)].
\end{aligned}$$

■

Proposition 2.9.6 ***(Application of the Class Equation).***
If G is simple, $H < G$ proper, and $[G : H] = n$, then there exists an injective map $\varphi : G \hookrightarrow S_n$.

Proof .
This action induces φ; it is nontrivial since $gH = H$ for all g implies $H = G$; $\ker\varphi \trianglelefteq G$ and G simple implies $\ker\varphi = 1$.

■

Corollary 2.9.7 ***(Burnside's Lemma).***
For G a finite group acting on X,

$$\#X/G = \frac{1}{\#G}\sum_{g\in G} \#X^g$$

Mnemonic: the number of orbits is equal to the average number of fixed points, i.e.

Proof (of Burnside's Lemma).
Strategy: form the set $A \coloneqq \left\{(g,x) \in G \times X \;\middle|\; g \curvearrowright x = x\right\}$ and write/count it in two different ways. First union over G:

$$A = \coprod_{g\in G} \left\{(g,x) \;\middle|\; gx = x\right\} \cong \coprod_{g\in G} \{g\} \times X^g.$$

Then union over X:

$$A = \coprod_{x\in X} \left\{(g,x) \;\middle|\; gx = x\right\} \cong \coprod_{x\in X} G_x \times \{g\}.$$

Taking cardinalities, and using the fact that $\{p\} \times A \cong A$ as sets for any set A,

$$\coprod_{g\in G} X^g \cong \coprod_{x\in X} G_x \implies \sum_{g\in G} \#X^g = \sum_{x\in X} \#G_x.$$

Apply orbit-stabilizer:

$$\begin{aligned}
\#G_x = \frac{\#G}{\#Gx} \implies \sum_{g\in G} X^g &= \sum_{x\in X} \#G_x \\
&= \sum_{x\in X} \frac{\#G}{\#Gx} \\
&= \#G \sum_{x\in X} \frac{1}{\#Gx},
\end{aligned}$$

so it suffices to show $\displaystyle\sum_{x\in X} \frac{1}{\#Gx} = \#X/G$. Proceed by grouping terms in this sum according to which orbit they're in:

$$\begin{aligned}
\sum_{x\in X} \frac{1}{\#Gx} &= \sum_{Gx\in X/G} \sum_{y\in Gx} \frac{1}{\#Gx} \\
&= \sum_{Gx\in X/G} \frac{1}{\#Gx} \sum_{y\in Gx} 1 \\
&= \sum_{Gx\in X/G} \frac{1}{\#Gx} \#G_x \\
&= \sum_{Gx\in X/G} 1 \\
&= \#X/G.
\end{aligned}$$

■

2.9.2 Conjugation on Subgroups

Example 2.9.8 ***(?):*** Let G act on $X \coloneqq \left\{H \;\middle|\; H \leq G\right\}$ (its set of *subgroups*) by conjugation.

- The orbit $\mathcal{O}(H) = \left\{gHg^{-1} \;\middle|\; g \in G\right\}$ is the **set of conjugate subgroups** of H.
  - This action is transitive iff all subgroups are conjugate.
- The fixed points $\mathrm{Fix}(G)$ form the set of **normal subgroups** of G.
- The stabilizer $\mathrm{Stab}(H) = N_G(H)$ is the **normalizer** of H in G.
- The kernel is the intersection of all normalizers.

Corollary 2.9.9.
Given $H \leq G$, the number of conjugate subgroups is $[G : N_G(H)]$, i.e.

$$\left|\left\{gHg^{-1} \;\middle|\; g \in G\right\}\right| = [G : N_G(H)].$$

2.9.3 Left Translation on Cosets

Example 2.9.10 ***(?):*** For a fixed proper subgroup $H < G$, let G act on its cosets $X \coloneqq G/H \coloneqq \left\{gH \;\middle|\; g \in G\right\}$ by left translation.

- The orbit $\mathcal{O}(xH) = G/H$, the entire set of cosets.
  - Note that this is a *transitive* action.
- The stabilizer $\mathrm{Stab}(xH) = xHx^{-1}$, a **conjugate subgroup** of H
- The fixed points form $\mathrm{Fix}(G) = \emptyset$.
- The kernel of this action is $\bigcap_{g \in G} gHg^{-1}$, the intersection of all conjugates of H.

Proposition 2.9.11 ***(Application of translation action on cosets).***
If G is a finite group and $p \coloneqq [G : H]$ is the smallest prime dividing $\#G$, then $H \trianglelefteq G$.

Proof (?).

- Let $\varphi : G \curvearrowright X \coloneqq \{xH\}$, noting that $\#X = p$ and $\mathrm{Sym}(X) \cong S_p$.
- Then $K \coloneqq \ker \varphi \subseteq H$.
- Since G is finite and $K \leq G$, we have $[G : K] = \#(G/K) = \#G/\#K$ so $\#(G/K)$ divides $\#G$.
- Since $G/K \cong K'$ is isomorphic to a subgroup of S_p, $\#(G/K)$ divides $\#S_p = p!$
- So $\#(G/K)$ divides $\gcd(\#G, p!)$, which is p since it was the minimal prime dividing $\#G$.
- p is prime, so if $\#G/K \neq 1$ we have $\#G/K = p$.

- Since $K \subset H$ and $[G:H] = p = [G:K]$, we have $K = H$.
- But $K = \ker \varphi \trianglelefteq G$, so $H \trianglelefteq G$.

■

3 | Sylow Theorems

Definition 3.0.1
A **p-group** is a group G such that every element is order p^k for some k. If G is a finite p-group, then $|G| = p^j$ for some j.

Write

- $|G| = p^k m$ where $(p, m) = 1$,
- S_p a Sylow-p subgroup, and
- n_p the number of Sylow-p subgroups.

3.1 Sylow 1 (Cauchy for Prime Powers)

Theorem 3.1.1 ***(Sylow 1).***

$$\forall p^n \text{ dividing } |G|, \text{ there exists a subgroup of size } p^n.$$

Slogan 3.1.2
Sylow p-subgroups exist for any p dividing $|G|$, and are maximal in the sense that every p-subgroup of G is contained in a Sylow p-subgroup. If $|G| = \prod p_i^{\alpha_i}$, then there exist subgroups of order $p_i^{\beta_i}$ for every i and every $0 \leq \beta_i \leq \alpha_i$. In particular, Sylow p-subgroups always exist.

3.2 Sylow 2 (Sylows are Conjugate)

Theorem 3.2.1 ***(Sylow 2).***
All Sylow-p subgroups S_p are conjugate, i.e.

$$S_p^i, S_p^j \in \mathrm{Syl}_p(G) \implies \exists g \text{ such that } g S_p^i g^{-1} = S_p^j$$

Corollary 3.2.2.

$$n_p = 1 \iff S_p \trianglelefteq G.$$

3.3 Sylow 3 (Numerical Constraints)

Theorem 3.3.1 ***(Sylow 3).***

1. $n_p \mid m$ (in particular, $n_p \leq m$),
2. $n_p \equiv 1 \pmod{p}$,
3. $n_p = [G : N_G(S_p)]$ where N_G is the normalizer.

3.4 Corollaries and Applications

Corollary 3.4.1.
By Sylow 3, p does not divide n_p.

Proposition 3.4.2.
Every p-subgroup of G is contained in a Sylow p-subgroup.

Proof .
Let $H \leq G$ be a p-subgroup. If H is not *properly* contained in any other p-subgroup, it is a Sylow p-subgroup by definition. Otherwise, it is contained in some p-subgroup H^1. Inductively this yields a chain $H \subsetneq H^1 \subsetneq \cdots$, and by Zorn's lemma $H := \cup_i H^i$ is maximal and thus a Sylow p-subgroup.

■

3.5 Proof of Sylow Theorems

Proof of Sylows

3.6 Exercises

- Let G be a group of order p with v and e positive integers, p prime, $p > v$, and v is not a multiple of p. Show that G has a normal Sylow p-subgroup.

3.7 Isomorphism Theorems

Theorem 3.7.1 ***(1st Isomorphism Theorem).***
If $\varphi : G \to H$ is a group morphism then

$$G/\ker\varphi \cong \operatorname{im}\varphi.$$

Note: for this to make sense, we also have

- $\ker\varphi \trianglelefteq G$
- $\operatorname{im}\varphi \leq G$

Corollary 3.7.2.
If $\varphi : G \to H$ is surjective then $H \cong G/\ker\varphi$.

Theorem 3.7.3 ***(Diamond Theorem / 2nd Isomorphism Theorem).***
If $S \leq G$ and $N \trianglelefteq G$, then

$$\frac{SN}{N} \cong \frac{S}{S \cap N} \quad \text{and} \quad |SN| = \frac{|S||N|}{|S \cap N|}.$$



Figure 1: The 2nd "Diamond" Isomorphism Theorem

Remark 3.7.4: For this to make sense, we also have

- $SN \leq G$,
- $S \cap N \trianglelefteq S$,

If we relax the conditions to $S, N \leq G$ with $S \in N_G(N)$, then $S \cap N \trianglelefteq S$ (but is not normal in G) and the 2nd Isomorphism Theorem still holds.

Theorem 3.7.5 ***(Cancellation / 3rd Isomorphism Theorem).***
Suppose $N, K \leq G$ with $N \trianglelefteq G$ and $N \subseteq K \subseteq G$.

1. If $K \leq G$ then $K/N \leq G/N$ is a subgroup
2. If $K \trianglelefteq G$ then $K/N \trianglelefteq G/N$.
3. Every subgroup of G/N is of the form K/N for some such $K \leq G$.

4. Every *normal* subgroup of G/N is of the form K/N for some such $K \trianglelefteq G$.
5. If $K \trianglelefteq G$, then we can cancel normal subgroups:

$$\frac{G/N}{K/N} \cong \frac{G}{K}.$$

Theorem 3.7.6 ***(The Correspondence Theorem / 4th Isomorphism Theorem).***
Suppose $N \trianglelefteq G$, then there exists a correspondence:

$$\left\{H < G \;\middle|\; N \subseteq H\right\} \rightleftharpoons \left\{H \;\middle|\; H < \frac{G}{N}\right\}$$
$$\left\{\begin{smallmatrix}\text{Subgroups of } G \\ \text{containing } N\end{smallmatrix}\right\} \rightleftharpoons \left\{\begin{smallmatrix}\text{Subgroups of the} \\ \text{quotient } G/N\end{smallmatrix}\right\}.$$

In words, subgroups of G containing N correspond to subgroups of the quotient group G/N. This is given by the map $H \mapsto H/N$.

Fact 3.7.7
$N \trianglelefteq G$ and $N \subseteq H < G \implies N \trianglelefteq H$.

3.8 Products

Theorem 3.8.1 ***(Chinese Remainder Theorem).***

$$\gcd(p, q) = 1 \implies \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/q\mathbb{Z} \cong \mathbb{Z}/pq\mathbb{Z}.$$

Theorem 3.8.2 ***(Recognizing Direct Products).***
We have $G \cong H \times K$ when

1. $H, K \trianglelefteq G$
2. $G = HK$.
3. $H \cap K = \{e\} \subset G$

Note: can relax to $[h, k] = 1$ for all h, k.

Proof (?).
With these conditions, the following map is an isomorphism:

$$\begin{aligned} \Gamma : H \times K &\to G \\ (h, k) &\mapsto hk. \end{aligned}$$

- This is a group morphism by condition (1):

  $$\Gamma(h_1,k_1)\Gamma(h_2,k_2) \coloneqq (h_1k_1)(h_2k_2) = h_1(k_1h_2)k_2 = h_1(h_2k_1)k_2 = (h_1h_2)(k_1k_2) \coloneqq \Gamma((h_1,k_1)(h_2,k_2)).$$

- This is surjective by condition (2)
- This is injective by condition(3) and checking the kernel:

  $$\ker\Gamma = \left\{(h,k) \;\middle|\; hk = 1_G,\, hk = 1_G\right\} \implies h = k^{-1} \implies hk \in K \cap H = \{1_G\}.$$

■

Theorem 3.8.3 ***(Recognizing Generalized Direct Products).***
We have $G \cong \prod_{i=1}^{n} H_i$ when

- $H_i \trianglelefteq G$ for all i.
- $G = H_1 \cdots H_n$
- $H_k \cap H_1 \cdots \widehat{H_k} \cdots H_n = \emptyset$

Note on notation: intersect H_k with the amalgam leaving out H_k.

Theorem 3.8.4 ***(Recognizing Semidirect Products).***
We have $G \cong N \rtimes_\psi H$ when

- $N \trianglelefteq G$
- $G = NH$
- $H \curvearrowright N$ by conjugation via a map

  $$\begin{aligned}\psi : H &\to \mathrm{Aut}(N)\\ h &\mapsto h(-)h^{-1}.\end{aligned}$$

Relaxed condition: $H, N \trianglelefteq G$ for direct product, or just $H \leq G$ for a semidirect product.

Proposition 3.8.5.
If $H, K \leq G$ and $H \leq N_G(K)$ (or $K \trianglelefteq G$) then $HK \leq G$ is a subgroup.

3.9 Automorphism Groups

Fact 3.9.1

- If $\sigma \in \mathrm{Aut}(H)$ and $\tau \in \mathrm{Aut}(N)$, then $N \rtimes_\psi H \cong N \rtimes_{\tau\circ\psi\circ\sigma} H$.
- $\mathrm{Aut}\left((\mathbb{Z}/p\mathbb{Z})^n\right) \cong \mathrm{GL}(n, \mathbb{F}_p)$, which has size

$$|\mathrm{Aut}(\mathbb{Z}/(p)^n)| = (p^n - 1)(p^n - p)\cdots(p^n - p^{n-1}).$$

  - If this occurs in a semidirect product, it suffices to consider similarity classes of matrices (i.e. just use canonical forms)

- $\mathrm{Aut}(\mathbb{Z}/n\mathbb{Z}) \cong (\mathbb{Z}/n\mathbb{Z})^\times \cong \mathbb{Z}/\varphi(n)\mathbb{Z}$ where φ is the totient function.
  - $\varphi(p^k) = p^{k-1}(p-1)$
- If G, H have coprime order then $\mathrm{Aut}(G \times H) \cong \mathrm{Aut}(G) \times \mathrm{Aut}(H)$.

3.10 Special Classes of Groups

Definition 3.10.1 (2 out of 3 Property)
The **"2 out of 3 property"** is satisfied by a class of groups $\mathcal{C}$ iff whenever $G \in \mathcal{C}$, then $N, G/N \in \mathcal{C}$ for any $N \trianglelefteq G$.

Definition 3.10.2 (p-groups)
If $|G| = p^k$, then G is a **p-group.**

Definition 3.10.3 (Normalizers Grow)
If for every proper $H < G$, $H \trianglelefteq N_G(H)$ is again proper, then "normalizers grow" in G.

3.11 Classification of Groups

General strategy: find a normal subgroup (usually a Sylow) and use recognition of semidirect products.

- Keith Conrad: Classifying Groups of Order 12
- Order p: cyclic.
- Order p^2: abelian, 2 choices.

- Order pq: cases, letting $q < p$ and checking $q \mid p - 1$.
- Order pqr: ?
- Order p^2q: ?

Proposition 3.11.1 ***(PQR Theorem).***
If $|G| = pqr$ where $p < q < r$ are distinct primes then G is solvable.

3.11.1 Finitely Generated Abelian Groups

Definition 3.11.2 (Invariant Factor Decomposition)

$$G \cong \mathbb{Z}^r \times \prod_{j=1}^{m} \mathbb{Z}/n_j\mathbb{Z} \quad \text{where } n_1 \mid \cdots \mid n_m.$$

Invariant factors → Elementary Divisors:

- Take prime factorization of each factor
- Split into coprime pieces

Example 3.11.3:

$$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_{2^3 \cdot 5^2 \cdot 7} \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_{2^3} \times \mathbb{Z}_{5^2} \times \mathbb{Z}_7$$

Elementary divisors → invariant factors:

- Bin up by primes occurring (keeping exponents)
- Take highest power from each prime as *last* invariant factor
- Take highest power from all remaining primes as next, etc

Example 3.11.4: Given the invariant factor decomposition

$$G = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_{5^2}$$

$p = 2$	$p = 3$	$p = 5$
$2, 2, 2$	$3, 3$	5^2

$\implies n_m = 5^2 \cdot 3 \cdot 2$

$p=2$	$p=3$	$p=5$
$2,2$	3	$\emptyset$

$\implies n_{m-1} = 3 \cdot 2$

$p=2$	$p=3$	$p=5$
2	$\emptyset$	$\emptyset$

$\implies n_{m-2} = 2$

and thus

$$G \cong \mathbb{Z}_2 \times \mathbb{Z}_{3\cdot 2} \times \mathbb{Z}_{5^2 \cdot 3 \cdot 2}$$

3.11.2 Classifying Abelian Groups of a Given Order

Let $p(x)$ be the integer partition function.

Example 3.11.5 ***(of an integer partition):*** Example: $p(6) = 11$, given by

$$\begin{aligned}
&[6],\\
&[5,1],\\
&[4,2],\\
&[4,1,1],\\
&[3,3],\\
&[3,2,1],\\
&[3,1,1,1],\\
&[2,2,2],\\
&[2,2,1,1],\\
&[2,1,1,1,1],\\
&[1,1,1,1,1,1].
\end{aligned}$$

Write $G = p_1^{k_1} p_2^{k_2} \cdots$; then there are $p(k_1)p(k_2)\cdots$ choices, each yielding a distinct group.

Add example

3.12 Series of Groups

Definition 3.12.1 (Normal Series)
A **normal series** of a group G is a sequence $G \to G^1 \to G^2 \to \cdots$ such that $G^{i+1} \trianglelefteq G_i$ for every i.

Definition 3.12.2 (Central Series)
A **central series** for a group G is a terminating normal series $G \to G^1 \to \cdots \to \{e\}$ such that each quotient is **central**, i.e. $[G, G^i] \leq G^{i-1}$ for all i.

Definition 3.12.3 (Composition Series)
A **composition series** of a group G is a finite normal series such that G^{i+1} is a *maximal proper* normal subgroup of G^i.

Theorem 3.12.4 ***(Jordan-Holder).***
Any two composition series of a group have the same length and isomorphic composition factors (up to permutation).

Definition 3.12.5 (Simple Groups)
A group G is **simple** iff $H \trianglelefteq G \implies H = \{e\}, G$, i.e. it has no non-trivial proper subgroups.

Proposition 3.12.6.
If G is *not* simple, then G is an extension of any of its normal subgroups. I.e. for any $N \trianglelefteq G$, $G \cong E$ for some extension of the form $N \to E \to G/N$.

Definition 3.12.7 (Lower Central Series)
Set $G^0 = G$ and $G^{i+1} = [G, G^i]$, then $G^0 \geq G^1 \geq \cdots$ is the *lower central series* of G.

Mnemonic: "lower" because the chain is descending. Iterate the adjoint map $[-, G]$, if this terminates then the map is nilpotent, so call G nilpotent!

Definition 3.12.8 (Upper Central Series)
Set $Z_0 = 1$, $Z_1 = Z(G)$, and $Z_{i+1} \leq G$ to be the subgroup satisfying $Z_{i+1}/Z_i = Z(G/Z_i)$. Then $Z_0 \leq Z_1 \leq \cdots$ is the *upper central series* of G.
Equivalently, since $Z_i \trianglelefteq G$, there is a quotient map $\pi : G \to G/Z_i$, so define $Z_{i+1} := \pi^{-1}(Z(G/Z_i))$ (?).

Mnemonic: "upper" because the chain is ascending. "Take higher centers".

Definition 3.12.9 (Derived Series)
Set $G^{(0)} = G$ and $G^{(i+1)} = [G^{(i)}, G^{(i)}]$, then $G^{(0)} \geq G^{(1)} \geq \cdots$ is the *derived series* of G.

Definition 3.12.10 (Solvable)
A group G is **solvable** iff G has a terminating normal series with abelian composition factors, i.e.

$$G \to G^1 \to \cdots \to \{e\} \text{ with } G^i/G^{i+1} \text{ abelian for all } i.$$

Theorem 3.12.11 ***(Characterization of Solvable).***
A group G is solvable iff its derived series terminates.

Theorem 3.12.12 ***(S_n is Almost Always Solvable).***
If $n \geq 4$ then S_n is solvable.

Lemmas:

- G is solvable iff G has a terminating *derived series*.
- Solvable groups satisfy the 2 out of 3 property
- Abelian $\implies$ solvable
- Every group of order less than 60 is solvable.

4 | Ring Theory

4.1 Definitions

Notation:

- $\langle a \rangle \coloneqq Ra \coloneqq \left\{ ra \;\middle|\; r \in R \right\}$ is the ideal generated by a single element.

4.1.1 Undergrad Review

Definition 4.1.1 (Divisibility of Elements)
An element $r \in R$ is **divisible** by $q \in R$ if and only if there exists some $c \in R$ such that $r = qc$. In this case, we sometimes write $q \mid r$.

Definition 4.1.2 (Irreducible Element)
An element $r \in R$ is **irreducible** iff

$$r = ab \implies a \in R^\times \text{ or } b \in R^\times$$

Definition 4.1.3 (Prime Element)
An element $p \in R$ is **prime** iff

$$a, b \in R^\times \setminus \{0\}, \quad ab \mid p \implies a \mid p \text{ or } b \mid p.$$

Fact 4.1.4
If R is an integral domain, prime $\implies$ irreducible. If R is a UFD, then prime $\iff$ irreducible.

Definition 4.1.5 (Zero Divisor)
An element $r \in R$ is a **zero-divisor** iff there exists an $a \in R \setminus \{0\}$ such that $ar = ra = 0$. Equivalently, the map

$$\begin{aligned} r_- : R &\to R \\ x &\mapsto rx \end{aligned}$$

fails to be injective.

Definition 4.1.6 (Associate Elements)
$a, b \in R$ are **associates** iff there exists a $u \in R^\times$ such that $a = ub$. Equivalently, $a \mid b$ and $b \mid a$.

Definition 4.1.7 (Irreducible Ideal)
An ideal $I \trianglelefteq R$ is **irreducible** if it can not be written as the intersection of two larger ideals, i.e. there are not $J_1, J_2 \supseteq I$ such that $J_1 \cap J_2 = I$.

Definition 4.1.8 (Prime Ideal)
$\mathfrak{p}$ is a **prime** ideal $\iff$

$$ab \in \mathfrak{p} \implies a \in \mathfrak{p} \text{or} b \in \mathfrak{p}.$$

Definition 4.1.9 (Maximal Ideal)
$\mathfrak{m}$ is **maximal** $\iff$ $(I \trianglelefteq R, I \neq R \implies I \subseteq \mathfrak{m}) \iff R/I$ is a field.

Remark 4.1.10:

- A non-maximal, non-prime ideal: $\left\langle x^4 + 2x^2 + 1 \right\rangle \trianglelefteq \mathbb{C}[x]$
- An ideal that is both prime and maximal: $\langle f \rangle$ for any $f \in \mathbb{Q}[x]^{\text{irr}}$
- A prime ideal that is not maximal: $\langle x - c \rangle \in \mathbb{R}[x]$.

Proposition 4.1.11 *(?).*

- $R/\mathfrak{m}$ is an integral domain $\iff$ $\mathfrak{m} \in \text{mSpec}(R)$ is maximal.
- $R/\mathfrak{p} \in \mathsf{Field} \iff \mathfrak{p} \in \text{Spec}(R)$ is prime.

Definition 4.1.12 (Prime Spectrum)
The **prime spectrum** (or just the **spectrum**) of R is defined as

$$\mathrm{Spec}(R) = \left\{ \mathrm{pr} \trianglelefteq R \;\middle|\; \mathrm{pr} \text{ is prime} \right\}.$$

Definition 4.1.13 (Max Spectrum)
The **max spectrum** of R is defined as

$$\mathrm{mSpec}(R) = \left\{ \mathfrak{m} \trianglelefteq R \;\middle|\; \mathfrak{m} \text{ is maximal} \right\}.$$

Definition 4.1.14 (Integral Domain)
A ring is an **integral domain** if and only if it has no nonzero zero divisors:

$$a, b \in R \setminus \{0\}, ab = 0 \implies a = 0 \text{or} b = 0.$$

Definition 4.1.15 (Principal Ideal)
An ideal $I \trianglelefteq R$ if **principal** if there exists an $a \in R$ such that $I = \langle a \rangle$.

Definition 4.1.16 (Principal Ideal Domain)
A ring R is a **principal ideal domain** iff every ideal is principal.

Definition 4.1.17 (Unique Factorization Domain)
A ring R is a **unique factorization domain** iff R is an integral domain and every $r \in R \setminus \{0\}$ admits a decomposition

$$r = u \prod_{i=1}^{n} p_i$$

where $u \in R^\times$ and the p_i irreducible, which is unique up to associates.

4.1.2 Types of Rings

Definition 4.1.18 (Simple Modules)
A module M is **simple** iff every submodule $M' \leq M$ is either 0 or M. A ring R is simple if and only if it is simple as an R-module, i.e. there are no nontrivial proper ideals.

Definition 4.1.19 (Semisimple Modules)
A module M is **simple** if and only if it admits a decomposition

$$M = \bigoplus_{j \in J} M_j$$

with each M_j simple.

Definition 4.1.20 (Noetherian)
A ring R is **Noetherian** if the ACC holds: every ascending chain of ideals $I_1 \leq I_2 \cdots$ stabilizes in the sense that there exists some N such that $I_N = I_{N+1} = \cdots$.

4.1.3 Commutative Algebra

Definition 4.1.21 (Primary Ideal)
An ideal $I \trianglelefteq R$ is **primary** iff whenever $pq \in I$, $p \in I$ and $q^n \in I$ for some n.

Definition 4.1.22 (Nilradical)
$\mathfrak{N}(R) \coloneqq \left\{x \in R \;\middle|\; x^n = 0 \text{ for some } n\right\}$ is the **nilradical** of R.

Definition 4.1.23 (Jacobson Radical)
The **Jacobson radical** $\mathfrak{J}(R)$ is the intersection of all maximal ideals, i.e.

$$\mathfrak{J}(R) = \cap \left\{\mathfrak{m} \;\middle|\; \mathfrak{m} \in \operatorname{maxSpec}(R)\right\}.$$

Definition 4.1.24 (Reduced Ring)
A ring R is **reduced** if R contains no nonzero nilpotent elements.

Definition 4.1.25 (Local Ring)
A ring R is **local** iff it contains a unique maximal ideal.

Definition 4.1.26 (Radical of an Ideal)
For an ideal $I \trianglelefteq R$, the **radical** $\operatorname{rad}(I) \coloneqq \left\{r \in R \;\middle|\; r^n \in I \text{ for some } n \geq 0\right\}$, so $x^n \in I \iff x \in I$.

Definition 4.1.27 (Radical Ideal)
An ideal is **radical** iff $\operatorname{rad}(I) = I$.

4.2 Structure Theorems

Proposition 4.2.1 ***(Characterizations of Rings).***

- R a commutative division ring $\implies$ R is a field
- R a finite integral domain $\implies$ R is a field.
- $\mathbb{F}$ a field $\implies$ $\mathbb{F}[x]$ is a Euclidean domain.
- $\mathbb{F}$ a field $\implies$ $\mathbb{F}[x]$ is a PID.
- $\mathbb{F}$ is a field $\iff$ $\mathbb{F}$ is a commutative simple ring.
- R is a UFD $\iff$ $R[x]$ is a UFD.

- R a PID $\implies$ $R[x]$ is a UFD
- R a PID $\implies$ R Noetherian
- $R[x]$ a PID $\implies$ R is a field.

Proposition 4.2.2.
Fields $\subset$ Euclidean domains $\subset$ PIDs $\subset$ UFDs $\subset$ Integral Domains $\subset$ Rings

Example 4.2.3:

- A Euclidean Domain that is not a field: $\mathbb{F}[x]$ for $\mathbb{F}$ a field
  - *Proof*: Use previous lemma, and x is not invertible
- A PID that is not a Euclidean Domain: $\mathbb{Z}\left[\frac{1+\sqrt{-19}}{2}\right]$.
  - *Proof*: complicated.
- A UFD that is not a PID: $\mathbb{F}[x, y]$.
  - *Proof*: $\langle x, y\rangle$ is not principal
- An integral domain that is not a UFD: $\mathbb{Z}[\sqrt{-5}]$
  - *Proof*: $(2+\sqrt{-5})(2-\sqrt{-5}) = 9 = 3 \cdot 3$, where all factors are irreducible (check norm).
- A ring that is not an integral domain: $\mathbb{Z}/(4)$
  - *Proof*: $2 \pmod 4$ is a zero divisor.

Proposition 4.2.4.
In R a UFD, an element $r \in R$ is prime $\iff$ r is irreducible.

Note: For R an integral domain, prime $\implies$ irreducible, but generally not the converse.

$x^2 \pmod{()x^2 + x} \in \mathbb{Q}[x]/(x^2 + x)$. Check that x is prime directly, but $x = x \cdot x$ and x is not a unit.

Proposition 4.2.5.
If R is a PID, then every element in R has a unique prime factorization.

Theorem 4.2.6 ***(Krull).***
Every ring has proper maximal ideals, and any proper ideal is contained in a maximal ideal.

Theorem 4.2.7 ***(Artin-Wedderburn?).***
If R is a nonzero, unital, *semisimple* ring then $R \cong \bigoplus_{i=1}^{m} \mathrm{Mat}(n_i, D_i)$, a finite sum of matrix rings over division rings.

Corollary 4.2.8.
If M is a simple ring over R a division ring, the M is isomorphic to a matrix ring.

4.3 Zorn's Lemma

Theorem 4.3.1 ***(Zorn's Lemma).***
If P is a poset in which every chain has an upper bound, then P has a maximal element.

Proposition 4.3.2.
Fields are simple rings.

Proposition 4.3.3.
If $I \trianglelefteq R$ is a proper ideal $\iff$ I contains no units.

Proof .
$r \in R^\times \cap I \implies r^{-1}r \in I \implies 1 \in I \implies x \cdot 1 \in I \quad \forall x \in R.$
■

Proposition 4.3.4.
If $I_1 \subseteq I_2 \subseteq \cdots$ are ideals then $\cup_j I_j$ is an ideal.

Proposition 4.3.5.
Every proper ideal is contained in a maximal ideal.

Proof .
Let $0 < I < R$ be a proper ideal, and consider the set

$$S = \left\{ J \;\middle|\; I \subseteq J < R \right\}.$$

Note $I \in S$, so S is nonempty. The claim is that S contains a maximal element M.
S is a poset, ordered by set inclusion, so if we can show that every chain has an upper bound, we can apply Zorn's lemma to produce M.
Let $C \subseteq S$ be a chain in S, so $C = \{C_1 \subseteq C_2 \subseteq \cdots\}$ and define $\widehat{C} = \cup_i C_i$.
$\widehat{C}$ **is an upper bound for** C**:** This follows because every $C_i \subseteq \widehat{C}$.
$\widehat{C}$ **is in** S**:** Use the fact that $I \subseteq C_i < R$ for every C_i and since no C_i contains a unit, $\widehat{C}$ doesn't contain a unit, and is thus proper.
■

Example 4.3.6 ***(An irreducible element that is not prime.):*** $3 \in \mathbb{Z}[\sqrt{-5}]$. Check norm to see irreducibility, but $3 \mid 9 = (2+\sqrt{-5})(2-\sqrt{-5})$ and doesn't divide either factor.

Example 4.3.7: Maximal ideals of $R[x]$ are of the form $I = (x - a_i)$ for some $a_i \in R$.

4.4 Unsorted

Proposition 4.4.1 *(Correspondence Theorem).*

todo

Fact 4.4.2
Division algorithm for Euclidean domains.

todo

Definition 4.4.3 (Localization)

todo

Definition 4.4.4 (Fraction Field)

todo

Theorem 4.4.5 *(Hilbert Basis Theorem).*

todo

5 Ring Theory

6 Field Theory

6.1 Facts and Definitions

Let k denote a field, and L/k extensions.

Remark 6.1.1:

- $[L:K] = \dim_{\mathsf{Vect}_K} L$, the dimension of L as a K-vector space
- $\mathrm{Aut}(L/k) \coloneqq \left\{\sigma : L \to L' \;\middle|\; \sigma|_K = \mathbb{1}_K\right\}$, the lifts of the identity on K.
- $\{L:K\} \coloneqq \#\mathrm{Aut}(L/k) \coloneqq \left\{\sigma : L \to L' \;\middle|\; \sigma|_K = \mathbb{1}_K\right\}$, the number of lifts of the identity on K.
- $\mathrm{Gal}(E/F) \coloneqq \mathrm{Aut}_{\mathsf{Fields}_k}$ if E is finite, normal, and separable.

Definition 6.1.2 (Simple Extensions)
An extension L/k is **simple** iff $L = K(\alpha)$ for some $\alpha \in L$.

Definition 6.1.3 (Fixed Field)
For $H \leq \mathrm{Aut}_{\mathsf{Fields}_k}(L)$,

$$L^H := \left\{ \ell \in L \;\middle|\; \sigma(l) = \ell \right\}.$$

Definition 6.1.4 (Prime Subfield)
The **prime subfield** of a field F is the subfield generated by 1.

Fact 6.1.5

- Irreducible $\implies$ separable.
- The minimal polynomial of an algebraic element is always irreducible.
- Every irreducible polynomial over a perfect field is separable.
- Every finite extension of a perfect field is separable.

Definition 6.1.6 (Field Automorphisms)

$$\mathrm{Aut}(L/k) = \left\{ \sigma : L \to L \;\middle|\; \sigma|_k = \mathrm{id}_k \right\}.$$

Definition 6.1.7 (Reducible and Irreducible Polynomials)
For $\mathbb{F}$ a field, a polynomial $f \in \mathbb{F}[x]$ is **reducible** if and only if f can be factored as $f(x) = g(x)h(x)$ for some $g, h \in \mathbb{F}[x]$ with $\deg g, \deg h \geq 1$ (so g, h are nonconstant). f is **irreducible** if f is not reducible.

Remark 6.1.8: Note that in general,

$$|\mathrm{Aut}(L/k)| \leq [L : K].$$

Definition 6.1.9 (Characterizations of Perfect Fields)
The following are equivalent:

- k is a **perfect** field.
- Every irreducible polynomial $p \in k[x]$ is separable
- Every finite extension F/k is separable.
- If $\mathrm{ch}\, k > 0$, the Frobenius is an automorphism of k.

Fact 6.1.10

- The characteristic of any field k is either 0 or p a prime.
- All fields are simple rings (no proper nontrivial ideals).
- If L/k is algebraic, then $\min(\alpha, L)$ divides $\min(\alpha, k)$.
- Every field morphism is either zero or injective.

Theorem 6.1.11 ***(Gauss' Lemma).***
Let R be a UFD and F its field of fractions. Then a primitive $p \in R[x]$ is irreducible in $R[x] \iff p$ is irreducible in $F[x]$.

Corollary 6.1.12.
A primitive polynomial $p \in \mathbb{Q}[x]$ is irreducible $\iff p$ is irreducible in $\mathbb{Z}[x]$.

Theorem 6.1.13 ***(Eisenstein's Criterion).***
If $f(x) = \sum_{i=0}^{n} \alpha_i x^i \in \mathbb{Q}[x]$ and $\exists p$ such that

- p divides every coefficient *except* a_n and
- p^2 does not divide a_0,

then f is irreducible over $\mathbb{Q}[x]$, and by Gauss' lemma, over $\mathbb{Z}[x]$.

Definition 6.1.14 (Elementary Symmetric Functions)
Todo

6.2 Extensions

Definition 6.2.1 (Algebraic Field Extension)
A field extension L/k is **algebraic** iff every $\alpha \in L$ is the root of some polynomial $f \in k[x]$.

Proposition 6.2.2 ***(Normal Field Extension).***
Let L/k be a finite extension. Then TFAE:

- L/k is **normal**.
- Every irreducible polynomial $f \in k[x]$ that has one root in L has *all* of its roots in L
  - i.e. every polynomial splits into linear factors
- Either f splits in L or f has no roots in L.
- Every embedding $\sigma : L \hookrightarrow \bar{k}$ that is a lift of the identity on k satisfies $\sigma(L) = L$.
- If L is separable: L is the splitting field of some irreducible $f \in k[x]$.

Proof (?).

todo

■

Definition 6.2.3 (Separable Field Extension)
Let L/k be a field extension, $\alpha \in L$ be arbitrary, and $f(x) \coloneqq \min(\alpha, k)$. The following are equivalent

- L/k is **separable**
- Every element $\alpha \in L$ has separable minimal polynomial $\min_{\alpha,L}(x) \in \bar{k}[x]$ (D&F's definition, p. 551).
- f has no repeated factors/roots, i.e. f has distinct roots in L.
- $\gcd(f, f') = 1$.
- $f' \not\equiv 0$

If L/k is a finite extension, then, TFAE:

- L/k is separable.
- $L = k(\alpha)$ for α a separable element.
- $L = k(\{\alpha_i\})$ for α_i separable elements
- $[L : k] = \{L : k\} \coloneqq \#\mathrm{Aut}_{\mathsf{Fields}_k}(L)$

Definition 6.2.4 (Galois Extension and Galois Group)
Let L/k be a finite field extension. The following are equivalent:

1. L/k is a **Galois extension**.
2. $\#\mathrm{Aut}_{\mathsf{Fields}_k}(L) = [L : k] = \{L : k\}$ (D&F's definition).
3. The fixed field of $\mathrm{Aut}(L/k)$ is exactly k.
4. L is the splitting field of a separable polynomial $p \in K[x]$.
5. L is finite, normal, and separable (most general definition?)

- L is a finite separable splitting field of an irreducible polynomial.
- L/k is separable and normal.
- The fixed field L^H for $H \coloneqq \mathrm{Aut}_{\mathsf{Fields}_k}(L)$ is precisely k.

In this case, we define the **Galois group** as

$$\mathrm{Gal}(L/k) \coloneqq \mathrm{Aut}_{\mathsf{Fields}_k}(L).$$

Definition 6.2.5 (Simple Extension)
An extension F/k is **simple** if $F = k[\alpha]$ for a single element α.

Definition 6.2.6 (Primitive Extension)
For R a UFD, a polynomial $p \in R[x]$ is **primitive** iff the greatest common divisors of its coefficients is a unit.

Proposition 6.2.7 ***(Classification of quadratic extensions).***

todo

If $\mathbb{F}$ is a field with $\mathrm{ch}(\mathbb{F}) \neq 2$ and $E_{/\mathbb{F}}$ is a degree 2 extension, then E is Galois and $E = F(\sqrt{a})$ for some squarefree $a \in \mathbb{F}$.

Corollary 6.2.8 ***(Quadratic extensions of rationals).***

If $E_{/\mathbb{Q}}$ is a quadratic extension, $E = \mathbb{Q}(\sqrt{\frac{p}{q}})$ for some $p, q \in \mathbb{Z}$.

Proposition 6.2.9 ***(?).***

For $\mathbb{F}_p$ a finite field of prime order, all quadratic extensions $E/\mathbb{F}_p$ are isomorphic.

Theorem 6.2.10 ***(Finite Extensions are Algebraic).***

Every finite extension is algebraic.

Proof .

If K/F and $[K : F] = n$, then pick any $\alpha \in K$ and consider $1, \alpha, \alpha^2, \ldots$. This yields $n+1$ elements in an n-dimensional vector space, and thus there is a linear dependence

$$f(\alpha) \coloneqq \sum_{j=1}^{n} c_j \alpha^j = 0.$$

But then α is the root of the polynomial f.

■

Theorem 6.2.11 ***(Primitive Element Theorem).***

Every finite separable extension is simple.

Corollary 6.2.12.

$\mathbb{GF}(p^n)$ is a simple extension over $\mathbb{F}_p$.

Proposition 6.2.13 ***(?).***

If L/k is separable, then

$$[L : k] = \{L : k\}.$$

If L/k is a splitting field, then

$$[L : K] = \#\mathrm{Aut}_{\mathsf{Fields}_k}(L) \coloneqq \#\,\mathrm{Gal}(L/k).$$

6.3 Finite Fields

Theorem 6.3.1 ***(Characterization of Prime Subfields).***
The prime subfield of any field is isomorphic to either $\mathbb{Q}$ or $\mathbb{F}_p$ for some p.

Proposition 6.3.2.
If $\operatorname{ch} k = p$ then $(a+b)^p = a^p + b^p$ and $(ab)^p = a^p b^p$.

Proof .
Todo

■ Proof

Theorem 6.3.3 ***(Construction of Finite Fields).***
$\mathbb{GF}(p^n) \cong \dfrac{\mathbb{F}_p}{(f)}$ where $f \in \mathbb{F}_p[x]$ is any irreducible of degree n, and $\mathbb{GF}(p^n) \cong \mathbb{F}[\alpha] \cong \operatorname{span}_{\mathbb{F}} \left\{1, \alpha, \cdots, \alpha^{n-1}\right\}$ for any root α of f.

Proposition 6.3.4 ***(Prime Subfields of Finite Fields).***
Every finite field F is isomorphic to a unique field of the form $\mathbb{GF}(p^n)$ and if $\operatorname{ch} F = p$, it has prime subfield $\mathbb{F}_p$.

Proposition 6.3.5 ***(Containment of Finite Fields).***
$\mathbb{GF}(p^\ell) \leq \mathbb{GF}(p^k) \iff \ell$ divides k.

Proposition 6.3.6 ***(Identification of Finite Fields as Splitting Fields).***
$\mathbb{GF}(p^n)$ is the splitting field of $\rho(x) = x^{p^n} - x$, and the elements are exactly the roots of ρ.

Proof .
Todo. Every element is a root by Cauchy's theorem, and the p^n roots are distinct since its derivative is identically -1.

■

Proposition 6.3.7 ***(Splits Product of Irreducibles).***
Let $\rho_n \coloneqq x^{p^n} - x$. Then $f(x) \mid \rho_n(x) \iff \deg f \mid n$ and f is irreducible.

Corollary 6.3.8.
$x^{p^n} - x = \prod f_i(x)$ over all irreducible monic $f_i \in \mathbb{F}_p[x]$ of degree d dividing n.

Proof .
$\Longleftarrow$:

- Suppose f is irreducible of degree d.
- Then $f \mid x^{p^d} - x$, by considering $F[x]/\langle f \rangle$.

- Thus $x^{p^d} - x \mid x^{p^n} - x \iff d \mid n$.

$\implies$:

- $\alpha \in \mathbb{GF}(p^n) \iff \alpha^{p^n} - \alpha = 0$, so every element is a root of φ_n and $\deg \min(\alpha, \mathbb{F}_p) \mid n$ since $\mathbb{F}_p(\alpha)$ is an intermediate extension.
- So if f is an irreducible factor of φ_n, f is the minimal polynomial of some root α of φ_n, so $\deg f \mid n$.
- $\varphi_n'(x) = p^n x^{p^n - 1} \neq 0$, so φ_n is squarefree and thus has no repeated factors. So φ_n is the product of all such irreducible f.

■

Proposition 6.3.9 ***(Finite fields are not algebraically closed).***
If $\mathbb{F}$ is a finite field then $F \neq \overline{F}$.

Proof .
If $k = \{a_1, a_2, \cdots a_n\}$ then define the polynomial

$$f(x) \coloneqq 1 + \prod_{j=1}^{n} (x - a_j) \in k[x].$$

This has no roots in k.

■ Proof

6.4 Cyclotomic Polynomials

Definition 6.4.1 (Cyclotomic Field)
Any subfield of the splitting field E of $f(x) = x^m - 1$ is a **cyclotomic field**.

Proposition 6.4.2.
$\deg \Phi_n(x) = \varphi(n)$ for φ the totient function.

Proof .
$\deg \Phi_n(x)$ is the number of nth primitive roots, which is the number of numbers less than and coprime to n.

■

Proposition 6.4.3 ***(Computing Totient Functions).***
Computing Φ_n:

1.

$$\Phi_n(z) = \prod_{d|n, d>0} \left(z^d - 1\right)^{\mu\left(\frac{n}{d}\right)}$$

where

$$\mu(n) \equiv \begin{cases} 0 & \text{if } n \text{ has one or more repeated prime factors} \\ 1 & \text{if } n = 1 \\ (-1)^k & \text{if } n \text{ is a product of } k \text{ distinct primes,} \end{cases}$$

2.

$$x^n - 1 = \prod_{d|n} \Phi_d(x) \implies \Phi_n(x) = \frac{x^n - 1}{\prod_{\substack{d|n \\ d<n}} \Phi_d(x)},$$

so just use polynomial long division.

Proposition 6.4.4.

$$\Phi_p(x) = x^{p-1} + x^{p-2} + \cdots + x + 1$$
$$\Phi_{2p}(x) = x^{p-1} - x^{p-2} + \cdots - x + 1$$

Proposition 6.4.5.

$$k \mid n \implies \Phi_{nk}(x) = \Phi_n\left(x^k\right)$$

Definition 6.4.6 (Cyclotomic Polynomials)
Let $\zeta_n = e^{2\pi i/n}$, then the n**th cyclotomic polynomial** is given by

$$\Phi_n(x) = \prod_{\substack{k=1 \\ (j,n)=1}}^{n} \left(x - \zeta_n^k\right) \in \mathbb{Z}[x],$$

which is a product over primitive roots of unity. It is the unique irreducible polynomial which is a divisor of $x^n - 1$ but *not* a divisor of $x^k - 1$ for any $k < n$.

Proposition 6.4.7 ***(Table of cyclotomic polynomials).***

todo

Proposition 6.4.8 ***(Galois Groups of Cyclotomic Fields).***
For ζ any primitive root of unity, $\mathrm{Gal}(\mathbb{Q}(\zeta_m)_{/\mathbb{Q}}) \cong (\mathbb{Z}/m\mathbb{Z})^\times$.

Theorem 6.4.9 ***(Kronecker-Weber).***
If $K_{/\mathbb{Q}}$ is an abelian extension, then $K \subseteq \mathbb{Q}(\zeta_m)$ for some m.

6.5 Splitting Fields

Proposition 6.5.1 ***(Splitting Fields of Cyclotomic Polynomials).***
The splitting field of $x^m - 1$ is $\mathbb{Q}(\zeta_m)$ for ζ_m any primitive root of unity.

6.6 Unsorted

Proposition 6.6.1 ***(?).***
A polynomial f has multiple roots $\iff$ $\gcd(f, f') \neq 1$, and the multiple roots of f are precisely the roots of $\gcd(f, f')$.

Proposition 6.6.2 ***(?).***
Irreducible polynomials have distinct roots after passing to a splitting field.

Proof (?).

todo

■

7 Galois Theory

Proposition 7.0.1.
If $\operatorname{ch} k = 0$ or k is finite, then every *algebraic* extension L/k is separable.

Proposition 7.0.2.
If L/k is algebraic, then $\operatorname{Aut}(L/k)$ permutes the roots of irreducible polynomials.

Proposition 7.0.3.
$|\operatorname{Aut}(L/k)| \leq [L : k]$ with equality precisely when L/k is normal.

Theorem 7.0.4 ***(Dirichlet's Theorem on Arithmetic Progressions).***

todo

Definition 7.0.5 (Constructible)

todo

Theorem 7.0.6 ***(Constructibility of Regular n-gons).***

7.0.1 Lemmas About Towers

Let $L/F/k$ be a finite tower of field extensions.

Proposition 7.0.7 ***(Towers are multiplicative in degree).***

$$[L:k] = [L:F][F:k].$$

Proposition 7.0.8 ***(Normal/Algebraic/Galois in towers).***
L/k normal/algebraic/Galois $\implies$ L/F normal/algebraic/Galois.

Proof (for normality).
$\min(\alpha, F) \mid \min(\alpha, k)$, so if the latter splits in L then so does the former.
■

Corollary 7.0.9 ***(?).***
$\alpha \in L$ algebraic over $k \implies \alpha$ algebraic over F.

Corollary 7.0.10 ***(?).***
E_1/k normal and E_2/k normal $\implies$ E_1E_2/k normal and $E_1 \cap E_2/k$ normal.



Link to diagram

Proposition 7.0.11 ***(Algebraicity is transitive).***
F/k algebraic and L/F algebraic $\implies$ L/k algebraic.



Proposition 7.0.12 ***(Separability is transitive).***
For $L/F/k$, then L/k is separable $\iff$ L/F, F/k are separable.



⚠ **Warning 7.0.13**
Being Galois is **not** transitive. Take $\mathbb{Q}\left(\sqrt[4]{2}\right)/\mathbb{Q}\left(\sqrt{2}\right)/\mathbb{Q}$.

Proposition 7.0.14 ***(?).***
If L/k is algebraic, then F/k separable:



Link to diagram

Moreover, L/F is additionally separable $\iff$ L/k separable:



Link to diagram

Proposition 7.0.15 ***(?).***
If L/k is Galois, then L/F is **always** Galois. Moreover, F/k is Galois if and only if $\mathrm{Gal}(L/F) \trianglelefteq \mathrm{Gal}(L/k)$



Link to diagram

In this case,

$$\mathrm{Gal}(F/k) \cong \frac{\mathrm{Gal}(L/k)}{\mathrm{Gal}(L/F)}.$$

7.0.2 Fundamental Theorem of Galois Theory

Theorem 7.0.16 ***(Fundamental Theorem of Galois Theory).***
Let L/k be a Galois extension, then there is a correspondence:

$$\left\{\text{Subgroups } H \leq \mathrm{Gal}(L/k)\right\} \rightleftharpoons \left\{\begin{matrix}\text{Fields } F \text{ such} \\ \text{that } L/F/k\end{matrix}\right\}$$
$$H \to \left\{E^H := \text{ The fixed field of } H\right\}$$
$$\left\{\mathrm{Gal}(L/F) := \left\{\sigma \in \mathrm{Gal}(L/k) \;\middle|\; \sigma(F) = F\right\}\right\} \leftarrow F$$

- This is contravariant with respect to subgroups/subfields.
- $[F : k] = [G : H]$, so degrees of extensions over the base field correspond to indices of

subgroups.

- $[K : F] = |H|$
- L/F is Galois and $Gal(K/F) = H$
- F/k is Galois $\iff$ H is normal, and $\mathrm{Gal}(F/k) = \mathrm{Gal}(L/k)/H$.
- The compositum F_1F_2 corresponds to $H_1 \cap H_2$.
- The subfield $F_1 \cap F_2$ corresponds to H_1H_2.

7.0.3 Examples

Example 7.0.17 ***(Cyclotomic Fields):*** $\mathrm{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q}) \cong \mathbb{Z}/(n)^\times$ and is generated by maps of the form $\zeta_n \mapsto \zeta_n^j$ where $(j, n) = 1$. I.e., the following map is an isomorphism:

$$\begin{aligned} \mathbb{Z}/(n)^\times &\to \mathrm{Gal}(\mathbb{Q}(\zeta_n), \mathbb{Q}) \\ r \pmod n &\mapsto (\varphi_r : \zeta_n \mapsto \zeta_n^r) \end{aligned}$$

Example 7.0.18 ***(Finite Fields):*** $\mathrm{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p) \cong \mathbb{Z}/\langle n \rangle$, a cyclic group generated by powers of the Frobenius automorphism:

$$\begin{aligned} \varphi_p : \mathbb{F}_{p^n} &\to \mathbb{F}_{p^n} \\ x &\mapsto x^p \end{aligned}$$

See D&F p.566 example 7.

Example 7.0.19 ***(Quadratic Fields):*** Every degree 2 extension L/k is Galois, except possibly in characteristic 2: if $\alpha \in L \setminus k$ then $\min_\alpha(x) \in L[x]$ must split in $L[x]$ since $\alpha \in L \implies \min_\alpha(x) = (x - \alpha)g(x)$ which forces $\deg(g) = 1$. So L is a splitting field. If $\mathrm{ch}(k) \neq 2$, then $\dfrac{\partial}{\partial x} \min_\alpha(x) = 2x - \cdots \not\equiv 0$, making L separable.

Proposition 7.0.20.
If K is the splitting field of an irreducible polynomial of degree n, then $\mathrm{Gal}(K/\mathbb{Q}) \leq S_n$ is a transitive subgroup.

Corollary 7.0.21.
n divides the order $|\mathrm{Gal}(K/\mathbb{Q})|$.

Theorem 7.0.22 ***(Splitting + Perfect implies Galois).***

- If $\mathrm{ch}\, k = 0$ or k is finite, then k is perfect.

- $k = \mathbb{C}, \mathbb{R}, \mathbb{Q}, \mathbb{F}_p$ are perfect, so any finite normal extension is Galois.
- Every splitting field of a polynomial over a perfect field is Galois.

Proposition 7.0.23 ***(Composite Extensions).***
If F/k is finite and Galois and L/k is arbitrary, then FL/L is Galois and

$$\mathrm{Gal}(FL/L) = \mathrm{Gal}(F/F \cap L) \subset \mathrm{Gal}(F/k).$$

7.0.4 Counterexamples

Example 7.0.24 ***(?):***

- $\mathbb{Q}(\zeta_3, 2^{1/3})$ is normal but $\mathbb{Q}(2^{1/3})$ is not since the irreducible polynomial $x^3 - 2$ has only one root in it.
- $\mathbb{Q}(2^{1/3})$ is not Galois since its automorphism group is too small (only of size 1 instead of 3?).
- $\mathbb{Q}(2^{1/4})$ is not Galois since its automorphism group is too small (only of size 2 instead of 4). However, the intermediate extensions $\mathbb{Q}(2^{1/4})/\mathbb{Q}(2^{1/2})$ and $\mathbb{Q}(\sqrt{2})/\mathbb{Q}$ are Galois since they are quadratic. Slogan: "Being Galois is not transitive in towers."
- A quadratic extension that is not Galois: $\mathrm{SF}(x^2 + y) \in \mathbb{F}_2(y)[x]$, which factors as $(x - \sqrt{y})^2$, making the extension not separable.

8 Modules

Definition 8.0.1 (R-modules)
Four properties:

- $r(x + y) = rx + ry$
- $(r + s)x = rs + sx$
- $(rs)x = r(s(x))$
- $1_R x = x$

Proposition 8.0.2 ***(The one-step submodule test).***
$M \in$ R-Mod iff for every $r \in R$ and $x, y \in M$, we have $rx + y \in M$.

8.1 General Theory

Definition 8.1.1 (Module Morphisms)
A map $f : M \to N$ is a **morphism of modules** iff $f(rm + n) = rf(m) + f(n)$.

Proposition 8.1.2 ***(Recognizing direct sums).***
If $M_1, M_2 \leq M$ are submodules, then $M = M_1 \oplus M_2$ if the following conditions hold:

- $M_1 + M_2 = M$
- $M_1 \cap M_2 = 0$

Definition 8.1.3 (Simple modules)
A module is **simple** iff it has no nontrivial proper submodules.

Definition 8.1.4 (Indecomposable modules)
A module M is **decomposable** iff it admits a direct sum decomposition $M \cong M_1 \oplus M_2$ with $M_1, M_2 \neq 0$. An **indecomposable** module is defined in the obvious way.

Definition 8.1.5 (Cyclic modules)
A module M is **cyclic** if there exists a single generator $m \in M$ such that $M = mR := \langle m \rangle$.

8.2 Free and Projective Modules

Definition 8.2.1 (Free Module)
A **free** module M is a module satisfying any of the following conditions:

- A universal property: There is a set $\mathcal{B}$ and a set map $M \xrightarrow{\iota} \mathcal{B}$ such that every set map $\mathcal{B} \xrightarrow{N}$ lifts:



Link to Diagram

- Existence of a basis:

  There is linearly independent (so $\sum r_i\beta_i = 0 \implies r_i = 0$) spanning set (so $m \in M \implies m = \sum r_i\beta_i$) of the form $\mathcal{B} := \{\beta_i\}_{i \in I}$,

- Direct sum decomposition:

M decomposes as $M \cong \bigoplus_{i \in I} \beta_i R$, a sum of cyclic submodules.

Example 8.2.2 ***(A non-free module):*** $\mathbb{Z}/6$ is a $\mathbb{Z}$-module that is *not* free, since the element $[3]$ is a torsion element, where $2[3] = [6] = [0]$. This uses the fact that free modules over a PID are torsionfree.

Definition 8.2.3 (Free rank)
If a module M is free, the **free rank** of M is the cardinality of any basis.

Proposition 8.2.4 ***(?).***
Every free R-module admits a basis (spanning R-linearly independent set).

Definition 8.2.5 (Torsion and torsionfree)
An element $m \in M$ is a **torsion element** if there exists a nonzero $r \in R$ such that $rm = 0_M$. A module M is **torsion-free** if and only if for every $x \in M$, $mx = 0_M \implies m = 0_M$, i.e. M has no nonzero torsion elements. Equivalently, defining $M_t \coloneqq \left\{ m \in M \;\middle|\; \exists r \in R, rm = 0_M \right\}$ as the set of all torsion elements, M is torsion free iff $M_t = 0$. If $M_t = M$, we say M is a **torsion module**.

Proposition 8.2.6 ***(Free implies torsionfree (generally)).***
For R an integral domain, any free R-module M is torsionfree.

Prove

Example 8.2.7 ***(A torsionfree module that is not free):*** $\mathbb{Q} \in \mathbb{Z}\text{-Mod}$ is torsionfree, but not free as a $\mathbb{Z}$-module. This follows because any two elements $a/b, p/q$ are in a single ideal, since taking $d \coloneqq \gcd(b, q)$ we have $1/a = 1/d + \cdots 1/d$ and similarly $p/q = 1/a + \cdots + 1/a$, so these are in $\langle 1/d \rangle$. So any basis has size one, which would mean $\mathbb{Q} = \{\pm 1/d, \pm 2/d, \cdots\}$ which in particular doesn't include the average of the first two terms.

Definition 8.2.8 (Projective Modules)
A module P is **projective** iff it satisfies any of the following conditions:

- A universal property: for every surjective $N \xrightarrow{g} M$ and $P \xrightarrow{f} M$, the following lift exists:



Link to Diagram

- Direct summand:

  P is a direct summand of a free module F, so $F = P \oplus T$ for some module $T \leq F$.

Proposition 8.2.9 ***(Free implies projective).***
Any free $M \in$ R-Mod is projective.

Proof (?).

Todo: proof.

■

Example 8.2.10 ***(Projective $\not\Longrightarrow$ free):*** Let R_1, R_2 be two nontrivial rings and set $R \coloneqq R_1 \oplus R_2$. Then R_1, R_2 are projective R-modules by construction, but each factor contains R-torsion: setting $e \coloneqq (0,1) \in R$ we have $e \curvearrowright R_1 = 0_{R_1}$. Since free implies torsionfree, R_1 can not be a free R-module.

8.3 Exact Sequences

Definition 8.3.1 (Exact Sequences)
A sequence of R-module morphisms

$$0 \xrightarrow{d_1} A \xrightarrow{d_2} B \xrightarrow{d_3} C \to 0$$

is *exact* iff $\operatorname{im} d_i = \ker d_{i+1}$.

Definition 8.3.2 (Split Exact Sequences)
A short exact sequence

$$\xi : 0 \to A \xrightarrow{d_1} B \xrightarrow{d_2} C \to 0$$

has a **right-splitting** iff there exists a map $s : C \to B$ such that $d_2 \circ s = \mathbb{1}_C$. ξ has a **left-splitting** iff there exists a map $t : B \to A$ such that $t \circ d_1 = \mathbb{1}_A$.

Proposition 8.3.3 ***(Classifying split SESs).***
Let $\xi : 0 \to A \to B \to C \to 0$ be a SES, then TFAE

- ξ admits a right-splitting.
- ξ admits a left-splitting.
- ξ is isomorphic to a SES of the form $0 \to A \to A \oplus C \to C \to 0$.

Proposition 8.3.4 ***(Splitting Exact Sequences).***
A SES ξ splits if any of the following conditions hold:

- C is free.
- C is projective.
- A is injective.

8.4 Classification of Modules over a PID

Proposition 8.4.1 ***(STFGMPID).***
Let M be a finitely generated modules over a PID R. Then there is an invariant factor decomposition

$$M \cong F \bigoplus_{i=1}^{m} R/(r_i) \quad \text{where } r_1 \mid r_2 \mid \cdots$$

and similarly an elementary divisor decomposition:

Elementary divisor decomposition

Proposition 8.4.2 ***(Principal Ideals are Free).***
If $I \trianglelefteq R$ is an ideal of R, then I is a free R-module iff I is a principal ideal.

Proof (?).
$\implies$:
Suppose I is free as an R-module, and let $B = \{\mathbf{m}_j\}_{j\in J} \subseteq I$ be a basis so we can write $M = \langle B\rangle$. Suppose that $|B| \geq 2$, so we can pick at least 2 basis elements $\mathbf{m}_1 \neq \mathbf{m}_2$, and consider

$$\mathbf{c} = \mathbf{m}_1\mathbf{m}_2 - \mathbf{m}_2\mathbf{m}_1,$$

which is also an element of M . Since R is an integral domain, R is commutative, and so

$$\mathbf{c} = \mathbf{m}_1\mathbf{m}_2 - \mathbf{m}_2\mathbf{m}_1 = \mathbf{m}_1\mathbf{m}_2 - \mathbf{m}_1\mathbf{m}_2 = \mathbf{0}_M$$

However, this exhibits a linear dependence between $\mathbf{m}_1$ and $\mathbf{m}_2$, namely that there exist $\alpha_1, \alpha_2 \neq 0_R$ such that $\alpha_1\mathbf{m}_1 + \alpha_2\mathbf{m}_2 = \mathbf{0}_M$; this follows because $M \subset R$ means that we can take $\alpha_1 = -m_2, \alpha_2 = m_1$. This contradicts the assumption that B was a basis, so we must have $|B| = 1$ and so $B = \{\mathbf{m}\}$ for some $\mathbf{m} \in I$. But then $M = \langle B\rangle = \langle \mathbf{m}\rangle$ is generated by a single element, so M is principal.
$\impliedby$: Suppose $M \trianglelefteq R$ is principal, so $M = \langle \mathbf{m}\rangle$ for some $\mathbf{m} \neq \mathbf{0}_M \in M \subset R$.
Then $x \in M \implies x = \alpha\mathbf{m}$ for some element $\alpha \in R$ and we just need to show that $\alpha\mathbf{m} = \mathbf{0}_M \implies \alpha = 0_R$ in order for $\{\mathbf{m}\}$ to be a basis for M, making M a free R-module. But since $M \subset R$, we have $\alpha, m \in R$ and $\mathbf{0}_M = 0_R$, and since R is an integral domain, we have $\alpha m = 0_R \implies \alpha = 0_R$ or $m = 0_R$. Since $m \neq 0_R$, this forces $\alpha = 0_R$, which allows $\{m\}$ to be a linearly independent set and thus a basis for M as an R-module.
■

8.5 Algebraic Properties

Definition 8.5.1 (Module structure on tensor products)

$$r \curvearrowright (m \otimes n) \coloneqq (r \curvearrowright m) \otimes n.$$

Proposition 8.5.2 *(?).*
If $\dim_k V, \dim_k W < \infty$ then there is an isomorphism

$$\begin{aligned} V^{\vee} \otimes_k W &\xrightarrow{\sim} \underset{\text{k-Mod}}{\operatorname{Hom}}(V, W) \\ \tilde{v} \otimes w &\mapsto \tilde{v}(-)w. \end{aligned}$$

Proposition 8.5.3 *(?).*
If either of $\dim_k V, \dim_k W$ is finite, then

$$\begin{aligned} V^{\vee} \otimes_k W^{\vee} &\xrightarrow{\sim} (V \otimes W)^{\vee} \\ v \otimes w &\mapsto (x \otimes y \mapsto v(x)w(y)). \end{aligned}$$

Proposition 8.5.4 *(?).*

$$\begin{aligned} \underset{\text{k-Mod}}{\operatorname{Hom}}(V, W) &\xrightarrow{\sim} \underset{\text{k-Mod}}{\operatorname{Hom}}(W, V)^{\vee} \\ T &\mapsto \operatorname{Tr}(T \circ -). \end{aligned}$$

Proposition 8.5.5 *(?).*
If $T : V \hookrightarrow W$ is injective, then $T \otimes \mathbb{1}_X : V \otimes X \hookrightarrow W \otimes X$ is also injective for any X. Thus $F(-) = (- \otimes X)$ is right-exact for any X.

Example 8.5.6 ***(Computing tensor products):*** $\mathbb{Z}/2 \otimes_{\mathbb{Z}} \mathbb{Z}/3 = 0$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/3 & \xrightarrow{(-\times 2)\times \mathbb{1}} & \mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/3 & \longrightarrow & \mathbb{Z}/3 \otimes_{\mathbb{Z}} \mathbb{Z}/2 \\ & & \downarrow{\scriptstyle \mathrm{proj}_2} & & \downarrow{\scriptstyle \mathrm{proj}_2} & & \downarrow{\scriptstyle \cong} \\ 0 & \longrightarrow & \mathbb{Z}/3 & \xrightarrow{(-\times 2)} & \mathbb{Z}/3 & \longrightarrow & 0 \end{array}$$

Link to Diagram

9 | Linear Algebra

9.1 Definitions

Definition 9.1.1 (Invariant Factor)
Todo
todo

Definition 9.1.2 (Elementary Divisor)
Todo
todo

Definition 9.1.3 (Nondegenerate Bilinear Form)
todo

Definition 9.1.4 (Quadratic Form)
todo

Definition 9.1.5 (Gram Matrix)
todo

Definition 9.1.6 (Normal Matrix)
A matrix $A \in \mathrm{Mat}(n \times n; \mathbb{C})$ is **normal** iff $A^*A = AA^*$ where A^* is the conjugate transpose.

Proposition 9.1.7 *(?).*
Any linear map $T : V \to V$ decomposes as $T = D + N$ with D diagonal, N nilpotent, and $[DN] = 0$.

Proposition 9.1.8 *(?).*

$$\sum (W_i)^\perp = \int W_i^\perp.$$

9.2 Minimal / Characteristic Polynomials

Remark 9.2.1: Fix some notation:

$$\begin{aligned} \min_A(x): &\quad \text{The minimal polynomial of } A \\ \chi_A(x): &\quad \text{The characteristic polynomial of } A. \end{aligned}$$

Definition 9.2.2 (?)
The **minimal polynomial** of a linear morphism is the unique monic polynomial $\min_A(x)$ of minimal degree such that $\min_A(A) = 0$.

Definition 9.2.3 (?)
The **characteristic polynomial** of A is given by

$$\chi_A(x) = \det(A - xI)) = \det(SNF(A - xI)).$$

Fact 9.2.4
If A is upper triangular, then $\det(A) = \prod_i a_{ii}$

Theorem 9.2.5 ***(Cayley-Hamilton).***
The minimal polynomial divides the characteristic polynomial, and in particular $\chi_A(A) = 0$.

Proof (?).
By minimality, $\min_A$ divides χ_A. Every λ_i is a root of $\min_A(x)$: Let $(\mathbf{v}_i, \lambda_i)$ be a nontrivial eigenpair. Then by linearity,

$$\min_A(\lambda_i)\mathbf{v}_i = \min_A(A)\mathbf{v}_i = \mathbf{0},$$

which forces $\min_A(\lambda_i) = 0$.

■

Definition 9.2.6 (Similar Matrices)
Two matrices A, B are **similar** (i.e. $A = PBP^{-1}$) $\iff$ A, B have the same Jordan Canonical Form (JCF).

Definition 9.2.7 (Equivalent Matrices)
Two matrices A, B are **equivalent** (i.e. $A = PBQ$) $\iff$

- They have the same rank,
- They have the same invariant factors, *and*
- They have the same (JCF)

9.3 Finding Minimal Polynomials

Proposition 9.3.1 ***(How to find the minimal polynomial).***
Let $m(x)$ denote the minimal polynomial A.

1. Find the characteristic polynomial $\chi(x)$; this annihilates A by Cayley-Hamilton. Then $m(x) \mid \chi(x)$, so just test the finitely many products of irreducible factors.
2. Pick any $\mathbf{v}$ and compute $T\mathbf{v}, T^2\mathbf{v}, \cdots T^k\mathbf{v}$ until a linear dependence is introduced. Write this as $p(T) = 0$; then $\min_A(x) \mid p(x)$.

Definition 9.3.2 (Companion Matrix)
Given a monic $p(x) = a_0 + a_1 x + a_2 x^2 + \cdots + a_{n-1}x^{n-1} + x^n$, the **companion matrix** of p is given by

$$C_p \coloneqq \begin{bmatrix} 0 & 0 & \dots & 0 & -a_0 \\ 1 & 0 & \dots & 0 & -a_1 \\ 0 & 1 & \dots & 0 & -a_2 \\ \vdots & & \ddots & & \vdots \\ 0 & 0 & \dots & 1 & -a_{n-1} \end{bmatrix}.$$

9.4 Canonical Forms

Proposition 9.4.1 ***(?).***
Let $T : V \to V$ be a linear map where $n \coloneqq \dim_k V$. TFAE:

- There exists a basis $\{e_i\}$ of V such that

$$T(e_i) = \begin{cases} e_{i-1} & i \geq 2 \\ 0 & i = 1. \end{cases}$$

- There exists a cyclic vector $\mathbf{v}$ such that $\left\{T^k\mathbf{v} \;\middle|\; k = 1, 2, \cdots, n\right\}$ form a basis for V.
- $T^{n-1} \neq 0$
- $\dim_k \ker T^\ell = \ell$ for each $1 \leq \ell \leq n$.
- $\dim_k \ker T = 1$.

9.4.1 Rational Canonical Form

Corresponds to the **Invariant Factor Decomposition** of T.

Theorem 9.4.2 ***(Structure Theorem).***
For R a PID and M a finitely-generated R-module, there exists an invariant factor decomposition

$$M \cong R^r \bigoplus_{i=1}^{\ell} R/(a_i) \quad a_1 \mid a_2 \mid \cdots \mid a_\ell$$

where each a_i is an invariant factor.

Proposition 9.4.3 ***(RCG Relates to Invariant Factors).***
$RCF(A)$ is a block matrix where each block is the companion matrix of an invariant factor of A.

Proof (?).
The derivation:

- Let $k[x] \curvearrowright V$ using T, makes V into a $k[x]$-module.
- k a field implies $k[x]$ a PID, so apply structure theorem to obtain invariant factors a_i,
- Note that $T \curvearrowright V$ by multiplication by x
- Write $\bar{x} = \pi(x)$ where $F[x] \xrightarrow{\pi} F[x]/(a_i)$; then span $\{\overline{x}\} = F[x]/(a_i)$.
- Write $a_i(x) = \sum b_i x^i$, note that $V \to F[x]$ pushes $T \curvearrowright V$ to $T \curvearrowright k[x]$ by multiplication by $\overline{x}$
- WRT the basis $\overline{x}$, T then acts via the companion matrix on this summand.
- Each invariant factor corresponds to a block of the RCF.

■

9.4.2 Jordan Canonical Form

Corresponds to the **Elementary Divisor Decomposition** of T.

Lemma 9.4.4 ***(?).***
The elementary divisors of A are the minimal polynomials of the Jordan blocks.

Lemma 9.4.5 ***(JCF from Minimal and Characteristic Polynomials).***

Writing $\text{Spec}(A) = \{(\lambda_i, b_i)\}$,

$$\min_A(x) = \prod (x - \lambda_i)^{a_i}$$
$$\chi_A(x) = \prod (x - \lambda_i)^{b_i}$$

- The roots both polynomials are precisely the eigenvalues of A
- The spectrum of A corresponds precisely to the **characteristic** polynomial
- $a_i \leq b_i$
- a_i is the size of the **largest** Jordan block associated to λ_i,
- b_i is the **sum of sizes** of all Jordan blocks associated to λ_i and the number of times λ_i appears on the diagonal of $JCF(A)$.
- $\dim E_{\lambda_i}$ is the **number of Jordan blocks** associated to λ_i

9.4.3 Finding Possible Canonical Forms

Show how to find RCF and JCF from eigenvalues, or minimal/char polynomials.

9.4.4 Using Canonical Forms

Lemma 9.4.6 *(?).*
The characteristic polynomial is the *product of the invariant factors*, i.e.

$$\chi_A(x) = \prod_{j=1}^{n} f_j(x).$$

Lemma 9.4.7 *(?).*
The minimal polynomial of A is the *invariant factor of highest degree*, i.e.

$$\min_A(x) = f_n(x).$$

Proposition 9.4.8 *(?).*
For a linear operator on a vector space of nonzero finite dimension, TFAE:

- The minimal polynomial is equal to the characteristic polynomial.
- The list of invariant factors has length one.
- The Rational Canonical Form has a single block.
- The operator has a matrix similar to a companion matrix.

- There exists a *cyclic vector* $\mathbf{v}$ such that $\operatorname{span}_k \left\{ T^j \mathbf{v} \;\middle|\; j = 1, 2, \cdots \right\} = V$.
- T has $\dim V$ distinct eigenvalues

9.5 Diagonalizability

Remark 9.5.1: *Notation:* A^* denotes the conjugate transpose of A.

Lemma 9.5.2 ***(?).***
Let V be a vector space over k an algebraically closed and $A \in \operatorname{End}(V)$. Then if $W \subseteq V$ is an invariant subspace, so $A(W) \subseteq W$, the A has an eigenvector in W.

Theorem 9.5.3 ***(The Spectral Theorem).***

1. Hermitian matrices (i.e. $A^* = A$) are diagonalizable over $\mathbb{C}$.
2. Symmetric matrices (i.e. $A^t = A$) are diagonalizable over $\mathbb{R}$.

Proof (?).

- Suppose A is Hermitian.
- Since V itself is an invariant subspace, A has an eigenvector $\mathbf{v}_1 \in V$.
- Let $W_1 = \operatorname{span}_k \{\mathbf{v}_1\} \perp$.
- Then for any $\mathbf{w}_1 \in W_1$,

$$\langle \mathbf{v}_1,\ A\mathbf{w}_1 \rangle = \langle A\mathbf{v}_1,\ \mathbf{w}_1 \rangle = \lambda \langle \mathbf{v}_1,\ \mathbf{w}_1 \rangle = 0,$$

  so $A(W_1) \subseteq W_1$ is an invariant subspace, etc.
- Suppose now that A is symmetric.
- Then there is an eigenvector of norm 1, $\mathbf{v} \in V$.

$$\lambda = \lambda \langle \mathbf{v},\ \mathbf{v} \rangle = \langle A\mathbf{v},\ \mathbf{v} \rangle = \langle \mathbf{v},\ A\mathbf{v} \rangle = \overline{\lambda} \implies \lambda \in \mathbb{R}.$$

■

Proposition 9.5.4 ***(Simultaneous Diagonalizability).***
A set of operators $\{A_i\}$ pairwise commute $\iff$ they are all simultaneously diagonalizable.

Proof (?).
By induction on number of operators

- A_n is diagonalizable, so $V = \bigoplus E_i$ a sum of eigenspaces
- Restrict all $n-1$ operators A to E_n.
- The commute in V so they commute in E_n
- **(Lemma)** They were diagonalizable in V, so they're diagonalizable in E_n
- So they're simultaneously diagonalizable by I.H.
- But these eigenvectors for the A_i are all in E_n, so they're eigenvectors for A_n too.
- Can do this for each eigenspace.

Full details here

■

Theorem 9.5.5 ***(Characterizations of Diagonalizability).***
M is diagonalizable over $\mathbb{F} \iff \min_M(x, \mathbb{F})$ splits into distinct linear factors over $\mathbb{F}$, or equivalently iff all of the roots of $\min_M$ lie in $\mathbb{F}$.

Proof (?).
$\implies$: If $\min_A$ factors into linear factors, so does each invariant factor, so every elementary divisor is linear and $JCF(A)$ is diagonal.
$\impliedby$: If A is diagonalizable, every elementary divisor is linear, so every invariant factor factors into linear pieces. But the minimal polynomial is just the largest invariant factor.

■

9.6 Matrix Counterexamples

Example 9.6.1 ***(?):***

Example 9.6.2 ***(?):*** A matrix that:

- Is not diagonalizable over $\mathbb{R}$ but diagonalizable over $\mathbb{C}$
- Has *no* eigenvalues over $\mathbb{R}$ but has *distinct* eigenvalues over $\mathbb{C}$
- $\min_M(x) = \chi_M(x) = x^2 + 1$

$$M = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \sim \left(\begin{array}{c|c} -1\sqrt{-1} & 0 \\ \hline 0 & 1\sqrt{-1} \end{array}\right).$$

Example 9.6.3 ***(?):*** A matrix that:

- Is not diagonalizable over $\mathbb{C}$,

- Has eigenvalues $[1, 1]$ (repeated, multiplicity 2)
- $\min_M(x) = \chi_M(x) = x^2 - 2x + 1$

$$M = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Example 9.6.4 *(?):* Non-similar matrices with the same characteristic polynomial

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

Example 9.6.5 *(?):* A full-rank matrix that is not diagonalizable:

$$\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

Example 9.6.6 *(?):* Matrix roots of unity, i.e. representations of i:

$$M_1 \coloneqq \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \quad M_2 \coloneqq \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

9.7 Matrix Groups

Definition 9.7.1 (General Linear Group)

$$\mathrm{GL}_n(\mathbb{R}) = \left\{ A \;\middle|\; A = \overline{A} \right\}.$$

:::{.definition title="Special Linear Group"}

$$\mathrm{SL}_n(\mathbb{C}) \coloneqq \left\{ A \;\middle|\; \det A = 1 \right\}.$$

::

Definition 9.7.2 (Orthogonal Group)

$$O_n(\mathbb{C}) \coloneqq \left\{ A \;\middle|\; A^t A = A A^t = I \right\}.$$

Dimension: $n(n-1)/2$.

Definition 9.7.3 (Special Orthogonal Group)

$$\mathrm{SO}_n(\mathbb{R}) = \left\{ A \;\middle|\; AA^t = I \right\} = \ker(\mathrm{GL}_n(\mathbb{R}) \to k^\times).$$

Definition 9.7.4 (Unitary Group)

$$U_n(\mathbb{C}) \coloneqq \left\{ A \;\middle|\; A^\dagger A = AA^\dagger = 1 \right\}.$$

Definition 9.7.5 (Special Unitary Group)

$$\mathrm{SU}_n(\mathbb{C}) \coloneqq \left\{ A \in U_n(\mathbb{C}) \;\middle|\; \det A = 1 \right\}.$$

Definition 9.7.6 (Symplectic Group)

Matrix group definitions.

Proposition 9.7.7 ***(Order of*** GL_n***).***

todo

10 Representation Theory

Theorem 10.0.1 ***(Schur's Lemma).***
If $M \in$ G-Mod is an irreducible representation of G with $\dim_k M < \infty$ and $k = \bar{k}$, then there is an isomorphism

$$M \xrightarrow{\sim} \mathrm{Aut}_G(M, M).$$

Theorem 10.0.2 ***(Maschke's Theorem).***
Let k be a field with $\mathrm{ch}(k)$ not dividing $\#G$. Then any finite-dimensional representation of G decomposes into a direct sum of irreducible representations.

Definition 10.0.3 (Characters)

The **character** of a representation M is the trace of the map

$$T_g : M \to M$$
$$m \mapsto g \curvearrowright m.$$

11 | Extra Problems

11.1 Commutative Algebra

- Show that a finitely generated module over a Noetherian local ring is flat iff it is free using Nakayama and Tor.
- Show that $\langle 2, x \rangle \trianglelefteq \mathbb{Z}[x]$ is not a principal ideal.
- Let R be a Noetherian ring and A, B algebras over R. Suppose A is finite type over R and finite over B. Then B is finite type over R.

11.2 Group Theory

11.2.1 Basic Structure

Just Structure

- Show that the intersection of two subgroups is again a subgroup.
- Show that the intersection of two subgroups with coprime orders is trivial.
- Show that subgroups with the *same* prime order are either equal or intersect trivially.
- Give a counterexample where $H, K \leq G$ but HK is not a subgroup of G.
- Show that $G = H \times K$ iff the conditions for recognizing direct products hold.
- Show that if $H, K \trianglelefteq G$ and $H \cap K = \emptyset$, then $hk = kh$ for all $h \in H, k \in K$.
- Show that if $H, K \trianglelefteq G$ are normal subgroups that intersect trivially, then $[H, K] = 1$ (so $hk = kh$ for all k and h).
- Show that the order of any element in a group divides the order of the group.
- Show that $|G|/|H| = [G : H]$.

Centers

- Show that if $G/Z(G)$ is cyclic then G is abelian.
- Show that G/N is abelian iff $[G, G] \leq N$.

- Show that every normal subgroup of G is contained in $Z(G)$.

Cyclic Groups

- Show that any cyclic group is abelian.
- Show that every subgroup of a cyclic group is cyclic.
- Show that

$$\varphi(n) = n \prod_{p \Big| n} \left(1 - \frac{1}{p}\right).$$

- Compute $\mathrm{Aut}(\mathbb{Z}/n\mathbb{Z})$ for n composite.
- Compute $\mathrm{Aut}((\mathbb{Z}/p\mathbb{Z})^n)$.

Conjugacy

- Show that the size of a conjugacy class divides the order of a group.
- Let G be finite with $H \leq G$ and show that G is not the union of the conjugates of H.

Hint: consider the intersection and count.

Hint: Orbit-stabilizer

11.2.2 Centralizing and Normalizing

- Show that $C_G(H) \subseteq N_G(H) \leq G$.
- Show that $Z(G) \subseteq C_G(H) \subseteq N_G(H)$.
- Given $H \subseteq G$, let $S(H) = \bigcup_{g \in G} gHg^{-1}$, so $|S(H)|$ is the number of conjugates to H. Show that $|S(H)| = [G : N_G(H)]$.
  - That is, the number of subgroups conjugate to H equals the index of the normalizer of H.
- Show that $Z(G) = \bigcap_{a \in G} C_G(a)$.
- Show that the centralizer $G_G(H)$ of a subgroup is again a subgroup.
- Show that $C_G(H) \trianglelefteq N_G(H)$ is a normal subgroup.
- Show that $C_G(G) = Z(G)$.

- Show that for $H \leq G$, $C_H(x) = H \cap C_G(x)$.
- Let $H, K \leq G$ a finite group, and without using the normalizers of H or K, show that $|HK| = |H||K|/|H \cap K|$.
- Show that if $H \leq N_G(K)$ then $HK \leq H$, and give a counterexample showing that this condition is necessary.
- Show that HK is a subgroup of G iff $HK = KH$.
- Prove that the kernel of a homomorphism is a normal subgroup.

11.2.3 Primes in Group Theory

- Show that any group of prime order is cyclic and simple.
- Analyze groups of order pq with $q < p$.

  Hint: consider the cases when p does or does not divide $q - 1$.

  - Show that if q does not divide $p - 1$, then G is cyclic.
  - Show that G is never simple.

- Analyze groups of order p^2q.

  Hint: Consider the cases when q does or does not divide $p^2 - 1$.

- Show that no group of order p^2q^2 is simple for $p < q$ primes.
- Show that a group of order p^2q^2 has a normal Sylow subgroup.
- Show that a group of order p^2q^2 where q does not divide $p^2 - 1$ and p does not divide $q^2 - 1$ is abelian.
- Show that every group of order pqr with $p < q < r$ primes contains a normal Sylow subgroup.

  - Show that G is never simple.

- Let p be a prime and $|G| = p^3$. Prove that G has a normal subgroup N of order p^2.

  - Suppose $N = \langle h \rangle$ is cyclic and classify all possibilities for G if:
    - ♢ $|h| = p^3$
    - ♢ $|h| = p$.

Hint: Sylow and semidirect products.

- Show that any normal p- subgroup is contained in every Sylow p-subgroup of G.
- Show that the order of $1+p$ in $\left(\mathbb{Z}/p^2\mathbb{Z}\right)^\times$ is equal to p. Use this to construct a non-abelian group of order p^3.

11.2.4 p-Groups

- Show that every p-group has a nontrivial center.
- Show that every p-group is nilpotent.
- Show that every p-group is solvable.
- Show that every maximal subgroup of a p-group has index p.
- Show that every maximal subgroup of a p-group is normal.
- Show that every group of order p is cyclic.
- Show that every group of order p^2 is abelian and classify them.
- Show that every normal subgroup of a p-group is contained in the center.

Hint: Consider $G/Z(G)$.

- Let $O_P(G)$ be the intersection of all Sylow p-subgroups of G. Show that $O_p(G) \trianglelefteq G$, is maximal among all normal p-subgroups of G
- Let $P \in \mathrm{Syl}_p(H)$ where $H \trianglelefteq G$ and show that $P \cap H \in \mathrm{Syl}_p(H)$.
- Show that Sylow p_i-subgroups S_{p_1}, S_{p_2} for distinct primes $p_1 \neq p_2$ intersect trivially.
- Show that in a p group, every normal subgroup intersects the center nontrivially.

11.2.5 Symmetric Groups

Specific Groups

- Show that the center of S_3 is trivial.
- Show that $Z(S_n) = 1$ for $n \geq 3$
- Show that $\mathrm{Aut}(S_3) = \mathrm{Inn}(S_3) \cong S_3$.

- Show that the transitive subgroups of S_3 are S_3, A_3
- Show that the transitive subgroups of S_4 are $S_4, A_4, D_4, \mathbb{Z}_2^2, \mathbb{Z}_4$.
- Show that S_4 has two normal subgroups: $A_4, \mathbb{Z}_2^2$.
- Show that $S_{n\geq 5}$ has one normal subgroup: A_n.
- $Z(A_n) = 1$ for $n \geq 4$
- Show that $[S_n, S_n] = A_n$
- Show that $[A_4, A_4] \cong \mathbb{Z}_2^2$
- Show that $[A_n, A_n] = A_n$ for $n \geq 5$, so $A_{n\geq 5}$ is nonabelian.

General Structure

- Show that an m-cycle is an odd permutation iff m is an even number.
- Show that a permutation is odd iff it has an odd number of even cycles.
- Show that the center of S_n for $n \geq 4$ is nontrivial.
- Show that disjoint cycles commute.
- Show directly that any k-cycle is a product of transpositions, and determine how many transpositions are needed.

Generating Sets

- Show that S_n is generated by any of the following types of cycles:

Group	Generating Set	Size
S_n, $n \geq 2$	(ij)'s	$\frac{n(n-1)}{2}$
	$(12), (13), \ldots, (1n)$	$n-1$
	$(12), (23), \ldots, (n-1\ n)$	$n-1$
	$(12), (12\ldots n)$ if $n \geq 3$	2
	$(12), (23\ldots n)$ if $n \geq 3$	2
	$(ab), (12\ldots n)$ if $(b-a, n) = 1$	2
A_n, $n \geq 3$	3-cycles	$\frac{n(n-1)(n-2)}{3}$
	$(1ij)$'s	$(n-1)(n-2)$
	$(12i)$'s	$n-2$
	$(i\ i+1\ i+2)$'s	$n-2$
	$(123), (12\ldots n)$ if $n \geq 4$ odd	2
	$(123), (23\ldots n)$ if $n \geq 4$ even	2

  - Show that S_n is generated by transpositions.
  - Show that S_n is generated by *adjacent* transpositions.
  - Show that S_n is generated by $\{(12), (12\cdots n)\}$ for $n \geq 2$
  - Show that S_n is generated by $\{(12), (23\cdots n)\}$ for $n \geq 3$
  - Show that S_n is generated by $\{(ab), (12\cdots n)\}$ where $1 \leq a < b \leq n$ iff $\gcd(b-a, n) = 1$.

- Show that S_p is generated by any arbitrary transposition and any arbitrary p-cycle.

11.2.6 Alternating Groups

- Show that A_n is generated 3-cycles.
- Prove that A_n is normal in S_n.
- Argue that A_n is simple for $n \geq 5$.
- Show that $\mathrm{Out}(A_4)$ is nontrivial.

11.2.7 Dihedral Groups

- Show that if $N \trianglelefteq D_n$ is a normal subgroup of a dihedral group, then D_n/N is again a dihedral group.

11.2.8 Other Groups

- Show that $\mathbb{Q}$ is not finitely generated as a group.
- Show that the Quaternion group has only one element of order 2, namely -1.

11.2.9 Classification

- Show that no group of order 36 is simple.
- Show that no group of order 90 is simple.
- Classifying all groups of order 99.
- Show that all groups of order 45 are abelian.
- Classify all groups of order 10.
- Classify the five groups of order 12.
- Classify the four groups of order 28.
- Show that if $|G| = 12$ and has a normal subgroup of order 4, then $G \cong A_4$.
- Suppose $|G| = 240 = s^4 \cdot 3 \cdot 5$.
  - How many Sylow-p subgroups does G have for $p \in \{2, 3, 5\}$?
  - Show that if G has a subgroup of order 15, it has an element of order 15.
  - Show that if G does not have such a subgroup, the number of Sylow-3 subgroups is either 10 or 40.

Hint: Sylow on the subgroup of order 15 and semidirect products.

11.2.10 Group Actions

- Show that the stabilizer of an element G_x is a subgroup of G.
- Show that if x, y are in the same orbit, then their stabilizers are conjugate.
- Show that the stabilizer of an element need not be a normal subgroup?
- Show that if $G \curvearrowright X$ is a group action, then the stabilizer G_x of a point is a subgroup.

11.2.11 Series of Groups

- Show that A_n is simple for $n \geq 5$
- Give a necessary and sufficient condition for a cyclic group to be solvable.
- Prove that every simple abelian group is cyclic.
- Show that S_n is generated by disjoint cycles.
- Show that S_n is generated by transpositions.
- Show if G is finite, then G is solvable $\iff$ all of its composition factors are of prime order.
- Show that if N and G/N are solvable, then G is solvable.
- Show that if G is finite and solvable then every composition factor has prime order.
- Show that G is solvable iff its derived series terminates.
- Show that S_3 is not nilpotent.
- Show that G nilpotent $\implies$ G solvable
- Show that nilpotent groups have nontrivial centers.
- Show that Abelian $\implies$ nilpotent
- Show that p-groups $\implies$ nilpotent

11.2.12 Misc

- Prove Burnside's theorem.
- Show that $\text{Inn}(G) \trianglelefteq Aut(G)$
- Show that $\text{Inn}(G) \cong G/Z(G)$

- Show that the kernel of the map $G \to \text{Aut}(G)$ given by $g \mapsto (h \mapsto ghg^{-1})$ is $Z(G)$.
- Show that $N_G(H)/C_G(H) \cong A \leq Aut(H)$
- Give an example showing that normality is not transitive: i.e. $H \trianglelefteq K \trianglelefteq G$ with H *not* normal in G.

11.2.13 Nonstandard Topics

- Show that H char $G \Rightarrow H \trianglelefteq G$

Thus "characteristic" is a strictly stronger condition than normality

- Show that H char K char $G \Rightarrow H$ char G

So "characteristic" is a transitive relation for subgroups.

- Show that if $H \leq G$, $K \trianglelefteq G$ is a normal subgroup, and H char K then H is normal in G.

So normality is not transitive, but strengthening one to "characteristic" gives a weak form of transitivity.

12

12.1 Ring Theory

Basic Structure

- Show that if an ideal $I \trianglelefteq R$ contains a unit then $I = R$.
- Show that $R^\times$ need not be closed under addition.

Ideals

Problem 12.1.1 (Units or Zero Divisors)
Every $a \in R$ for a finite ring is either a unit or a zero divisor.

Solution:

- Let $a \in R$ and define $\varphi(x) = ax$.
- If φ is injective, then it is surjective, so $1 = ax$ for some $x \implies x^{-1} = a$.
- Otherwise, $ax_1 = ax_2$ with $x_1 \neq x_2 \implies a(x_1 - x_2) = 0$ and $x_1 - x_2 \neq 0$
- So a is a zero divisor.

Problem 12.1.2 (Maximal implies prime)
Maximal $\implies$ prime, but generally not the converse.

Solution:
- Suppose $\mathfrak{m}$ is maximal, $ab \in \mathfrak{m}$, and $b \notin \mathfrak{m}$.
- Then there is a containment of ideals $\mathfrak{m} \subsetneq \mathfrak{m} + (b) \implies \mathfrak{m} + (b) = R$.
- So

$$1 = m + rb \implies a = am + r(ab),$$

but $am \in \mathfrak{m}$ and $ab \in \mathfrak{m} \implies a \in \mathfrak{m}$.

Counterexample: $(0) \in \mathbb{Z}$ is prime since $\mathbb{Z}$ is a domain, but not maximal since it is properly contained in any other ideal.

- Show that every proper ideal is contained in a maximal ideal
- Show that if $x \in R$ a PID, then x is irreducible $\iff \langle x \rangle \trianglelefteq R$ is maximal.
- Show that intersections, products, and sums of ideals are ideals.
- Show that the union of two ideals need not be an ideal.
- Show that every ring has a proper maximal ideal.
- Show that $I \trianglelefteq R$ is maximal iff R/I is a field.
- Show that $I \trianglelefteq R$ is prime iff R/I is an integral domain.
- Show that $\cup_{\mathfrak{m} \in \operatorname{maxSpec}(R)} = R \setminus R^\times$.
- Show that $\operatorname{maxSpec}(R) \subsetneq \operatorname{Spec}(R)$ but the containment is strict.
- $\star$ Show that if x is not a unit, then x is contained in some maximal ideal.
- Show that every prime ideal is radical.
- Show that the nilradical is given by $\mathfrak{N}(R) = \operatorname{rad}(0)$.
- Show that $\operatorname{rad}(IJ) = \operatorname{rad}(I) \cap \operatorname{rad}(J)$
- Show that if $\operatorname{Spec}(R) \subseteq \operatorname{maxSpec}(R)$ then R is a UFD.
- Show that if R is Noetherian then every ideal is finitely generated.

Characterizing Certain Ideals

- Show that the nilradical of a ring is the intersection of all prime ideals $I \trianglelefteq R$.
- Show that for an ideal $I \trianglelefteq R$, its radical is the intersection of all prime ideals containing I.
- Show that $\operatorname{rad}(I)$ is the intersection of all prime ideals containing I.

Problem 12.1.3 (Jacobson radical is bigger than the nilradical)
The nilradical is contained in the Jacobson radical, i.e.

$$\mathfrak{N}(R) \subseteq \mathfrak{J}(R).$$

Solution:
Maximal $\implies$ prime, and so if x is in every prime ideal, it is necessarily in every maximal ideal as well.

Problem 12.1.4 (Mod by nilradical to kill nilpotents)
$R/\mathfrak{N}(R)$ has no nonzero nilpotent elements.

Solution:

$$\begin{aligned} a+\mathfrak{N}(R) \text{ nilpotent} &\implies (a+\mathfrak{N}(R))^n := a^n + \mathfrak{N}(R) = \mathfrak{N}(R) \\ &\implies a^n \in \mathfrak{N}(R) \\ &\implies \exists \ell \text{ such that } (a^n)^\ell = 0 \\ &\implies a \in \mathfrak{N}(R). \end{aligned}$$

Problem 12.1.5 (Nilradical is intersection of primes)
The nilradical is the intersection of all prime ideals, i.e.

$$\mathfrak{N}(R) = \cap_{\mathfrak{p}\in \operatorname{Spec}(R)} \mathfrak{p}$$

Solution: • $\mathfrak{N} \subseteq \cap \mathfrak{p}$:

- $x \in \mathfrak{N} \implies x^n = 0 \in \mathfrak{p} \implies x \in \mathfrak{p}$ or $x^{n-1} \in \mathfrak{p}$.
- $\mathfrak{N}^c \subseteq \cup \mathfrak{p}^c$:
- Define $S = \left\{ I \trianglelefteq R \;\middle|\; a^n \notin I \text{ for any } n \right\}$.
- Then apply Zorn's lemma to get a maximal ideal $\mathfrak{m}$, and maximal $\implies$ prime.

Misc

- Show that localizing a ring at a prime ideal produces a local ring.
- Show that R is a local ring iff for every $x \in R$, either x or $1-x$ is a unit.
- Show that if R is a local ring then $R \setminus R^\times$ is a proper ideal that is contained in $\mathfrak{J}(R)$.
- Show that if $R \neq 0$ is a ring in which every non-unit is nilpotent then R is local.
- Show that every prime ideal is primary.
- Show that every prime ideal is irreducible.
- Show that

12.2 Field Theory

General Algebra

- Show that any finite integral domain is a field.
- Show that every field is simple.
- Show that any field morphism is either 0 or injective.
- Show that if L/F and α is algebraic over both F and L, then the minimal polynomial of α over L divides the minimal polynomial over F.
- Prove that if R is an integral domain, then $R[t]$ is again an integral domain.
- Show that $ff(R[t]) = ff(R)(t)$.
- Show that $[\mathbb{Q}(\sqrt{2}+\sqrt{3}) : \mathbb{Q}] = 4$.
  - Show that $\mathbb{Q}(\sqrt{2}+\sqrt{3}) = \mathbb{Q}(\sqrt{2}-\sqrt{3}) = \mathbb{Q}(\sqrt{2},\sqrt{3})$.
- Show that the splitting field of $f(x) = x^3 - 2$ is $\mathbb{Q}(\sqrt[3]{2}, \zeta_2)$.

Extensions?

- What is $[\mathbb{Q}(\sqrt{2}+\sqrt{3}) : \mathbb{Q}]$?
- What is $[\mathbb{Q}(2^{\frac{3}{2}}) : \mathbb{Q}]$?
- Show that if $p \in \mathbb{Q}[x]$ and $r \in \mathbb{Q}$ is a rational root, then in fact $r \in \mathbb{Z}$.
- If $\{\alpha_i\}_{i=1}^n \subset F$ are algebraic over K, show that $K[\alpha_1, \cdots, \alpha_n] = K(\alpha_1, \cdots, \alpha_n)$.
- Show that α/F is algebraic $\iff$ $F(\alpha)/F$ is a finite extension.
- Show that every finite field extension is algebraic.
- Show that if α, β are algebraic over F, then $\alpha \pm \beta, \alpha\beta^{\pm 1}$ are all algebraic over F.
- Show that if $L/K/F$ with K/F algebraic and L/K algebraic then L is algebraic.

Special Polynomials

- Show that a field with p^n elements has exactly one subfield of size p^d for every d dividing n.
- Show that $x^{p^n} - x = \prod f_i(x)$ over all irreducible monic f_i of degree d dividing n.
- Show that $x^{p^d} - x \mid x^{p^n} - x \iff d \mid n$
- Prove that $x^{p^n} - x$ is the product of all monic irreducible polynomials in $\mathbb{F}_p[x]$ with degree dividing n.
- Prove that an irreducible $\pi(x) \in \mathbb{F}_p[x]$ divides $x^{p^n} - x \iff \deg \pi(x)$ divides n.

12.3 Galois Theory

12.3.1 Theory

- Show that if K/F is the splitting field of a separable polynomial then it is Galois.
- Show that any quadratic extension of a field F with $\mathrm{ch}(F) \neq 2$ is Galois.
- Show that if $K/E/F$ with K/F Galois then K/E is always Galois with $g(K/E) \leq g(K/F)$.
  - Show additionally E/F is Galois $\iff$ $g(K/E) \trianglelefteq g(K/F)$.
  - Show that in this case, $g(E/F) = g(K/F)/g(K/E)$.
- Show that if $E/k, F/k$ are Galois with $E \cap F = k$, then EF/k is Galois and $G(EF/k) \cong G(E/k) \times G(F/k)$.

12.3.2 Computations

- Show that the Galois group of $x^n - 2$ is D_n, the dihedral group on n vertices.
- Compute all intermediate field extensions of $\mathbb{Q}(\sqrt{2}, \sqrt{3})$, show it is equal to $\mathbb{Q}(\sqrt{2} + \sqrt{3})$, and find a corresponding minimal polynomial.



- Compute all intermediate field extensions of $\mathbb{Q}(2^{\frac{1}{4}}, \zeta_8)$.
- Show that $\mathbb{Q}(2^{\frac{1}{3}})$ and $\mathbb{Q}(\zeta_3 2^{\frac{1}{3}})$
- Show that if L/K is separable, then L is normal $\iff$ there exists a polynomial $p(x) = \prod_{i=1}^{n} x - \alpha_i \in K[x]$ such that $L = K(\alpha_1, \cdots, \alpha_n)$ (so L is the splitting field of p).
- Is $\mathbb{Q}(2^{\frac{1}{3}})/\mathbb{Q}$ normal?
- Show that $\mathbb{GF}(p^n)$ is the splitting field of $x^{p^n} - x \in \mathbb{F}_p[x]$.
- Show that $\mathbb{GF}(p^d) \leq \mathbb{GF}(p^n) \iff d \mid n$
- Compute the Galois group of $x^n - 1 \in \mathbb{Q}[x]$ as a function of n.
- Identify all of the elements of the Galois group of $x^p - 2$ for p an odd prime (note: this has a complicated presentation).
- Show that $\mathrm{Gal}(x^{15} + 2)/\mathbb{Q} \cong S_2 \rtimes \mathbb{Z}/15\mathbb{Z}$ for S_2 a Sylow 2-subgroup.
- Show that $\mathrm{Gal}(x^3 + 4x + 2)/\mathbb{Q} \cong S_3$, a symmetric group.

12.4 Modules and Linear Algebra

- Prove the Cayley-Hamilton theorem.
- Prove that the minimal polynomial divides the characteristic polynomial.
- Prove that the cokernel of $A \in \mathrm{Mat}(n \times n, \mathbb{Z})$ is finite $\iff \det A \neq 0$, and show that in this case $|\mathrm{coker}(A)| = |\det(A)|$.
- Show that a nilpotent operator is diagonalizable.
- Show that if A, B are diagonalizable and $[A, B] = 0$ then A, B are simultaneously diagonalizable.
- Does diagonalizable imply invertible? The converse?
- Does diagonalizable imply distinct eigenvalues?

- Show that if a matrix is diagonalizable, its minimal polynomial is squarefree.
- Show that a matrix representing a linear map $T : V \to V$ is diagonalizable iff V is a direct sum of eigenspaces $V = \bigoplus_i \ker(T - \lambda_i I)$.
- Show that if $\{\mathbf{v}_i\}$ is a basis for V where $\dim(V) = n$ and $T(\mathbf{v}_i) = \mathbf{v}_{i+1 \pmod n}$ then T is diagonalizable with minimal polynomial $x^n - 1$.
- Show that if the minimal polynomial of a linear map T is irreducible, then every T-invariant subspace has a T-invariant complement.

12.5 Linear Algebra

Sort out from module section.

13 Even More Algebra Questions

Remark 13.0.1: (DZG): These all come from a random PDF I found, but I couldn't find the original author/source!

13.1 Groups

13.1.1 Question 1.1

What is a normal subgroup? Can you get some natural map from a normal subgroup? What topological objects can the original group, normal subgroup, and quotient group relate to?

13.1.2 Question 1.2

Prove that a subgroup of index two is normal.

13.1.3 Question 1.3

Find all normal subgroups of A_4.

13.1.4 Question 1.4

Give an interesting example of a non-normal subgroup. Is SO(2) normal inside $SL_2(R)$?

13.1.5 Question 1.5

Is normality transitive? That is, is a normal subgroup of a normal subgroup normal in the biggest group?

13.1.6 Question 1.6.

Define a solvable group. Give an example of a solvable nonabelian group.

Show A_4 is solvable. Do the Sylow theorems tell you anything about whether this index 3 subgroup of A_4 is normal?

13.1.7 Question 1.7

Define lower central series, upper central series, nilpotent and solvable groups.

13.1.8 Question 1.8

Define the derived series. Define the commutator. State and prove two nontrivial theorems about derived series.

13.1.9 Question 1.9

Prove that $SL_2(Z)$ is not solvable.

13.1.10 Question 1.10

What are all possible orders of elements of $\mathrm{SL}_2(Z)$?

13.1.11 Question 1.11

Can you show that all groups of order p^n for p prime are solvable? Do you know how to do this for groups of order $p^r q^s$?

13.1.12 Question 1.12

Suppose a p-group acts on a set whose cardinality is not divisible by p (p prime). Prove that there is a fixed point for the action.

13.1.13 Question 1.13

Prove that the centre of a group of order pr (p prime) is not trivial.

13.1.14 Question 1.14

Give examples of simple groups. Are there infinitely many?

13.1.15 Question 1.15

State and prove the Jordan-Holder theorem for finite groups.

13.1.16 Question 1.16

What's Cayley's theorem? Give an example of a group of order n that embeds in S_m for some m smaller than n.

Give an example of a group where you have to use S_n.

13.1.17 Question 1.17

Is A_4 a simple group? What are the conjugacy classes in S_4? What about in A_4?

13.1.18 Question 1.18

Talk about conjugacy classes in the symmetric group S_n.

13.1.19 Question 1.19

When do conjugacy classes in S_n split in A_n?

13.1.20 Question 1.20

What is the centre of S_n? Prove it.

13.1.21 Question 1.21

Prove that the alternating group A_n is simple for $n \geq 5$.

13.1.22 Question 1.22

Prove the alternating group on n letters is generated by the 3-cycles for $n \geq 3$.

13.1.23 Question 1.23

Prove that for p prime, Sp is generated by a p-cycle and a transposition.

13.1.24 Question 1.24

What is the symmetry group of a tetrahedron? Cube? Icosahedron?

13.1.25 Question 1.25

How many ways can you color the tetrahedron with C colors if we identify symmetric colorings?

13.1.26 Question 1.26.

What is the symmetry group of an icosahedron? What's the stabiliser of an edge?

How many edges are there? How do you know the symmetry group of the icosahedron is the same as the symmetry group of the dodecahedron?

Do you know the classification of higher-dimensional polyhedra?

13.1.27 Question 1.27

Do you know what the quaternion group is? How many elements are there of each order?

13.1.28 Question 1.28

What is the group of unit quaternions topologically? What does it have to do with SO(3)?

13.1.29 Question 1.29

What's the stabiliser of a point in the unit disk under the group of conformal automorphisms?

13.1.30 Question 1.30

What group-theoretic construct relates the stabiliser of two points?

13.1.31 Question 1.31

Consider $\mathrm{SL}_2(R)$ acting on $\mathbb{R}^2$ by matrix multiplication. What is the stabiliser of a point? Does it depend which point? Do you know what sort of subgroup this is? What if $\mathrm{SL}_2(R)$ acts by Möbius transformations instead?

13.1.32 Question 1.32

What are the polynomials in two real variables that are invariant under the action of D_4, the symmetry group of a square, by rotations and reflections on the plane that the two variables form?

13.1.33 Question 1.33

Give an interesting example of a subgroup of the additive group of the rationals.

13.1.34 Question 1.34

Talk about the isomorphism classes of subgroups of $\mathbb{Q}$. How many are there? Are the ones you've given involving denominators divisible only by certain primes distinct? So that gives you the

cardinality. Are these all of them?

13.1.35 Question 1.35

Is the additive group of the reals isomorphic to the multiplicative group of the positive reals? Is the same result true with reals replaced by rationals?

13.1.36 Question 1.36

What groups have nontrivial automorphisms?

13.1.37 Question 1.37

A subgroup H of a group G that meets every conjugacy class is in fact G. Why is that true?

13.1.38 Question 1.38

Let G be the group of invertible 3×3 matrices over $\mathbb{F}_p$, for p prime. What does basic group theory tell us about G?

How many conjugates does a Sylow p-subgroup have? Give a matrix form for the elements in this subgroup.

Explain the conjugacy in terms of eigenvalues and eigenvectors. give a matrix form for the normaliser of the Sylow p-subgroup.

13.1.39 Question 1.39

Let's look at $\mathrm{SL}_2(\mathbb{F}_3)$. How many elements are in that group? What is its centre? Identify $\mathrm{PSL}_2(\mathbb{F}_3)$ as a permutation group.

13.1.40 Question 1.40

How many elements does $\mathfrak{gl}_2(\mathbb{F}_q)$ have? How would you construct representations?

What can you say about the 1-dimensional representations? What can you say about simplicity of some related groups?

13.1.41 Question 1.41.

A subgroup of a finitely-generated free abelian group is?

A subgroup of a finitely-generated free group is..? Prove your answers.

13.1.42 Question 1.42

What are the subgroups of $\mathbb{Z}^2$?

13.1.43 Question 1.43

What are the subgroups of the free group F_2? How many generators can you have?

Can you find one with 3 generators? 4 generators? Countably many generators?

Is the subgroup with 4 generators you found normal? Why? Can you find a normal one?

13.1.44 Question 1.44

Talk about the possible subgroups of $\mathbb{Z}^3$. Now suppose that you have a subgroup of $\mathbb{Z}^3$. What theorem tells you something about the structure of the quotient group?

13.2 Classification of Finite groups

13.2.1 Question 2.1

Given a finite abelian group with at most n elements of order divisible by n, prove it's cyclic.

13.2.2 Question 2.2

Suppose I asked you to classify groups of order 4. Why isn't there anything else? Which of those could be realised as a Galois group over $\mathbb{Q}$?

13.2.3 Question 2.3

State/prove the Sylow theorems.

13.2.4 Question 2.4

Classify groups of order 35.

13.2.5 Question 2.5

Classify groups of order 21.

13.2.6 Question 2.6

Discuss groups of order 55.

13.2.7 Question 2.7

Classify groups of order 14. Why is there a group of order 7? Are all index-2 subgroups normal?

13.2.8 Question 2.8

How many groups are there of order 15? Prove it.

13.2.9 Question 2.9

Classify all groups of order 8.

13.2.10 Question 2.10

Classify all groups of order p^3 for p prime.

13.2.11 Question 2.11

What are the groups of order p^2? What about pq? What if q is congruent to 1 (mod p)?

13.2.12 Question 2.12

What are the groups of order 12? Can there be a group of order 12 with 2 nonisomorphic subgroups of the same order?

13.2.13 Question 2.13

How would you start finding the groups of order 56? Is there in fact a way for $\mathbb{Z}/7\mathbb{Z}$ to act on a group of order 8 nontrivially?

13.2.14 Question 2.14

How many abelian groups are there of order 36?

13.2.15 Question 2.15

What are the abelian groups of order 16?

13.2.16 Question 2.16.

What are the abelian groups of order 9? Prove that they are not isomorphic. groups of order 27?

13.2.17 Question 2.17

How many abelian groups of order 200 are there?

13.2.18 Question 2.18

Prove there is no simple group of order 132.

13.2.19 Question 2.19

Prove that there is no simple group of order 160. What can you say about the structure of groups of that order?

13.2.20 Question 2.20

Prove that there is no simple group of order 40.

13.3 Fields and Galois Theory

13.3.1 Question 3.1

What is the Galois group of a finite field? What is a generator? How many elements does a finite field have? What can you say about the multiplicative group? Prove it.

13.3.2 Question 3.2

Classify finite fields, their subfields, and their field extensions. What are the automorphisms of a finite field?

13.3.3 Question 3.3

Take a finite field extension $\mathbb{F}_p^n$ over $\mathbb{F}_p$. What is Frobenius? What is its characteristic polynomial?

13.3.4 Question 3.4

What are the characteristic and minimal polynomial of the Frobenius automorphism?

13.3.5 Question 3.5

What's the field with 25 elements?

13.3.6 Question 3.6

What is the multiplicative group of $\mathbb{F}_9$?

13.3.7 Question 3.7

What is a separable extension? Can $\mathbb{Q}$ have a non-separable extension? How about $\mathbb{Z}/p\mathbb{Z}$? Why not? Are all extensions of characteristic 0 fields separable? Of finite fields? Prove it.

Give an example of a field extension that's not separable.

13.3.8 Question 3.8

Are there separable polynomials of any degree over any field?

13.3.9 Question 3.9

What is a perfect field and why is this important? Give an example of a non-perfect field.

13.3.10 Question 3.10

What is Galois theory? State the main theorem. What is the splitting field of $x^5 - 2$ over $\mathbb{Q}$? What are the intermediate extensions? Which extensions are normal, which are not, and why? What are the Galois groups (over Q) of all intermediate extensions?

13.3.11 Question 3.11

What is a Galois extension?

13.3.12 Question 3.12

Take a quadratic extension of a field of characteristic 0. Is it Galois? Take a degree 2 extension on top of that. Does it have to be Galois over the base field? What statement in group theory can you think of that reflects this?

13.3.13 Question 3.13.

Is Abelian Galois extension transitive? That is, if K has abelian Galois group over E, E has abelian Galois group over F , and K is a Galois extension of F, is it necessarily true that $\mathrm{Gal}(K/F)$ is also abelian? Give a counterexample involving number fields as well as one involving function fields.

13.3.14 Question 3.14

What is a Kummer extension?

13.3.15 Question 3.15

Say you have a field extension with only finitely many intermediate fields. Show that it is a simple extension.

13.3.16 Question 3.16

Tell me a condition on the Galois group which is implied by irreducibility of the polynomial. What happens when the polynomial has a root in the base field?

13.3.17 Question 3.17

What is the discriminant of a polynomial?

13.3.18 Question 3.18

If we think of the Galois group of a polynomial as contained in S_n, when is it contained in A_n?

13.3.19 Question 3.19

Is $\mathbb{Q}(\sqrt[3]{21})$ normal? What is its splitting field? What is its Galois group? Draw the lattice of subfields.

13.3.20 Question 3.20

What's the Galois group of $x^2 + 1$ over Q? What's the integral closure of $\mathbb{Z}$ in $\mathbb{Q}(i)$?

13.3.21 Question 3.21

What's the Galois group of $x^2 + 9$?

13.3.22 Question 3.22

What is the Galois group of $x^2 - 2$? Why is $x^2 - 2$ irreducible?

13.3.23 Question 3.23

What is the Galois group of

$$\mathbb{Q}(\sqrt{2}, \sqrt{3}) \,/\, \mathbb{Q}?$$

13.3.24 Question 3.24

What is the Galois group of

$$\mathbb{Q}\left(\sqrt{n_1}, \cdots, \sqrt{n_m}\right) \,/\, \mathbb{Q}(\sqrt{n_1} + \cdots + \sqrt{n_m})?$$

13.3.25 Question 3.25

What are the Galois groups of irreducible cubics?

13.3.26 Question 3.26

If an irreducible cubic polynomial has Galois group NOT contained in A3, does it necessarily have to be all of S_3?

13.3.27 Question 3.27

Compute the Galois group of $x^3 - 2$ over the rationals.

13.3.28 Question 3.28

How would you find the Galois group of $x^3 + 2x + 1$? Adjoin a root to $\mathbb{Q}$. Can you say something about the roots of $x^3 + 3x + 1$ in this extension?

13.3.29 Question 3.29

Compute the Galois group of $x^3 + 6x + 3$.

13.3.30 Question 3.30

Find the Galois group of $x^4 - 2$ over Q.

13.3.31 Question 3.31

What's the Galois group of $x^4 - 3$?

13.3.32 Question 3.32

What is the Galois group of $x^4 - 2x^2 + 9$?

13.3.33 Question 3.33

Calculate the Galois group of $x^5 - 2$.

13.3.34 Question 3.34.

Discuss sufficient conditions on a polynomial of degree 5 to have Galois group S_5 over $\mathbb{Q}$ and prove your statements.

13.3.35 Question 3.35

Show that if f is an irreducible quintic with precisely two non-real roots, then its Galois group is S_5.

13.3.36 Question 3.36

Suppose you have a degree 5 polynomial over a field. What are necessary and sufficient conditions for its Galois group to be of order divisible by 3? Can you give an example of an irreducible polynomial in which this is not the case?

13.3.37 Question 3.37

What is the Galois group of $x^7 - 1$ over the rationals?

13.3.38 Question 3.38

What is the Galois group of the polynomial $x^n - 1$ over $\mathbb{Q}$?

13.3.39 Question 3.39

Describe the Galois theory of cyclotomic extensions.

13.3.40 Question 3.40

What is the maximal real field in a cyclotomic extension $\mathbb{Q}(\zeta_n)/\mathbb{Q}$?

13.3.41 Question 3.41

Compute the Galois group of $p(x) = x^7 - 3$.

13.3.42 Question 3.42

What Galois stuff can you say about $x^{2n} - 2$?

13.3.43 Question 3.43

What are the cyclic extensions of (prime) order p?

13.3.44 Question 3.44

Can you give me a polynomial whose Galois group is $\mathbb{Z}/3\mathbb{Z}$?

13.3.45 Question 3.45

Which groups of order 4 can be realised as a Galois group over $\mathbb{Q}$?

13.3.46 Question 3.46

Give a polynomial with S_3 as its Galois group.

13.3.47 Question 3.47

Give an example of a cubic with Galois group S_3.

13.3.48 Question 3.48

How do you construct a polynomial over $\mathbb{Q}$ whose Galois group is S_n? Do it for $n = 7$ in particular.

13.3.49 Question 3.49

What's a Galois group that's not S_n or A_n?

13.3.50 Question 3.50

Which finite groups are Galois groups for some field extension?

13.3.51 Question 3.51

What Galois group would you expect a cubic to have?

13.3.52 Question 3.52

Draw the subgroup lattice for S_3.

13.3.53 Question 3.53

Do you know what the quaternion group is? How many elements are there of each order? Suppose I have a field extension of the rationals with Galois group the quaternion group. How many quadratic extensions does it contain? Can any of them be imaginary?

13.3.54 Question 3.54

Suppose you are given a finite Galois extension $K/\mathbb{Q}$ by $f(x) \in \mathbb{Z}[x]$ such that $\deg(f) = n$ and $\mathrm{Gal}(K/Q) = S_n$. What can you say about the roots?

13.3.55 Question 3.55

How many automorphisms does the complex field have? How can you extend a simple automorphism $\sqrt{2} \mapsto -\sqrt{2}$ of an algebraic field into $\mathbb{C}$? How can you extend a subfield automorphism? What feature of $\mathbb{C}$ allows you to?

13.3.56 Question 3.56.

Can it happen that a proper subfield of C is isomorphic to C? How?

13.3.57 Question 3.57

Consider the minimal polynomial $f(x)$ for a primitive mth root of unity. Prove that if p divides $f(a)$ for some integer a and $\gcd(p, m) = 1$ then m divides $p - 1$. Use this fact to show that there are infinitely many primes congruent to 1 (mod m).

13.3.58 Question 3.58

What is Dirichlet's theorem about primes in arithmetic progression? What can you say about the density of such primes?

13.3.59 Question 3.59

How many irreducible polynomials of degree six are there over $\mathbb{F}_2$?

13.3.60 Question 3.60

Can you have a degree 7 irreducible polynomial over $\mathbb{F}_p$? How about a degree 14 irreducible polynomial?

13.3.61 Question 3.61

How many irreducible polynomials are there of degree 4 over $\mathbb{F}_2$?

13.3.62 Question 3.62

For each prime p, give a polynomial of degree p that is irreducible over $\mathbb{F}_p$. You can do it in a "uniform" way.

13.3.63 Question 3.63

Can we solve general quadratic equations by radicals? And what about cubics and so on? Why can't you solve 5th degree equations by radicals?

13.3.64 Question 3.64

Talk about solvability by radicals. Why is S_5 not solvable? Why is A_5 simple?

13.3.65 Question 3.65

For which n can a regular n-gon be constructed by ruler and compass?

13.3.66 Question 3.66

How do you use Galois theory (or just field theory) to prove the impossibility of trisecting an angle? Doubling a cube? Squaring a circle?

13.3.67 Question 3.67

Which numbers are constructible? Give an example of a non-constructible number whose degree is nevertheless a power of 2.

13.3.68 Question 3.68

State and prove Eisenstein's Criterion.

13.3.69 Question 3.69

Why is $(x^p - 1)/(x - 1)$ irreducible over $\mathbb{Q}$?

13.3.70 Question 3.70

Can you prove the fundamental theorem of algebra using Galois theory? What do you need from analysis to do so?

13.3.71 Question 3.71

What are the symmetric polynomials?

13.3.72 Question 3.72

State the fundamental theorem of symmetric polynomials.

13.3.73 Question 3.73

Is the discriminant of a polynomial always a polynomial in the coefficients? What does this have to do with symmetric polynomials?

13.3.74 Question 3.74

Find a non-symmetric polynomial whose square is symmetric.

13.3.75 Question 3.75

Let f be a degree 4 polynomial with integer coefficients. What's the smallest finite field in which f necessarily has four roots?

13.3.76 Question 3.76

Define p-adic numbers. What is a valuation?

13.3.77 Question 3.77

What's Hilbert's theorem 90?

13.3.78 Question 3.78

Consider a nonconstant function between two compact Riemann Surfaces. How is it related to Galois theory?

13.4 Normal Forms

13.4.1 Question 4.1

What is the connection between the structure theorem for modules over a PID and conjugacy classes in the general linear group over a field?

13.4.2 Question 4.2

Explain how the structure theorem for finitely-generated modules over a PID applies to a linear operator on a finite dimensional vector space.

13.4.3 Question 4.3

I give you two matrices over a field. How would you tell if they are conjugate or not? What theorem are you using? State it. How does it apply to this situation? Why is $k[x]$ a PID? If two matrices are conjugate over the algebraic closure of a field, does that mean that they are conjugate over the base field too?

13.4.4 Question 4.4

If two real matrices are conjugate in $\mathrm{Mat}(n \times n, \mathbb{C})$, are they necessarily conjugate in $\mathrm{Mat}(n \times N, R)$ as well?

13.4.5 Question 4.5

Give the 4×4 Jordan forms with minimal polynomial $(x-1)(x-2)^2$.

13.4.6 Question 4.6

Talk about Jordan canonical form. What happens when the field is not algebraically closed?

13.4.7 Question 4.7

What are all the matrices that commute with a given Jordan block?

13.4.8 Question 4.8

How do you determine the number and sizes of the blocks for Jordan canonical form?

13.4.9 Question 4.9

For any matrix A over the complex numbers, can you solve $B^2 = A$?

13.4.10 Question 4.10

What is rational canonical form?

13.4.11 Question 4.11

Describe all the conjugacy classes of 3×3 matrices with rational entries which satisfy the equation $A^4 - A^3 - A + 1 = 0$. Give a representative in each class.

13.4.12 Question 4.12

What 3×3 matrices over the rationals (up to similarity) satisfy $f(A) = 0$, where $f(x) = (x^2 + 2)(x-1)^3$? List all possible rational forms.

13.4.13 Question 4.13

What can you say about matrices that satisfy a given polynomial (over an algebraically closed field)? How many of them are there? What about over a finite field? How many such matrices are there then?

13.4.14 Question 4.14

What is a nilpotent matrix?

13.4.15 Question 4.15

When do the powers of a matrix tend to zero?

13.4.16 Question 4.16

If the traces of all powers of a matrix A are 0, what can you say about A?

13.4.17 Question 4.17

When and how can we solve the matrix equation $\exp(A) = B$? Do it over the complex numbers and over the real numbers. give a counterexample with real entries.

13.4.18 Question 4.18

Say we can find a matrix A such that $\exp(A) = B$ for B in $SL_n(\mathbb{R})$. Does A also have to be in $\mathrm{SL}_n(R)$? Does A *need* to be in $SL_n(R)$?

13.4.19 Question 4.19

Is a square matrix always similar to its transpose?

13.4.20 Question 4.20

What are the conjugacy classes of $\mathrm{SL}_2(\mathbb{R})$?

13.4.21 Question 4.21

What are the conjugacy classes in $\mathrm{GL}_2(\mathbb{C})$?

13.5 Matrices and Linear Algebra

13.5.1 Question 5.1

What is a bilinear form on a vector space? When are two forms equivalent? What is an orthogonal matrix? What's special about them?

13.5.2 Question 5.2

What are the possible images of the unit circle under a linear transformation of $\mathbb{R}^2$?

13.5.3 Question 5.3

Explain geometrically how you diagonalise a quadratic form.

13.5.4 Question 5.4

Do you know Witt's theorem on real quadratic forms?

13.5.5 Question 5.5

Classify real division algebras.

13.5.6 Question 5.6

Consider the simple operator on C given by multiplication by a complex number. It decomposes into a stretch and a rotation. What is the generalisation of this to operators on a Hilbert space?

13.5.7 Question 5.7

Do you know about singular value decomposition?

13.5.8 Question 5.8

What are the eigenvalues of a symmetric matrix?

13.5.9 Question 5.9

What can you say about the eigenvalues of a skew-symmetric matrix?

13.5.10 Question 5.10

Prove that the eigenvalues of a Hermitian matrix are real and those of a unitary matrix are unitary.

13.5.11 Question 5.11

Prove that symmetric matrices have real eigenvalues and can be diagonalised by orthogonal matrices.

13.5.12 Question 5.12

To which operators does the spectral theorem for symmetric matrices generalise?

13.5.13 Question 5.13

Given a skew-symmetric/skew-Hermitian matrix S, show that $U = (S + I)(S - I) - 1$ is orthogonal/unitary. Then find an expression for S in terms of U.

13.5.14 Question 5.14

If a linear transformation preserves a nondegenerate alternating form and has k as an eigenvalue, prove that $1/k$ is also an eigenvalue.

13.5.15 Question 5.15

State/prove the Cayley–Hamilton theorem.

13.5.16 Question 5.16

Are diagonalisable $N \times N$ matrices over the complex numbers dense in the space of all $N \times N$ matrices over the complex numbers? How about over another algebraically closed field if we use the Zariski topology?

13.5.17 Question 5.17

For a linear ODE with constant coefficients, how would you solve it using linear algebra?

13.5.18 Question 5.18

What can you say about the eigenspaces of two matrices that commute with each other?

13.5.19 Question 5.19

What is a Toeplitz operator?

13.5.20 Question 5.20

What is the number of invertible matrices over $\mathbb{Z}/p\mathbb{Z}$?

13.6 Rings

13.6.1 Question 6.1

State the Chinese remainder theorem in any form you like. Prove it.

13.6.2 Question 6.2

What is a PID? What's an example of a UFD that is not a PID? Why? Is $k[x]$ a PID? Why?

13.6.3 Question 6.3

Is $\mathbb{C}[x, y]$ a PID? Is $\langle x, y \rangle$ a prime ideals in it?

13.6.4 Question 6.4

Do polynomials in several variables form a PID?

13.6.5 Question 6.5

Prove that the integers form a PID.

13.6.6 Question 6.6

Give an example of a PID with a unique prime ideal.

13.6.7 Question 6.7

What is the relation between Euclidean domains and PIDs?

13.6.8 Question 6.8

Do you know a PID that's not Euclidean?

13.6.9 Question 6.9

Give an example of a UFD which is not a Euclidean domain.

13.6.10 Question 6.10

Is a ring of formal power series a UFD?

13.6.11 Question 6.11

Is a polynomial ring over a UFD again a UFD?

13.6.12 Question 6.12

What does factorisation over $\mathbb{Q}[x]$ say about factorisation over $\mathbb{Z}[x]$?

13.6.13 Question 6.13

Give an example of a ring where unique factorisation fails.

13.6.14 Question 6.14

Factor 6 in two different ways in $\mathbb{Z}[\sqrt{-5}]$ Is there any way to explain the two factorisations? Factor the ideal generated by 6 into prime ideals.

13.6.15 Question 6.15

What's the integral closure of $\mathbb{Z}$ in $\mathbb{Q}(i)$?

13.6.16 Question 6.16

Find all primes in the ring of Gaussian integers.

13.6.17 Question 6.17

What is a ring of integers? What does "integral over $\mathbb{Z}$" mean?

13.6.18 Question 6.18

Let $\mathcal{O}$ be the ring of integers of $\mathbb{Q}(d)$, where $d > 0$. What can you say about the quotient of O by one of its prime ideals?

13.6.19 Question 6.19

Do you know about Dedekind domains and class numbers?

13.6.20 Question 6.20

Talk about factorisation and primes in a polynomial ring. What is irreducibility? For what rings R is it true that $R[x_1, \cdots, x_n]$ is a unique factorisation domain? What is wrong with unique factorisation if we don't have a domain? Now, PIDs are Noetherian, but are there UFDs which are not?

13.6.21 Question 6.21

What is the radical of an ideal? What is special about elements in the nilradical?

13.6.22 Question 6.22

Define the "radical" of an ideal. Prove it is an ideal. Prove that the ideal of all polynomials vanishing on the zero set of I is $\sqrt{I}$.

13.6.23 Question 6.23.

Do you know what the radical is? Use the fact that the intersection of all prime ideals is the set of all nilpotent elements to prove that $F[x]$ has an infinite number of prime ideals, where F is a field.

13.6.24 Question 6.24

What are the radical ideals in $\mathbb{Z}$?

13.6.25 Question 6.25

Give a prime ideal in $\urcorner[x, y]$. Why is it prime? What is the variety it defines? What is the Nullstellensatz? Can you make some maximal ideals?

13.6.26 Question 6.26

State/describe Hilbert's Nullstellensatz. Sketch a proof.

13.6.27 Question 6.27

What is an irreducible variety? Give an example of a non-irreducible one.

13.6.28 Question 6.28

What are the prime ideals and maximal ideals of $\mathbb{Z}[x]$?

13.6.29 Question 6.29

Is the following map an isomorphism?

$$\begin{aligned} \mathbb{Z}[t]/\langle t^p - 1\rangle &\to \mathbb{Z}[w] \\ t &\mapsto w \text{ where } w^p = 1. \end{aligned}$$

13.6.30 Question 6.30

Describe the left, right, and two-sided ideals in the ring of square matrices of a fixed size. Now identify the matrix algebra $\mathrm{Mat}(n \times n, K)$ with $\underset{K}{\mathrm{End}}(V)$ where V is an n-dimensional K-vector space. Try to geometrically describe the simple left ideals and also the simple right ideals via that identification.

13.6.31 Question 6.31

Give examples of maximal ideals in $K = R \times R \times R \times \cdots$, the product of countably many copies of R. What about for a product of countably many copies of an arbitrary commutative ring R?

13.6.32 Question 6.32

Consider a commutative ring, R, and a maximal ideal I, what can you say about the structure of R/I? What if I were prime?

13.6.33 Question 6.33

Define "Noetherian ring". give an example.

13.6.34 Question 6.34

Prove the Hilbert basis theorem.

13.6.35 Question 6.35

What is a Noetherian ring? If I is an ideal in a Noetherian ring with a unit, what is the intersection of I^n over all positive integers n?

13.6.36 Question 6.36

What is the Jacobson radical? If R is a finitely-generated algebra over a field what can you say about it?

13.6.37 Question 6.37

Give an example of an Artinian ring.

13.6.38 Question 6.38

State the structure theorem for semisimple Artinian rings.

13.6.39 Question 6.39

What is a semisimple algebra? State the structure theorem for semisimple algebras.

13.6.40 Question 6.40

What is a matrix algebra?

13.6.41 Question 6.41

Does L_1 have a natural multiplication with which it becomes an algebra?

13.6.42 Question 6.42.

Consider a translation-invariant subspace of L_1. What can you say about its relation to L_2 as a convolution algebra?

13.6.43 Question 6.43

State the structure theorem for simple rings.

13.6.44 Question 6.44

Do you know an example of a local ring? Another one? What about completions?

13.6.45 Question 6.45

Consider the space of functions from the natural numbers to $\mathbb{C}$ endowed with the usual law of addition and the following analogue of the convolution product:

$$(f * g)(n) = \sum_{d|n} f(d) g\left(\frac{n}{d}\right).$$

Show that this is a ring. What does this ring remind you of and what can you say about it?

13.6.46 Question 6.46

Prove that any finite division ring is a field (that is, prove commutativity). Give an example of a (necessarily infinite) division ring which is NOT a field.

13.6.47 Question 6.47

Prove that all finite integral domains are fields.

13.6.48 Question 6.48

Can a polynomial over a division ring have more roots than its degree?

13.6.49 Question 6.49

Classify (finite-dimensional) division algebras over $\mathbb{R}$.

13.6.50 Question 6.50

Give an example of a $\mathbb{C}$-algebra which is not semisimple.

13.6.51 Question 6.51

What is Wedderburn's theorem? What does the group ring generated by $\mathbb{Z}/5\mathbb{Z}$ over $\mathbb{Q}$ look like?

What if we take the noncyclic group of order 4 instead of $\mathbb{Z}/5\mathbb{Z}$? The quaternion group instead of $\mathbb{Z}/5\mathbb{Z}$?

13.6.52 Question 6.52

Tell me about group rings. What do you know about them?

13.7 Modules

13.7.1 Question 7.1

How does one prove the structure theorem for modules over PID? What is the module and what is the PID in the case of abelian groups?

13.7.2 Question 7.2

If M is free abelian, how can I put quotients of M in some standard form? What was crucial about the integers here (abelian groups being modules over $\mathbb{Z}$)? How does the procedure simplify if the ring is a Euclidean domain, not just a PID?

13.7.3 Question 7.3

Suppose D is an integral domain and the fundamental theorem holds for finitely-generated modules over D (i.e. they are all direct sums of finitely many cyclic modules).

Does D have to be a PID?

13.7.4 Question 7.4

Classify finitely-generated modules over $\mathbb{Z}$, over PIDs, and over Dedekind rings.

13.7.5 Question 7.5

Prove a finitely-generated torsion-free abelian group is free abelian.

13.7.6 Question 7.6.

What is a tensor product? What is the universal property? What do the tensors look like in the case of vector spaces?

13.7.7 Question 7.7

Now we'll take the tensor product of two abelian groups, that is, $\mathbb{Z}$-modules. Take $\mathbb{Z}/p\mathbb{Z}$ and $\mathbb{Z}/q\mathbb{Z}$, where p and q are distinct primes. What is their tensor product?

13.7.8 Question 7.8

What is a projective module?

13.7.9 Question 7.9

What is an injective module?

13.7.10 Question 7.10

Do you know an example of a flat module?

13.8 Representation Theory

13.8.1 Question 8.1

Define "representation" of a group. Define "irreducible representation". Why can you decompose representations of finite groups into irreducible ones? Construct an in- variant inner product.

13.8.2 Question 8.2

State and prove Maschke's theorem. What can go wrong if you work over the real field? What can go wrong in characteristic p?

13.8.3 Question 8.3

Do you know what a group representation is? Do you know what the trace of a group representation is?

13.8.4 Question 8.4

State/prove/explain Schur's lemma.

13.8.5 Question 8.5

What can you say about characters? What are the orthogonality relations? How do you use characters to determine if a given irreducible representation is a subspace of another given representation?

13.8.6 Question 8.6

What's the relation between the number of conjugacy classes in a finite group and the number of irreducible representations?

13.8.7 Question 8.7

What is the character table? What field do its entries lie in?

13.8.8 Question 8.8

Why is the character table a square?

13.8.9 Question 8.9

If $\chi(g)$ is real for every character χ, what can you say about g?

13.8.10 Question 8.10

What's the regular representation?

13.8.11 Question 8.11

Give two definitions of "induced representation". Why are they equivalent?

13.8.12 Question 8.12

If you have a representation of H, a subgroup of a group G, how can you induce a representation of G?

13.8.13 Question 8.13

If you have an irreducible representation of a subgroup, is the induced representation of the whole group still irreducible?

13.8.14 Question 8.14.

What can you say about the kernel of an irreducible representation? How about kernels of direct sums of irreducibles? What kind of functor is induction? Left or right exact?

13.8.15 Question 8.15

What is Frobenius reciprocity?

13.8.16 Question 8.16

Given a normal subgroup H of a finite group G, we lift all the representations of G/H to representations of G.

Show that the intersection of the kernels of all these representations is precisely H. What can you say when H is the commutator subgroup of G?

13.8.17 Question 8.17

If you have two linear representations π_1 and π_2 of a finite group G such that $\pi_1(g)$ is conjugate to $\pi_2(g)$ for every g in G, is it true that the two representations are isomorphic?

13.8.18 Question 8.18

Group representations: What's special about using $\mathbb{C}$ in the definition of group algebra?

Is it possible to work over other fields?

What goes wrong if the characteristic of the field divides the order of the group?

13.8.19 Question 8.19

Suppose you have a finite p-group, and you have a representation of this group on a finite-dimensional vector space over a finite field of characteristic p. What can you say about it?

13.8.20 Question 8.20

Let (π, V) be a faithful finite-dimensional representation of G. Show that, given any irreducible representation of G, the nth tensor power of $\mathrm{GL}(V)$ will contain it for some large enough n.

13.8.21 Question 8.21

What are the irreducible representations of finite abelian groups?

13.8.22 Question 8.22

What are the group characters of the multiplicative group of a finite field?

13.8.23 Question 8.23

Are there two nonisomorphic groups with the same representations?

13.8.24 Question 8.24

If you have a $\mathbb{Z}/5\mathbb{Z}$ action on a complex vector space, what does this action look like? What about an S_3 action? A dihedral group of any order?

13.8.25 Question 8.25

What are the representations of S_3? How do they restrict to S_2?

13.8.26 Question 8.26

Tell me about the representations of D_4. Write down the character table. What is the 2-dimensional representation? How can it be interpreted geometrically?

13.8.27 Question 8.27

How would you work out the orders of the irreducible representations of the dihedral group D_n?

Why is the sum of squares of dimensions equal to the order of the group?

13.8.28 Question 8.28

Do you know any representation theory? What about representations of A_4?

Give a nontrivial one. What else is there? How many irreducible representations do we have? What are their degrees? Write the character table of A_4.

13.8.29 Question 8.29

Write the character table for S_4.

13.8.30 Question 8.30

Start constructing the character table for S_5.

13.8.31 Question 8.31.

How many irreducible representations does S_n have?

What classical function in mathematics does this number relate to?

13.8.32 Question 8.32

Discuss representations of $\mathbb{Z}$, the infinite cyclic group. What is the group algebra of $\mathbb{Z}$?

13.8.33 Question 8.33

What is a Lie group? Define a unitary representation. What is the Peter–Weyl theorem? What is the Lie algebra? The Jacobi identity? What is the adjoint representation of a Lie algebra? What is the commutator of two vector fields on a manifold?

When is a representation of $\mathbb{Z}$ completely reducible? Why?

Which are the indecomposable modules?

13.8.34 Question 8.34

Talk about the representation theory of compact Lie groups. How do you know you have a finite-dimensional representation?

13.8.35 Question 8.35

How do you prove that any finite-dimensional representation of a compact Lie group is equivalent to a unitary one?

13.8.36 Question 8.36

Do you know a Lie group that has no faithful finite-dimensional representations?

13.8.37 Question 8.37

What do you know about representations of SO(2)? SO(3)?

13.9 Categories and Functors

13.9.1 Question 9.1

Which is the connection between Hom and tensor product? What is this called in representation theory?

13.9.2 Question 9.2

Can you get a long exact sequence from a short exact sequence of abelian groups together with another abelian group?

13.9.3 Question 9.3

Do you know what the Ext functor of an abelian group is? Do you know where it appears? What is $\mathrm{Ext}(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z}/n\mathbb{Z})$? What is $\mathrm{Ext}(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z})$?

14 | Appendix: Extra Topics

14.1 Characteristic Subgroups

Slogan 14.1.1
Normality is not transitive!

I.e. if $H \trianglelefteq G$ and $N \trianglelefteq H$, it's not necessarily the case that $N \trianglelefteq G$.

Definition 14.1.2 (Characteristic Subgroups)
A subgroup $H \leq G$ is **characteristic** in G, written $H \operatorname{ch} G$, iff for every $\varphi \in \operatorname{Aut}(G)$, $\varphi(H) \leq H$. Equivalently, $\varphi(H) = H$. I.e. H is fixed (not necessarily pointwise) under every automorphism of the ambient group G.

Remark 14.1.3 ***(Characteristic isn't equivalent to normalcy):*** Characteristic subgroups are normal, because $\psi_g(-) \coloneqq g(-)g^{-1}$ is an (inner) automorphic of G. Not every normal subgroup is characteristic: take $G \coloneqq H_1 \times H_2$ and $\psi(x, y) = (y, x)$.

Proposition 14.1.4 ***(Fixing transitivity of normality).***
Characteristic subgroups of normal subgroups are normal, i.e. if $H \trianglelefteq G$ and $N \operatorname{ch} H$, then $N \trianglelefteq G$.

Proof (?).
$A \operatorname{ch} B \trianglelefteq C \implies A \trianglelefteq C$:

- $A \operatorname{ch} B$ iff A is fixed by every $\psi \in \operatorname{Aut}(B)$., WTS $cAc^{-1} = A$ for all $c \in C$.
- Since $B \trianglelefteq C$, the automorphism $\psi(-) \coloneqq c(-)c^{-1}$ descends to an element of $\operatorname{Aut}(B)$.
- Then $\psi(A) = A$ since $A \operatorname{ch} B$, so $cAc^{-1} = A$ and $A \trianglelefteq C$.

■

Proposition 14.1.5 ***(Centers are characteristic).***
For any group G,

$$Z(G) \operatorname{ch} G.$$

Proof (?).

Let $\psi \in \mathrm{Aut}(H)$ and $x = \psi(y) \in \psi(Z(H))$ so $y \in Z(H)$, then for arbitrary $h \in H$,

$$\begin{aligned} \psi(y)h &= \psi(y)(\psi \circ \psi^{-1})(h) \\ &= \psi(y \cdot \psi^{-1}(h)) \\ &= \psi(\psi^{-1}(h) \cdot y) && \text{since } \psi^{-1}(h) \in H,\ y \in Z(H) \\ &= h\psi(y). \end{aligned}$$

■

14.2 Normal Closures and Cores

Definition 14.2.1 (Normal Closure of a Subgroup)
The smallest normal subgroup of G containing H:

$$H^G \coloneqq \{gHg^{-1} : g \in G\} = \bigcap \{N : H \leq N \trianglelefteq G\}.$$

Definition 14.2.2 (Normal Core of a subgroup)
The largest normal subgroup of G containing H:

$$H_G = \bigcap_{g \in G} gHg^{-1} = \langle N : N \trianglelefteq G \ \&\ N \leq H \rangle = \ker \psi.$$

where

$$\begin{aligned} \psi : G &\to \mathrm{Aut}(G/H) \\ g &\mapsto (xH \mapsto gxH) \end{aligned}$$

Theorem 14.2.3 ***(Fratini's Argument).***
If $H \trianglelefteq G$ and $P \in \mathrm{Syl}_p(G)$, then $HN_G(P) = G$ and $[G : H]$ divides $|N_G(P)|$.

14.2.1 Exercises

Exercise 14.2.4 (?)
Show that $Z(G) \leq G$ is always characteristic.

Solution:
Let $\psi \in \mathrm{Aut}(G)$. For one containment, we can show $\psi(g) = h = h\psi(g)$ for all $\psi(g) \in \psi(G)$

and $h \in G$. This is a computation:

$$\begin{aligned}\psi(g)h &= \psi(g)(\psi\psi^{-1})(h)\\ &= \psi(g)\psi(\psi^{-1}(h))\\ &= \psi(\psi^{-1}(h)g)\\ &= (\psi\psi^{-1})(h)\psi(g)\\ &= h\psi(g).\end{aligned}$$

This yields $\psi(Z(G)) \subseteq Z(G)$. Applying the same argument to ψ^{-1} yields $\psi^{-1}(Z(G)) \subseteq Z(G)$. Since ψ is a bijection, $\psi\psi^{-1}(A) = A$ for all $A \leq G$, so $Z(G) \subseteq \psi(Z(G))$.

14.3 Nilpotent Groups

Definition 14.3.1 (Nilpotent)
A group G is **nilpotent** iff G has a terminating upper central series.

Moral: the adjoint map is nilpotent.

Theorem 14.3.2 ***(Characterization of Nilpotent Groups).***
G is nilpotent iff G has an upper central series terminating at G.

Theorem 14.3.3 ***(Characterization of Nilpotent Groups).***
G is nilpotent iff G has a lower central series terminating at 1.

Theorem 14.3.4 ***(Nilpotents Have All Sylows Normal).***
A group G is nilpotent iff all of its Sylow p-subgroups are normal for every p dividing $|G|$.

Theorem 14.3.5 ***(Nilpotent Implies Maximal Normals).***
A group G is nilpotent iff every maximal subgroup is normal.

Proposition 14.3.6.
For G a finite group, TFAE:

- G is nilpotent
- Normalizers grow, i.e. if $H < G$ is proper then $H < N_G(H)$.
- Every Sylow-p subgroup is normal
- G is the direct product of its Sylow p-subgroups
- Every maximal subgroup is normal
- G has a terminating *Lower* Central Series
- G has a terminating *Upper* Central Series

Fact 14.3.7

- Nilpotent groups satisfy the 2 out of 3 property.
- G has normal subgroups of order d for *every* d dividing $|G|$

Todo. Speci

15 | UGA Fall 2019 Problem Sets

15.1 Problem Set One

15.1.1 Exercises

Problem 15.1.1 (Hungerford 1.6.3)
If $\sigma = (i_1 i_2 \cdots i_r) \in S_n$ and $\tau \in S_n$, then show that $\tau\sigma\tau^{-1} = (\tau(i_1)\tau(i_2)\cdots\tau(i_r))$.

Problem 15.1.2 (Hungerford 1.6.4)
Show that $S_n \cong \langle (12), (123\cdots n)\rangle$ and also that $S_n \cong \langle (12), (23\cdots n)\rangle$

Problem 15.1.3 (Hungerford 2.2.1)
Let G be a finite abelian group that is not cyclic. Show that G contains a subgroup isomorphic to $\mathbb{Z}_p \oplus \mathbb{Z}_p$ for some prime p.

Problem 15.1.4 (Hungerford 2.2.12.b.)
Determine (up to isomorphism) all abelian groups of order 64; do the same for order 96.

Problem 15.1.5 (Hungerford 2.4.1)
Let G be a group and $A \trianglelefteq G$ be a normal abelian subgroup. Show that G/A acts on A by conjugation and construct a homomorphism $\varphi : G/A \to \mathrm{Aut}(A)$.

Problem 15.1.6 (Hungerford 2.4.9).)
Let $Z(G)$ be the center of G. Show that if $G/Z(G)$ is cyclic, then G is abelian.

Note that Hungerford uses the notation $C(G)$ for the center.

Problem 15.1.7 (Hungerford 2.5.6)
Let G be a finite group and$H \trianglelefteq G$ a normal subgroup of order p^k. Show that H is contained in every Sylow p-subgroup of G.

Problem 15.1.8 (Hungerford 2.5.9)
Let $|G| = p^n q$ for some primes $p > q$. Show that G contains a unique normal subgroup of index q.

15.1.2 Qual Problems

Problem 15.1.9
Let G be a finite group and p a prime number. Let X_p be the set of Sylow-p subgroups of G and n_p be the cardinality of X_p. Let $\mathrm{Sym}(X)$ be the permutation group on the set X_p.

1. Construct a homomorphism $\rho : G \to \mathrm{Sym}(X_p)$ with image a transitive subgroup (i.e. with a single orbit).
2. Deduce that if G is simple then the order of G divides $n_p!$.
3. Show that for any $1 \leq a \leq 4$ and any prime power p^k, no group of order ap^k is simple.

Solution:

1. Define the required group action by

$$\begin{aligned} \rho : G &\to \mathrm{Sym}(X_p) \\ g &\mapsto (\gamma_g : P \mapsto gPg^{-1}). \end{aligned}$$

   The claim is that this action is transitive on X_p. This can be equivalently stated as

$$\forall P \in X_p, \exists g \in G, P' \in X_p \;\middle|\; gP'g^{-1} = P.$$

   However, by Sylow 2, all Sylow $p-$subgroups are conjugate to each other, and thus this condition is satisfied.

2. Suppose that G is simple, so that we have

$$H \trianglelefteq G \implies H = \{e\} \text{ or } H = G.$$

   Note that $\mathrm{Sym}(X_p) = (n_p)!$, and if we have an injective homomorphism $G \xrightarrow{\varphi} \mathrm{Sym}(X_p)$, then $|G| = |\varphi(G)|$, since $\varphi(G) \leq \mathrm{Sym}(X_p)$ will be a subgroup and thus have order dividing $(n_p)!$, which proves the statement.

   Using the φ defined in (1), we can apply the first isomorphism theorem

$$G/\ker\varphi \cong \mathrm{im}\varphi \leq \mathrm{Sym}(X_p),$$

   and so it suffices to show that $\ker\varphi = \{e\}$.

   Note that since $\ker\varphi \trianglelefteq G$ and G is simple, we can only have $\ker\varphi = \{e\}$ or $\ker\varphi = G$.

   Towards a contradiction, suppose $\ker\varphi = G$.

   By definition, we have

$$\begin{aligned} \ker\varphi &= \{g \in G \;|\; \gamma_g = \mathrm{id}_{X_p}\} \\ &= \{g \in G \;|\; \forall P \in X_p,\ gPg^{-1} = P\} \\ &= \bigcap_{P \in X_p} N_G(P), \end{aligned}$$

and so the kernel of φ is the intersection of all of the normalizers of the Sylow $p-$subgroups.

But this means that $N_G(P) = G$ for every Sylow $p-$subgroup, which means that $n_p = 1$ and there is a unique P which must be normal in G. Since G is simple, this forces P to be trivial or the whole group.

Towards a contradiction, suppose $P = G$. Then G is a $p-$group and thus has order p^n. But then G has normal subgroups of order p^k for all $0 < k < n$, contradicting the simplicity of G.

But the only other option is that P is trivial, whereas we know nontrivial Sylow $p-$subgroups exist by Sylow 1.

Thus we can not have $\ker\varphi = G$, and so $\ker\varphi$ is trivial as desired.

3. Suppose $|G| = ap^k$, where $1 \leq a \leq 4$. Then by Sylow 3, we have $n_p = 1 \pmod p$ and n_p divides a. If $a = 1$, then $n_p = 1$, and so G can not be simple. Moreover, if $p \geq a$, then $n_p \leq a$ and $n_p = 1 \pmod p$ forces $n_p = 1$ again.

   So we can restrict our attention to $2 \leq a \leq 4$ and $p = 2, 3$, which reduces to checking the cases $ap^k = 2(3^k), 4(3^k)$, or $3(2^k)$ for $k \geq 1$.

   If $ap^k = 2(3^k)$, we have $n_3 = 1 \pmod 3$ and $n_3 \mid 2$, which forces $n_3 = 1$, so this can not be a simple group.

   Similarly, if $ap^k = 4(3^k)$, then $n_3 = 1 \pmod 3$ and n_3 divides 4, which forces $n_3 = 1$ and thus G can't be simple.

   If $ap^k = 3(2^k)$, then $n_2 = 1 \pmod 2$ and n_2 divides 3, so $n_2 = 1, 3$. But then $n_3! = 6$, and if $k > 1$, we have $3(2^k) > 6 = n_3!$, so G can not be simple by the result in (2).

   If $k = 1$, then G is order 6, so G is isomorphic to either $\mathbb{Z}_6$ or S_3. The group S_3 is not simple, since $A_3 \trianglelefteq S_3$, and the only simple cyclic groups are of prime order, so $\mathbb{Z}_6$ is not simple. This exhausts all of the possible cases.

Problem 15.1.10
Let G be a finite group and let $N \trianglelefteq G$, and let p be a prime number and Q a subgroup of G such that $N \subset Q$ and Q/N is a Sylow $p-$subgroup of G/N.

1. Prove that Q contains a Sylow $p-$subgroup of G.
2. Prove that every Sylow $p-$subgroup of G/N is the image of a Sylow $p-$subgroup of G.

Solution:
Proof.

1. Since Q/N is a Sylow $p-$subgroup of G/N, we can write $|G/N| = p^k l$ where $\gcd(p, l) = 1$, and $|Q/N| = p^k$.

   We can then write $|G| = p^n m$ where $n \geq l$ and $l \mid m$.

By the third isomorphism theorem, we have

$$\frac{G/N}{Q/N} \cong G/Q$$

and so

$$\left|\frac{G/N}{Q/N}\right| = \frac{|G/N|}{|Q/N|} = \frac{p^k l}{p^k} = l$$

and so $|G/Q| = l$ where $(p, l) = 1$, and thus

$$|G/Q| = |G|/|Q| = l \implies |G| = |Q|\ l.$$

We then have

$$p^n m = |Q|\ l,$$

and since $(p, l) = 1$, it must be the case that p^n divides $|Q|$. But since $Q \leq G$, this means that Q itself must be a Sylow $p-$ subgroup of G.

2. Let $P_N \in \text{Syl}(p, G/N)$. By the subgroup correspondence theorem, $P_n = H/N$ for some $H \leq G$ such that $N \subseteq H$.

   So choose $P_H \in \text{Syl}(p, H)$; the claim is that $P_H \in \text{Syl}(p, G)$ and that $\dfrac{P_H N}{N} \cong P_N$, which exhibits P_N as the image of a Sylow $p-$subgroup of G.

   We first have $P_H \in \text{Syl}(p, G)$, which follows because we have $[G/N, H/N] = [G : P_H]$ from the fourth isomorphism theorem, and thus $[G/N, P_N] = [G : P_H]$. In particular, since P_N is a Sylow $p-$subgroup, p does not divide $[G/N, P_N]$ and thus p doesn't divide $[G : P_H]$, which makes P_H a maximal $p-$subgroup in G and thus a Sylow $p-$subgroup.

   We then have $P_H N/N = P_N$, which follows because $P_H \leq H \implies P_H N/N \leq H/N = P_N \leq G/N$.

   However, it is also the case that $P_H N/N \in \text{Syl}(p, G/N)$. This follows because

   1. $P_H N/N = P_H/P_H \cap N$ by the 2nd isomorphism theorem, so it is a $p-$group.
   2. $P_H \subseteq P_H N \subseteq G \implies p$ doesn't divide $[G : P_H N]$, since P_H is also a Sylow $p-$group of G and thus has maximal prime power dividing $|G|$.
   3. $N \subseteq P_H N \subseteq G \implies [G/N : P_H N/N] = [G : P_H N]$

   Taken together, this says that $P_H N/N$ is a p-group and p doesn't divide $[G/N, P_H N/N]$, so it is a maximal $p-$subgroup and $P_H N/N \in \text{Syl}(p, G/N)$.

   But since $P_H N/N \leq P_N$ and $|P_H N/N| = |P_N|$, we must have $P_H N/N = P_N$ as desired.

□

Problem 15.1.11
Let G be a finite group and $H < G$ a subgroup. Let n_H be the number of subgroups of G that are conjugate to H. Show that n_H divides the order of G.

Solution:
.* Let

$$C_H = \{gHg^{-1} \;\middle|\; g \in G\}$$

be the conjugacy class of H, so $|C_H| = n_H$.
We wish to show that n_H divides $|G|$.
Claim 1:

$$n_H = [G : N_G(H)],$$

where $N_G(H) \leq G$ is the normalizer of H in G.
Note that if this claim is true, then we can apply Lagrange's theorem, which states

$$A \leq G \implies |G| = [A : G]\ |A|,$$

which in this case translates to

$$|G| = [N_G(H) : G]\ |N_G(H)| = n_H\ |N_G(H)|.$$

Since n_H divides the right-hand side, it must divide the left-hand side as well, which is precisely what we would like to show.
Proof of Claim 1:
The normalizer of H in G, written $N_G(H)$, is the largest subgroup of G containing H such that $H \trianglelefteq N_G(H)$, i.e.

$$N_G(H) = \{g \in G \;\middle|\; gHg^{-1} = H\} \leq G.$$

Now consider S, the set of left cosets of $N_G(H)$. Suppose there are k of them, so

$$[G : N_G(H)] = |S| \coloneqq k.$$

Then S can be written as

$$S = \{g_1 N_G(H),\ g_2 N_G(H),\ \cdots,\ g_k N_G(H)\}.$$

where each g_i is a distinct element of G yielding a distinct coset $g_i N_G(H)$. In particular, if $i \neq j$, then $g_i \neq g_j$, and $g_i N_G(H) \notin g_j N_G(H)$.
In particular, S acts on C_H,

$$S \curvearrowright C_H$$
$$g_i N_G(H) \curvearrowright H = g_i H g_i^{-1},$$

taking H to one of its conjugate subgroups.
So define

$$K \coloneqq \{g_i H g_i^{-1} \;\middle|\; 1 \leq i \leq k\}.$$

Note that $K \subseteq C_H$, and has at most k elements.

We claim that K has k *distinct* elements, i.e. that each g_i takes H to a *distinct* conjugate subgroup. We have

$$\begin{aligned} g_i H g_i^{-1} &= g_j H g_j^{-1} \quad \implies \\ g_j^{-1} g_i H g_i^{-1} g_j &= H \quad \implies \\ (g_j^{-1} g_i) H (g_j^{-1} g_i)^{-1} &= H \quad \implies \\ g_j^{-1} g_i &\in N_G(H) \quad \implies \\ g_i &\in g_j N_G(H) \quad \implies \\ g_i &= g_j, \end{aligned}$$

where the last line follows because we assumed that each coset contains at most one g_i.
Thus K has k distinct elements, and so

$$= k = |K| \leq |C_H| = n_H.$$

We now claim that $k \geq n_H$ as well.
Let $H' \in C_H$ be any subgroup conjugate to H, so $H' = gHg^{-1}$ for some $g \in G$. Then $g = g_i$ for some i, so $g \in g_i N_G(H)$.
Thus $g = g_i n$ for some $n \in N_G(H)$, but $n \in N_G(H) \iff nHn^{-1} = H$ by definition, and so we have

$$\begin{aligned} H' &= gHg^{-1} \\ &= (g_i n) H (g_i n)^{-1} \\ &= g_i (nHn^{-1}) g_i^{-1} \\ &= g_i H g_i^{-1} \in K. \end{aligned}$$

Since $H' \in C_H$ was an arbitrary subgroup conjugate to H, this says that $C_H \subseteq K$ and thus

$$n_H = |C_H| \leq |K| = k$$

Thus

$$[G : N_G(H)] = k = |M| = |K| = n_H,$$

which is what we wanted to show. □

Problem 15.1.12
Let $G = S_5$, the symmetric group on 5 elements. Identify all conjugacy classes of elements in G, provide a representative from each class, and prove that this list is complete.

Solution:

Claim 1: Conjugacy classes in S_n are completely determined by cycle type.
This follows because of the result on homework 1, which says that for any two cycles $\tau, \sigma \in S_n$,

we have

$$\tau(s_1\ s_2\ \cdots\ s_k)\tau^{-1} = (\tau(s^1)\ \tau(s^2)\ \cdots\ \tau(s_k)).$$

In particular, this shows that the cycle type of a single cycle is invariant under conjugation. If an element $\sigma \in S_n$ is comprised of multiple cycles, say $\sigma = \sigma_1 \cdots \sigma_\ell$, then

$$\tau(\sigma)\tau^{-1} = \tau(\sigma_1 \cdots \sigma_\ell)\tau^{-1} = (\tau\sigma_1\tau^{-1}) \cdots (\tau\sigma_\ell\tau^{-1}),$$

which shows that the entire cycle type is preserved under conjugation. So each conjugacy class has exactly one cycle type, and distinct classes have distinct cycle types, so this completely determines the conjugacy classes.

Claim 2: Cycle types in S_n are in bijective correspondence with integer partitions of n.

This follows because any integer partition of n can be used to obtain a canonical representative of a conjugacy class of S_n: if $n = a_1 + a_2 + \cdots a_n$, we simply take a cycle of length a_1 the first a_1 integers in order, a cycle of length a_2 containing the integers $a_1 + 1$ to a_2 in order, and so on.

Conversely, any permutation can be written as a product of disjoint cycles, and when the cycles for fixed points are added in, every integer between 1 and n will appear, and the sum of the lengths of all cycles must sum to n. Thus taking the cycle lengths yields an integer partition of n.

All integer partitions of 5 are given below, along with a canonical representative of the associated conjugacy class.

5	$(1\ 2\ 3\ 4\ 5)$
$4+1$	$(1\ 2\ 3\ 4)(5)$
$3+2$	$(1\ 2\ 3)(4\ 5)$
$3+1+1$	$(1\ 2\ 3)(4)(5)$
$2+2+1$	$(1\ 2)(3\ 4)(5)$
$2+1+1+1$	$(1\ 2)(3)(4)(5)$
$1+1+1+1+1$	$(1)(2)(3)(4)(5)$

15.2 Problem Set Two

15.2.1 Exercises

Problem 15.2.1 (Hungerford 2.1.9)
Let G be a finitely generated abelian group in which no element (except 0) has finite order. Show that G is a free abelian group.

Problem 15.2.2 (Hungerford 2.1.10)

1. Show that the additive group of rationals $\mathbb{Q}$ is not finitely generated.
2. Show that $\mathbb{Q}$ is not free.
3. Conclude that Exercise 9 is false if the hypothesis "finitely generated" is omitted.

Problem 15.2.3 (Hungerford 2.5.8)
Show that if every Sylow $p-$subgroup of a finite group G is normal for every prime p, then G is the direct product of its Sylow subgroups.

Problem 15.2.4 (Hungerford 2.6.4)
What is the center of the quaternion group Q_8? Show that $Q_8/Z(Q_8)$ is abelian.

Problem 15.2.5 (Hungerford 2.6.9)
Classify up to isomorphism all groups of order 18. Do the same for orders 20 and 30.

Problem 15.2.6 (Hungerford 1.9.1)
Show that every non-identity element in a free group F has infinite order.

Problem 15.2.7 (Hungerford 1.9.3)
Let F be a free group and for a fixed integer n, let H_n be the subgroup generated by the set $\{x^n \mid x \in F\}$. Show that $H_n \trianglelefteq F$.

15.2.2 Qual Problems

Problem 15.2.8
List all groups of order 14 up to isomorphism.

Problem 15.2.9
Let G be a group of order p^3 for some prime p. Show that either G is abelian, or $|Z(G)| = p$.

Problem 15.2.10
Let p, q be distinct primes, and let k denote the smallest positive integer such that p divides $q^k - 1$. Show that no group of order pq^k is simple.

Problem 15.2.11
Show that S_4 is a solvable, nonabelian group.

15.3 Problem Set Three

15.3.1 Exercises

Problem 15.3.1 (Hungerford 2.7.10)
Show that S_n is solvable for$n \leq 4$ but S_3 and S_4 are not nilpotent.

Problem 15.3.2 (Hungerford 2.8.3)
Show that if N is a simple normal subgroup of a group G and G/N has a composition series, then G has a composition series.

Problem 15.3.3 (Hungerford 2.8.9)
Show that any group of order p^2q(for primes p, q) is solvable.

Problem 15.3.4 (Hungerford 5.1.1)
Let F/K be a field extension. Show that

1. $[F : K] = 1$ iff $F = K$.
2. If $[F : K]$ is prime, then there are no intermediate fields between F and K.
3. If $u \in F$ has degree n over K, then n divides $[F : K]$.

Problem 15.3.5 (Hungerford 5.1.8)
Show that if $u \in F$ is algebraic of odd degree over K, then so is u^2, and moreover $K(u) = K(u^2)$.

Problem 15.3.6 (Hungerford 5.1.14)

1. If $F = \mathbb{Q}(\sqrt{2}, \sqrt{3})$, compute $[F : \mathbb{Q}]$ and find a basis of $F/\mathbb{Q}$.
2. Do the same for $\mathbb{Q}(i, \sqrt{3}, \zeta_3)$ where ζ_3 is a complex third root of 1.

Problem 15.3.7 (Hungerford 5.1.16)
Show that in $\mathbb{C}$, the fields $\mathbb{Q}(i) \cong \mathbb{Q}(\sqrt{2})$ as vector spaces, but not as fields.

15.3.2 Qual Problems

Problem 15.3.8
Let R and S be commutative rings with multiplicative identity.

1. Prove that when R is a field, every non-zero ring homomorphism $\varphi : R \to S$ is injective.
2. Does (a) still hold if we only assume that R is a domain? If so, prove it, and if not

provide a counterexample.

Problem 15.3.9
Determine for which integers the ring $\mathbb{Z}/n\mathbb{Z}$ is a direct sum of fields. Carefully prove your answer.

Problem 15.3.10
Suppose that R is a commutative ring. Show that an element $r \in R$ is not invertible iff it is contained in a maximal ideal.

Problem 15.3.11

1. Give the definition that a group G must satisfy the be solvable.
2. Show that every group G of order 36 is solvable.

Hint: You may assume that S^4 is solvable.

15.4 Problem Set Four

15.4.1 Exercises

Problem 15.4.1 (Hungerford 5.3.7)
If F is algebraically closed and$E$ is the set of all elements in F that are algebraic over a field K, then E is an algebraic closure of K.

Problem 15.4.2 (Hungerford 5.3.8)
Show that no finite field is algebraically closed.
Hint: if $K = \{a_i\}_{i=0}^n$, consider

$$f(x) = a_1 + \prod_{i=0}^{n}(x - a_i) \in K[x]$$

where $a_1 \neq 0$.

Problem 15.4.3 (Hungerford 5.5.2)
Show that if $p \in \mathbb{Z}$ is prime, then $a^p = a$ for all $a \in \mathbb{Z}_p$, or equivalently $c^p \equiv c \pmod{p}$ for all $c \in \mathbb{Z}$.

Problem 15.4.4 (Hungerford 5.5.3)
Show that if $|K| = p^n$, then every element of K has a unique pth root in K.

Problem 15.4.5 (Hungerford 5.5.10)
Show that every element in a finite field can be written as the sum of two squares.

Problem 15.4.6 (Hungerford 5.6.1)
Let F/K be a field extension. Let $\text{char}K = p \neq 0$ and let $n \geq 1$ be an integer such that $(p, n) = 1$. If $v \in F$ and $nv \in K$, then $v \in K$.

Problem 15.4.7 (Hungerford 5.6.8)
If $\text{char}K = p \neq 0$ and $[F : K]$ is finite and not divisible by p, then F is separable over K.

15.4.2 Qual Problems

Problem 15.4.8
Suppose that α is a root in $\mathbb{C}$ of $P(x) = x^{17} - 2$. How many field homomorphisms are there from $\mathbb{Q}(\alpha)$ to:

1. $\mathbb{C}$,
2. $\mathbb{R}$,
3. $\overline{\mathbb{Q}}$, the algebraic closure of $\mathbb{Q}$?

Problem 15.4.9
Let C/F be an algebraic field extension. Prove that the following are equivalent:

1. Every non-constant polynomial $f \in F[x]$ factors into linear factors over $C[x]$.
2. For every (not necessarily finite) algebraic extension E/F, there is a ring homomorphism $\alpha : E \to C$ such that $\alpha \Big|_F$ is the identity on F.

Hint: use Zorn's Lemma.

Problem 15.4.10
Let R be a commutative ring containing a field k, and suppose that $\dim_k R < \infty$. Let $\alpha \in R$.

1. Show that there exist $n \in \mathbb{N}$ and $\{c_0, c_1, \cdots c_{n-1}\} \subseteq k$ such that
$$a^n + c_{n-1}a^{n-1} + \cdots + c_1 a + c_0 = 0.$$
2. Suppose that (a) holds and show that if $c_0 \neq 0$ then a is a unit in R.
3. Suppose that (a) holds and show that if a is not a zero divisor in R, then a is invertible.

15.5 Problem Set Five

15.5.1 Exercises

Problem 15.5.1 (Hungerford 5.3.5)
Show that if $f \in K[x]$ has degree n and F is a splitting field of f over K, the $[F : K]$ divides $n!$.

Problem 15.5.2 (Hungerford 5.3.12)
Let E be an intermediate field extension in $K \leq E \leq F$.

1. Show that if $u \in F$ is separable over over K, then u is separable over E.
2. Show that if F is separable over K, then F is separable over E and E is separable over K.

Problem 15.5.3 (Hungerford 5.3.13)
Show that if $[F : K] < \infty$, then the following conditions are equivalent:

1. F is Galois over K
2. F is separable over K and F is a splitting field of some polynomial $f \in K[x]$.
3. F is a splitting field over K of some polynomial $f \in K[x]$ whose irreducible factors are separable.

Problem 15.5.4 (Hungerford 5.4.1)
Suppose that $f \in K[x]$ splits in$F$ as

$$f = \prod_{i=1}^{k} (x - u_i)^{n_i}$$

with the u_i distinct and each $n_i \geq 1$. Let

$$g(x) = \prod_{i=1}^{k} (x - u_i) = \sum_{i=1}^{k} v_i x^i$$

and let $E = K(\{v_i\}_{i=1}^{k})$. Then show that the following hold:

1. F is a splitting field of g over E.
2. F is Galois over E.
3. $\mathrm{Aut}_E(F) = \mathrm{Aut}_K(F)$.

Problem 15.5.5 (Hungerford 5.4.10 a/g/h)
Determine the Galois groups of the following polynomials over the corresponding fields:

1. $x^4 - 5$ over $\mathbb{Q}, \mathbb{Q}(\sqrt{5}), \mathbb{Q}(i\sqrt{5})$.
2. $x^3 - 2$ over $\mathbb{Q}$.
3. $(x^3 - 2)(x^2 - 5)$ over $\mathbb{Q}$.

Problem 15.5.6 (Hungerford 5.6.11)
If $f \in K[x]$ is irreducible of degree $m > 0$ and $\text{char}(K)$ does not divide m, then f is separable.

15.5.2 Qual Problems

Problem 15.5.7
Let E/F be a Galois field extension, and let K/F be an intermediate field of E/F. Show that K is normal over F iff $\text{Gal}(E/K) \trianglelefteq \text{Gal}(E/F)$.

Problem 15.5.8
Let $F \subset L$ be fields such that L/F is a Galois field extension with Galois group equal to $D_8 = \left\langle \sigma, \tau \;\middle|\; \sigma^4 = \tau^2 = 1,\ \sigma\tau = \tau\sigma^3 \right\rangle$. Show that there are fields $F \subset E \subset K \subset L$ such that E/F and K/E are Galois field extensions, but K/F is not Galois.

Problem 15.5.9
Let $f(x) = x^3 - 7$.

1. Let K be the splitting field for f over $\mathbb{Q}$. Describe the Galois group of $K/\mathbb{Q}$ and the intermediate fields between $\mathbb{Q}$ and K. Which intermediate fields are not Galois over $\mathbb{Q}$?
2. Let L be the splitting field for f over $\mathbb{R}$. What is the Galois group $L/\mathbb{R}$?
3. Let M be the splitting field for f over $\mathbb{F}_{13}$, the field with 13 elements. What is the Galois group of $M/\mathbb{F}_{13}$?

15.6 Problem Set Six

15.6.1 Exercises

Problem 15.6.1 (Hungerford 5.4.11)
Determine all subgroups of the Galois group and all intermediate fields of the splitting (over $\mathbb{Q}$) of the polynomial $(x^3-2)(x^2-3) \in \mathbb{Q}[x]$.

Problem 15.6.2 (Hungerford 5.4.12)
Let K be a subfield of$\mathbb{R}$ and let $f \in K[x]$ be an irreducible quartic. If f has exactly 2 real roots, the Galois group of f is either S_4 or D_4.

Problem 15.6.3 (Hungerford 5.8.3)
Let φ be the Euler function.

1. $\varphi(n)$ is even for $n > 2$.
2. find all $n > 0$ such that $\varphi(n) = 2$.

Problem 15.6.4 (Hungerford 5.8.9)
If $n > 2$ and ζ is a primitive nth root of unity over $\mathbb{Q}$, then $[\mathbb{Q}(\zeta + \zeta^{-1}) : \mathbb{Q}] = \varphi(n)/2$.

Problem 15.6.5 (Hungerford 5.9.1)
If F is a radical extension field of K and E is an intermediate field, then F is a radical extension of E.

Problem 15.6.6 (Hungerford 5.9.3)
Let K be a field, $f \in K[x]$ an irreducible polynomial of degree $n \geq 5$ and F a splitting field of f over K. Assume that $Aut_k(F) \simeq S_n$. Let u be a root of f in F. Then,

1. $K(u)$ is not Galois over K; $[K(u) : K] = n$ and $Aut_K(K(u)) = 1$ (and hence solvable).
2. Every normal closure over K that contains u also contains an isomorphic copy of F.
3. There is no radical extension field E of K such that $K \subset K(u) \subset E$.

15.6.2 Qual Problems

Problem 15.6.7

1. Let K be a field. State the main theorem of Galois theory for a finite field extension L/K

2. Let $\zeta_{43} := e^{2\pi i/43}$. Describe the group of all field automorphisms $\sigma : \mathbb{Q}(\zeta_{43}) \to \mathbb{Q}(\zeta_{43})$.

3. How many proper subfields are there in the field $\mathbb{Q}(\zeta_{43})$?

Problem 15.6.8
Let F be a field and let $f(x) \in F[x]$.

1. Define what is a splitting field of $f(x)$ over F.

2. Let F be a finite field with q elements. Let E/F be a finite extension of degree $n > 0$. Exhibit an explicit polynomial $g(x) \in F[x]$ such that E/F is a splitting of $g(x)$ over F. Fully justify your answer.

3. Show that the extension E/F in (2) is a Galois extension.

Problem 15.6.9
Let $K \subset L \subset M$ be a tower of finite degree field extensions. In each of the following parts, either prove the assertion or give a counterexample (with justification).

1. If M/K is Galois, then L/K is Galois

2. If M/K is Galois, then M/L is Galois.

15.7 Problem Set Seven

15.7.1 Exercises

Problem 15.7.1 (Hungerford 4.1.3)
Let I be a left ideal of a ring$R$, and let A be an $R-$module.

1. Show that if S is a nonempty subset of A, then

$$IS := \left\{ \sum_{i=1}^{n} r_i a_i \;\middle|\; n \in \mathbb{N}^*; r_i \in I; a_i \in S \right\}$$

is a submodule of A.

Note that if $S = \{a\}$, then $IS = Ia = \{ra \mid r \in I\}$.

2. If I is a two-sided ideal, then A/IA is an R/I module with the action of R/I given by

$$(r + I)(a + IA) = ra + IA.$$

Problem 15.7.2 (Hungerford 4.1.5)
If R has an identity, then a nonzero unitary R-module is **simple** if its only submodules are 0 and A.

1. Show that every simple $R-$module is cyclic.
2. If A is simple, every $R-$module endomorphism is either the zero map or an isomorphism.

Problem 15.7.3 (Hungerford 4.1.7)

1. Show that if A, B are R-modules, then the set $\mathrm{Hom}_R(A, B)$ is all R-module homomorphisms $A \to B$ is an abelian group with $f + g$ given on $a \in A$ by

$$(f + g)(a) \coloneqq f(a) + g(a) \in B.$$

   Also show that the identity element is the zero map.

2. Show that $\mathrm{Hom}_R(A, A)$ is a ring with identity, where multiplication is given by composition of functions.

   Note that $\mathrm{Hom}_R(A, A)$ *is called the* ***endomorphism ring*** *of* A.

3. Show that A is a left $\mathrm{Hom}_R(A, A)$-module with an action defined by

$$a \in A, f \in \mathrm{Hom}_R(A, A) \implies f \curvearrowright a \coloneqq f(a).$$

Problem 15.7.4 (Hungerford 4.1.12)
Let the following be a commutative diagram of R-modules and R-module homomorphisms with exact rows:
Prove the following:

1. If α_1 is an epimorphisms and α_2, α_4 are monomorphisms then α_3 is a monomorphism.
2. If α_5 is a monomorphism and α_2, α_4 are epimorphisms then α_3 is an epimorphism.

Problem 15.7.5 (Hungerford 4.2.4)
Let R be a principal ideal domain,A a unitary left R-module, and $p \in R$ a prime (and thus irreducible) element. Define

$$pA \coloneqq \{pa \mid a \in A\}$$

$$A[p] \coloneqq \{a \in A \mid pa = 0\}.$$

Show the following:

1. $R/(p)$ is a field.
2. pA and $A[p]$ are submodules of A.
3. A/pA is a vector space over $R/(p)$, with

$$(r + (p))(a + pA) = ra + pA.$$

4. $A[p]$ is a vector space over $R/(p)$ with

$$(r + (p))a = ra.$$

Problem 15.7.6 (Hungerford 4.2.8)
If V is a finite dimensional vector space and

$$V^m \coloneqq V \oplus V \oplus \cdots \oplus V \quad (m \text{ summands}),$$

then for each $m \geq 1$, V^m is finite dimensional and $\dim V^m = m(\dim V)$.

Problem 15.7.7 (Hungerford 4.2.9)
If F_1, F_2 are free modules of a ring with the invariant dimension property, then

$$\text{rank}(F_1 \oplus F_2) = \text{rank}F_1 + \text{rank}F_2.$$

15.7.2 Qual Problems

Problem 15.7.8
Let F be a field and let $f(x) \in F[x]$.

1. State the definition of a splitting field of $f(x)$ over F.
2. Let F be a finite field with q elements. Let E/F be a finite extension of degree $n > 0$. Exhibit an explicit polynomial $g(x) \in F[x]$ such that E/F is a splitting field of g over F. Fully justify your answer.
3. Show that the extension in (b) is a Galois extension.

Problem 15.7.9
Let R be a commutative ring and let M be an R-module. Recall that for $\mu \in M$, the *annihilator* of μ is the set

$$\text{Ann}(\mu) = \{r \in R \,\big|\, r\mu = 0\}.$$

Suppose that I is an ideal in R which is maximal with respect to the property there exists a nonzero element $\mu \in M$ such that $I = \text{Ann}(\mu)$.
Prove that I is a *prime* ideal in R.

Problem 15.7.10
Suppose that R is a principal ideal domain and $I \trianglelefteq R$ is an ideal. If $a \in I$ is an irreducible element, show that $I = Ra$.

15.8 Problem Set Eight

15.8.1 Exercises

Problem 15.8.1 (Hungerford 4.4.1)
Show the following:

1. For any abelian group A and any positive integer m,
$$\mathrm{Hom}(\mathbb{Z}_m, A) \cong A[m] \coloneqq \{a \in A \;\middle|\; ma = 0\}.$$
2. $\mathrm{Hom}(\mathbb{Z}_m, \mathbb{Z}_n) \cong \mathbb{Z}_{\gcd(m,n)}$.
3. As a $\mathbb{Z}-$module, $\mathbb{Z}_m^* = 0$.
4. For each $k \geq 1$, $\mathbb{Z}_m$ is a $\mathbb{Z}_{mk}-$module, and as a $\mathbb{Z}_{mk}$ module, $\mathbb{Z}_m^* \cong \mathbb{Z}_m$.

Problem 15.8.2 (Hungerford 4.4.3)
Let$\pi : \mathbb{Z} \to \mathbb{Z}_2$ be the canonical epimorphism. Show that the induced map $\overline{\pi} : \mathrm{Hom}(\mathbb{Z}_2, \mathbb{Z}) \to \mathrm{Hom}(\mathbb{Z}_2, \mathbb{Z}_2)$ is the zero map. Conclude that $\overline{\pi}$ is not an epimorphism.

Problem 15.8.3 (Hungerford 4.4.5)
Let R be a unital ring, show that there is a ring homomorphism $\mathrm{Hom}_R(R, R) \to R^{op}$ where Hom_R denotes left $R-$module homomorphisms. Conclude that if R is commutative, then there is a ring isomorphism $\mathrm{Hom}_R(R, R) \cong R$.

Problem 15.8.4 (Hungerford 4.4.9)
Show that for any homomorphism$f : A \to B$ of left $R-$modules the following diagram is commutative:
where θ_A, θ_B are as in Theorem 4.12 and f^* is the map induced on $A^{**} \coloneqq \mathrm{Hom}_R(\mathrm{Hom}(A, R), R)$ by the map
$$\overline{f} : \mathrm{Hom}(B, R) \to \mathrm{Hom}_R(A, R).$$

Problem 15.8.5 (Hungerford 4.6.2)
Show that every free module over a unital integral domain is torsion-free. Show that the converse is false.

Problem 15.8.6 (Hungerford 4.6.3)
Let A be a cyclic $R-$module of order $r \in R$.

1. Show that if s is relatively prime to r, then $sA = A$ and $A[s] = 0$.
2. If s divides r, so $sk = r$, then $sA \cong R/(k)$ and $A[s] \cong R/(s)$.

Problem 15.8.7 (Hungerford 4.6.6)
Let A, B be cyclic modules over $R$of nonzero orders r, s respectively, where r is *not* relatively prime to s. Show that the invariant factors of $A \oplus B$ are $\gcd(r, s)$ and $\operatorname{lcm}(r, s)$.

15.8.2 Qual Problems

Problem 15.8.8
Let R be a PID. Let $n > 0$ and $A \in M_n(R)$ be a square $n \times n$ matrix with coefficients in R. Consider the R-module $M \coloneqq R^n/\operatorname{im}(A)$.

1. Give a necessary and sufficient condition for M to be a torsion module (i.e. every nonzero element is torsion). Justify your answer.
2. Let F be a field and now let $R \coloneqq F[x]$. Give an example of an integer $n > 0$ and an $n \times n$ square matrix $A \in M_n(R)$ such that $M \coloneqq R^n/\operatorname{im}(A)$ is isomorphic as an $R-$module to $R \times F$.

Problem 15.8.9

1. State the structure theorem for finitely generated modules over a PID.
2. Find the decomposition of the $\mathbb{Z}-$module M generated by w, x, y, z satisfying the relations

$$\begin{aligned} 3w + 12y + 3x + 6z &= 0 \\ 6y &= 0 \\ -3w - 3x + 6y &= 0. \end{aligned}$$

Problem 15.8.10
Let R be a commutative ring and M an $R-$module.

1. Define what a torsion element of M is .
2. Given an example of a ring R and a cyclic $R-$module M such that M is infinite and M contains a nontrivial torsion element m. Justify why m is torsion.
3. Show that if R is a domain, then the subset of elements of M that are torsion is an $R-$submodule of M. Clearly show where the hypothesis that R is a domain is used.

15.9 Problem Set Nine

15.9.1 Exercises

Problem 15.9.1 (Hungerford 7.1.3)

1. Show that the center of the ring $M_n(R)$ consists of matrices of the form rI_n where r is in the center of R.

   Hint: Every such matrix must commute with ϵ_{ij}, the matrix with 1_R in the i, j position and zeros elsewhere.

2. Show that $Z(M_n(R)) \cong Z(R)$.

Problem 15.9.2 (Hungerford 7.1.5)

1. Show that if A, B are (skew)-symmetric then $A + B$ is (skew)-symmetric.

2. Let R be commutative. Show that if A, B are symmetric, then AB is symmetric $\iff$ $AB = BA$. Also show that for any matrix $B \in M_n(R)$, both BB^t and $B + B^t$ are always symmetric, and $B - B^t$ is always skew-symmetric.

Problem 15.9.3 (Hungerford 7.1.7)
Show that similarity is an equivalence relation on $M_n(R)$, and *equivalence* is an equivalence relation on $M_{m\times n}(R)$.

Problem 15.9.4 (Hungerford 7.2.2)
Show that an $n \times m$ matrix Aover a division ring D has an $m \times n$ left inverse B (so $BA = I_m$) $\iff$ rank$A = m$. Similarly, show A has a right $m \times n$ inverse $\iff$ rank$A = n$.

Problem 15.9.5 (Hungerford 7.2.4)

1. Show that a system of linear equations

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1m}x_m &= b_1 \\ &\vdots \\ a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nm}x_m &= b_n \end{aligned}$$

   has a simultaneous solution $\iff$ the corresponding matrix equation $AX = B$ has a solution, where $A = (a_{ij}), X = [x_1, \cdots, x_m]^t$, and $B = [b_1, \cdots, b_n]^t$.

2. If A_1, B_1 are matrices obtained from A, B respectively by performing the same sequence of elementary **row** operations, then X is a solution of $AX = B$ $\iff$ X is a solution of $A_1X = B_1$.

3. Let C be the $n \times (m+1)$ matrix given by

$$C = \begin{pmatrix} a_{11} & \cdots & a_{1m} & b_1 \\ & & & \\ . & & & \\ a_{n1} & \cdots & a_{nm} & b_n \end{pmatrix}.$$

   Then $AX = B$ has a solution $\iff$ rank$A$ = rank$C$ and the solution is unique $\iff$ rank$(A) = m$.

Hint: use part 2.

4. If $B = 0$, so the system $AX = B$ is homogeneous, then it has a nontrivial solution $\iff$ rank$A < m$ and in particular $n < m$.

Problem 15.9.6 (Hungerford 7.2.5)
Let R be a PID. For each positive integer r and sequence of nonzero ideals $I_1 \supset I_2 \supset \cdots \supset I_r$, choose a sequence $d_i \in R$ such that $(d_i) = I_i$ and $d_i \mid d_{i+1}$.
For a given pair of positive integers n, m, let S be the set of all $n \times m$ matrices of the form $\begin{pmatrix} L_r & 0 \\ 0 & 0 \end{pmatrix}$ where $r = 1, 2, \cdots, \min(m, n)$ and L_r is a diagonal $r \times r$ matrix with main diagonal d_i.
Show that S is a set of canonical forms under equivalence for the set of all $n \times m$ matrices over R.

15.9.2 Qual Problems

Problem 15.9.7
Let R be a commutative ring.

1. Say what it means for R to be a unique factorization domain (UFD).
2. Say what it means for R to be a principal ideal domain (PID)
3. Give an example of a UFD that is not a PID. Prove that it is not a PID.

Problem 15.9.8
Let A be an $n \times n$ matrix over a field F such that A is diagonalizable. Prove that the following are equivalent:

1. There is a vector $v \in F^n$ such that $v, Av, \cdots A^{n-1}v$ is a basis for F^n.
2. The eigenvalues of A are distinct.

Problem 15.9.9
Let $x, y \in \mathbb{C}$ and consider the matrix

$$M = \begin{bmatrix} 1 & 0 & x \\ 0 & 1 & 0 \\ y & 0 & 1 \end{bmatrix}$$

1. Show that $[0, 1, 0]^t$ is an eigenvector of M.
2. Compute the rank of M as a function of x and y.
3. Find all values of x and y for which M is diagonalizable.

15.10 Problem Set Ten

15.10.1 Exercises

Problem 15.10.1 (Hungerford 7.3.1)
Let B be an R-module. Show that if $r + r \neq 0$ for all $r \neq 0 \in R$, then an n-linear form $B^n \to R$ is alternating $\iff$ it is skew-symmetric.

Problem 15.10.2 (Hungerford 7.3.5)
If R is a field and $A, B \in M_n(R)$ are invertible then the matrix $A + rB$ is invertible for all but a finite number of $r \in R$.

Problem 15.10.3 (Hungerford 7.4.4)
Show that if q is the minimal polynomial of a linear transformation $\varphi : E \to E$ with $\dim_k E = n$ then $\deg q \leq n$.

Problem 15.10.4 (Hungerford 7.4.8).)
Show that $A \in M_n(K)$ is similar to a diagonal matrix $\iff$ the elementary divisors of A are all linear.

Problem 15.10.5 (Hungerford 7.4.10)
Find all possible rational canonical forms for a matrix $A \in M_n(\mathbb{Q})$ such that

1. A is 6×6 with minimal polynomial $q(x) = (x-2)^2(x+3)$.
2. A is 7×7 with $q(x) = (x^2+1)(x-7)$.

Also find all such forms when $A \in M_n(\mathbb{C})$ instead, and find all possible Jordan Canonical Forms over $\mathbb{C}$.

Problem 15.10.6 (Hungerford 7.5.2)
Show that if φ is an endomorphism of a free k-module E of finite rank, then $p_\varphi(\varphi) = 0$.
Hint: If A is the matrix of φ and $B = xI_n - A$ then

$$B^a B = |B| I_n = p_\varphi I_n \in M_n(k[x]).$$

If E is a $k[x]$-module with structure induced by φ, and ψ is the $k[x]$-module endomorphism $E \to E$ with matrix given by B, then

$$\psi(u) = xu - \varphi(u) = \varphi(u) - \varphi(u) = 0 \qquad \forall u \in E.$$

Problem 15.10.7 (Hungerford 7.5.7)

1. Let φ, ψ be endomorphisms of a finite-dimensional vector space E such that $\varphi\psi = \psi\varphi$. Show that if E has a basis of eigenvectors of ψ, then it has a basis of eigenvectors for both ψ and φ simultaneously.
2. Interpret the previous part as a statement about matrices similar to a diagonal matrix.

15.10.2 Qual Problems

Problem 15.10.8
Let $M \in M_5(R)$ be a 5×5 square matrix with real coefficients defining a linear map $L : \mathbb{R}^5 \to \mathbb{R}^5$. Assume that when considered as an element of $M_5(\mathbb{C})$, then the scalars $0, 1+i, 1+2i$ are eigenvalues of M.

1. Show that the associated linear map L is neither injective nor surjective.
2. Compute the characteristic polynomial and minimal polynomial of M.
3. How many fixed points can L have?
   (That is, how many solutions are there to the equation $L(v) = v$ with $v \in \mathbb{R}^5$?)

Problem 15.10.9
Let n be a positive integer and let B denote the $n \times n$ matrix over $\mathbb{C}$ such that every entry is 1. Find the Jordan normal form of B.

Problem 15.10.10
Suppose that V is a 6-dimensional vector space and that T is a linear transformation on V such that $T^6 = 0$ and $T^5 \neq 0$.

1. Find a matrix for T in Jordan Canonical form.
2. Show that if S, T are linear transformations on a 6-dimensional vector space V which both satisfy $T^6 = S^6 = 0$ and $T^5, S^5 \neq 0$, then there exists a linear transformation A from V to itself such that $ATA^{-1} = S$.

Bibliography

[1] David Steven. Dummit and Richard M. Foote. *Abstract algebra*. John Wiley and Sons, 2004.

[2] Kenneth Hoffman and Ray Kunze. *Linear Algebra*. Prentice Hall, 1981.

[3] Thomas W. Hungerford. *Algebra*. Springer, 2008.

[4] Roy Smith. *Algebra Notes by Roy Smith*. URL: https://www.math.uga.edu/directory/people/roy-smith.